

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OPANA® ER safely and effectively. See full prescribing information for OPANA® ER.

OPANA® ER (oxymorphone hydrochloride) extended-release tablets, for oral use, CII

Initial U.S. Approval: 1959

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; and INTERACTION WITH ALCOHOL; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES AND OTHER CNS DEPRESSANTS.

See full prescribing information for complete boxed warning.

- OPANA ER exposes users to risks of addiction, abuse, and misuse, which can lead to overdose and death. Assess patient's risk before prescribing, and monitor regularly these behaviors and conditions. (5.1)
- To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a Risk Evaluation and Mitigation Strategy (REMS) for these products. (5.2)
- Serious life-threatening or fatal respiratory depression may occur. Monitor closely, especially upon initiation or following a dose increase. Instruct patients to swallow OPANA ER tablets whole to avoid exposure to a potentially fatal dose of oxymorphone. (5.3)
- Accidental ingestion of OPANA ER, especially by children, can result in fatal overdose of oxymorphone. (5.3)
- Prolonged use of OPANA ER during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available. (5.4)
- Instruct patients not to consume alcohol or any product containing alcohol while taking OPANA ER because co-ingestion can result in fatal plasma oxymorphone levels. (5.5)
- Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing for use in patients for whom alternative treatment options are inadequate; limit dosages and durations to the minimum required; and follow patients for signs and symptoms of respiratory depression and sedation. (5.5, 7)

RECENT MAJOR CHANGES

Dosage and Administration (2.4)	10/2019
Warnings and Precautions (5.3, 5.14)	10/2019

INDICATIONS AND USAGE

OPANA ER is an opioid agonist indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. (1)

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve OPANA ER for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain. (1)
- OPANA ER is not indicated as an as-needed (prn) analgesic. (1)

DOSAGE AND ADMINISTRATION

- To be prescribed only by healthcare providers knowledgeable in use of potent opioids for management of chronic pain. (2.2)
- Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals (2.2).
- Individualize dosing based on the severity of pain, patient response, prior analgesic experience, and risk factors for addiction, abuse, and misuse.

Administer on an empty stomach, at least 1 hour prior to or 2 hours after eating. (2.2)

- For opioid-naïve and opioid non-tolerant patients, initiate treatment with 5 mg tablets orally every 12 hours. (2.2)
- To convert to OPANA ER from another opioid, use available conversion factors to obtain estimated dose. (2.2)
- Dose can be increased every 3 to 7 days, using increments of 5 to 10 mg every 12 hours (i.e., 10 to 20 mg per day). (2.3)
- Do not abruptly discontinue OPANA ER in a physically dependent patient because rapid discontinuation of opioid analgesics has resulted in serious withdrawal symptoms, uncontrolled pain, and suicide. (2.4, 5.14)
- **Mild Hepatic Impairment:** For opioid-naïve patients, initiate treatment with 5 mg and titrate slowly. For patients on prior opioid therapy, reduce starting dose by 50% and titrate slowly. Monitor for signs of respiratory and central nervous system depression. (2.5)
- **Renal Impairment:** For opioid-naïve patients, initiate treatment with 5 mg and titrate slowly. For patients on prior opioid therapy, reduce starting dose by 50% and titrate slowly. Monitor for signs of respiratory and central nervous system depression. (2.6)
- **Geriatric Patients:** Initiate dosing with 5 mg, titrate slowly, and monitor for signs of respiratory and central nervous system depression. (2.7)

DOSAGE FORMS AND STRENGTHS

Extended-release tablets: 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, and 40 mg

CONTRAINDICATIONS

- Significant respiratory depression (4)
- Acute or severe bronchial asthma in an unmonitored setting or in absence of resuscitative equipment. (4)
- Hypersensitivity to oxymorphone (4)
- Moderate or severe hepatic impairment (4)
- Known or suspected gastrointestinal obstruction, including paralytic ileus (4)

WARNINGS AND PRECAUTIONS

- **Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly Cachectic or Debilitated Patients:** Monitor closely particularly during initiation and titration. (5.6)
- **Anaphylaxis, Angioedema, and Other Hypersensitivity Reactions:** If symptoms occur, stop administration immediately, discontinue permanently, and do not rechallenge with any other oxymorphone formulation. (5.7)
- **Adrenal Insufficiency:** If diagnosed, treat with physiologic replacement of corticosteroids, and wean patient off of the opioid. (5.8)
- **Severe Hypotension:** Monitor during dose initiation and titration. Avoid use of OPANA ER in patients with circulatory shock. (5.10)
- **Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury or Impaired Consciousness:** Monitor for sedation and respiratory depression. Avoid use of OPANA ER in patients with impaired consciousness or coma. (5.11)

ADVERSE REACTIONS

Adverse reactions in ≥2% of patients in placebo-controlled trials: nausea, constipation, dizziness, somnolence, vomiting, pruritus, headache, sweating increased, dry mouth, sedation, diarrhea, insomnia, fatigue, appetite decreased, and abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Endo Pharmaceuticals Inc. at 1-800-462-3636 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- **Serotonergic Drugs:** Concomitant use may result in serotonin syndrome. Discontinue OPANA ER if serotonin syndrome is suspected. (7)
- **Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics:** Avoid use with OPANA ER because they may reduce analgesic effect of OPANA ER or precipitate withdrawal symptoms. (7)
- **Monoamine Oxidase Inhibitors (MAOIs):** Can potentiate the effects of oxymorphone. Avoid concomitant use in patients receiving MAOIs or within 14 days of stopping treatment with an MAOI. (7)

USE IN SPECIFIC POPULATIONS

- Pregnancy: May cause fetal harm. (8.1)
- Lactation: Not recommended. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: 10/2019

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; INTERACTION WITH ALCOHOL; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

- 1 INDICATIONS AND USAGE**
- 2 DOSAGE AND ADMINISTRATION**
 - 2.1 Important Dosage and Administration Instructions
 - 2.2 Initial Dosing
 - 2.3 Titration and Maintenance of Therapy
 - 2.4 Safe Reduction or Discontinuation of OPANA ER
 - 2.5 Dosage Modifications in Patients with Hepatic Impairment
 - 2.6 Dosage Modifications in Patients with Renal Impairment
 - 2.7 Dosage Modifications in Geriatric Patients
- 3 DOSAGE FORMS AND STRENGTHS**
- 4 CONTRAINDICATIONS**
- 5 WARNINGS AND PRECAUTIONS**
 - 5.1 Addiction, Abuse, and Misuse
 - 5.2 Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)
 - 5.3 Life Threatening Respiratory Depression
 - 5.4 Neonatal Opioid Withdrawal Syndrome
 - 5.5 Risks from Concomitant Use with Benzodiazepines or Other CNS Depressants
 - 5.6 Risk of Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients
 - 5.7 Anaphylaxis, Angioedema, and Other Hypersensitivity Reactions
 - 5.8 Adrenal Insufficiency
 - 5.9 Use in Patients with Hepatic Impairment
 - 5.10 Severe Hypotension

- 5.11 Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness
- 5.12 Risks of Use in Patients with Gastrointestinal Conditions
- 5.13 Increased Risk of Seizures in Patients with Seizure Disorders
- 5.14 Withdrawal
- 5.15 Risks of Driving and Operating Machinery
- 6 ADVERSE REACTIONS**
 - 6.1 Clinical Trial Experience
 - 6.2 Post-marketing Experience
- 7 DRUG INTERACTIONS**
- 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.2 Lactation
 - 8.3 Females and Males Reproductive Potential
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Hepatic Impairment
 - 8.7 Renal Impairment
- 9 DRUG ABUSE AND DEPENDENCE**
 - 9.1 Controlled Substance
 - 9.2 Abuse
 - 9.3 Dependence
- 10 OVERDOSAGE**
- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION**

***Sections or subsections omitted from the full prescribing information are not listed.**

FULL PRESCRIBING INFORMATION

WARNING: ADDICTION, ABUSE, AND MISUSE; LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; INTERACTION WITH ALCOHOL; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

Addiction, Abuse, and Misuse

OPANA ER exposes patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing OPANA ER, and monitor all patients regularly for the development of these behaviors and conditions [see *Warnings and Precautions (5.1)*].

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS):

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a REMS for these products [see *Warnings and Precautions (5.2)*]. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to

- complete a REMS-compliant education program,
- counsel patients and/or their caregivers, with every prescription, on safe use, serious risks, storage, and disposal of these products,
- emphasize to patients and their caregivers the importance of reading the Medication Guide every time it is provided by their pharmacist, and
- consider other tools to improve patient, household, and community safety.

Life-threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of OPANA ER. Monitor for respiratory depression, especially during initiation of OPANA ER or following a dose increase. Instruct patients to swallow OPANA ER tablets whole; crushing, chewing, or dissolving OPANA ER tablets can cause rapid release and absorption of a potentially fatal dose of oxymorphone [see *Warnings and Precautions (5.3)*].

Accidental Ingestion

Accidental ingestion of even one dose of OPANA ER, especially by children, can result in a fatal overdose of oxymorphone [see *Warnings and Precautions (5.3)*].

Neonatal Opioid Withdrawal Syndrome

Prolonged use of OPANA ER during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see *Warnings and Precautions (5.4)*].

Interaction with Alcohol

Instruct patients not to consume alcoholic beverages or use prescription or non-prescription products that contain alcohol while taking OPANA ER. The co-ingestion of alcohol with OPANA ER may result in increased plasma levels and a potentially fatal overdose of oxymorphone [see *Warnings and Precautions (5.5)*].

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death [see *Warnings and Precautions 5.5, Drug Interactions (7)*].

- Reserve concomitant prescribing of OPANA ER and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.
- Limit dosages and durations to the minimum required.
- Follow patients for signs and symptoms of respiratory depression and sedation.

1 INDICATIONS AND USAGE

OPANA ER is indicated for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Limitations of Usage

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations [see *Warnings and Precautions (5.1)*], reserve OPANA ER

ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.

- OPANA ER is not indicated as an as-needed (prn) analgesic.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Instructions

OPANA ER should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.

- Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals [*see Warnings and Precautions (5)*]. Initiate the dosing regimen for each patient individually, taking into account the patient's severity of pain, patient response, prior analgesic treatment experience, and risk factors for addiction, abuse, and misuse [*see Warnings and Precautions (5.1)*].
- Monitor patients closely for respiratory depression, especially within the first 24-72 hours of initiating therapy and following dosage increases with OPANA ER and adjust the dosage accordingly [*see Warnings and Precautions (5.3)*].

Instruct patients to swallow OPANA ER tablets whole [*see Patient Counseling Information (17)*]. Crushing, chewing, or dissolving OPANA ER tablets will result in uncontrolled delivery of oxymorphone and can lead to overdose or death [*see Warnings and Precautions (5.3)*].

Administer on an empty stomach, at least 1 hour prior to or 2 hours after eating.

OPANA ER is administered orally twice daily (every 12 hours).

2.2 Initial Dosage

Use of OPANA ER as the First Opioid Analgesic

Initiate treatment with OPANA ER with the 5 mg tablet orally every 12-hours.

Use of OPANA ER in Patients who are not Opioid Tolerant

The starting dose for patients who are not opioid tolerant is OPANA ER 5 mg orally every 12 hours.

Patients considered opioid tolerant are those taking, for one week or longer, at least 60 mg oral morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, 60 mg oral hydrocodone per day, or an equianalgesic dose of another opioid.

Use of higher starting doses in patients who are not opioid tolerant may cause fatal respiratory depression.

Conversion from OPANA to OPANA ER

Patients receiving OPANA may be converted to OPANA ER by administering half the patient's total daily oral OPANA dose as OPANA ER, every 12 hours.

Conversion from Parenteral Oxymorphone to OPANA ER

The absolute oral bioavailability of OPANA ER is approximately 10%. Convert patients receiving parenteral oxymorphone to OPANA ER by administering 10 times the patient's total daily parenteral oxymorphone dose as OPANA ER in two equally divided doses (e.g., [IV dose x 10] divided by 2). Due to patient variability with regards to opioid analgesic response, upon conversion monitor patients closely to evaluate for adequate analgesia and side effects.

Conversion from Other Oral Opioids to OPANA ER

Discontinue all other around-the-clock opioid drugs when OPANA ER therapy is initiated.

While there are useful tables of opioid equivalents readily available, there is substantial inter-patient variability in the relative potency of different opioid drugs and products. As such, it is preferable to underestimate a patient's 24-hour oral oxymorphone requirements and provide rescue medication (e.g., immediate-release opioid) than to overestimate the 24-hour oral oxymorphone requirements which could result in adverse reactions. In an OPANA ER clinical trial with an open-label titration period, patients were converted from their prior opioid to OPANA ER using Table 1 as a guide for the initial OPANA ER dose.

Consider the following when using the information in Table 1:

- This is **not** a table of equianalgesic doses.
- The conversion factors in this table are only for the conversion **from** one of the listed oral opioid analgesics **to** OPANA ER.
- This table **cannot** be used to convert **from** OPANA ER **to** another opioid. Doing so will result in an over-estimation of the dose of the new opioid and may result in fatal overdose.

CONVERSION FACTORS TO OPANA ER	
Prior Oral Opioid	Approximate Oral Conversion Factor
Oxymorphone	1
Hydrocodone	0.5
Oxycodone	0.5
Methadone	0.5
Morphine	0.333

To calculate the estimated OPANA ER dose using Table 1:

- For patients on a single opioid, sum the current total daily dose of the opioid and then multiply the total daily dose by the conversion factor to calculate the approximate oral (active opioid) daily dose.
- For patients on a regimen of more than one opioid, calculate the approximate oral (active opioid) dose for each opioid and sum the totals to obtain the approximate total (active opioid) daily dose.
- For patients on a regimen of fixed-ratio opioid/non-opioid analgesic products, use only the opioid component of these products in the conversion

Always round the dose down, if necessary, to the appropriate OPANA ER strength(s) available.

Example conversion from a single opioid to OPANA ER:

Step 1: Sum the total daily dose of the opioid oxycodone 20 mg twice daily

20 mg former opioid 2 times daily = 40 mg total daily dose of former opioid

Step 2: Calculate the approximate equivalent dose of oral (active opioid) based on the total daily dose of the current opioid using Table 1

40 mg total daily dose of former opioid x 0.5 mg Conversion Factor = 20 mg of oral (active opioid) daily

Step 3: Calculate the approximate starting dose of OPANA ER to be given every 12 hours. Round down, if necessary, to the appropriate OPANA ER TABLETS strengths available.

10 mg OPANA ER every 12 hours

Conversion from Methadone to OPANA ER

Close monitoring is of particular importance when converting from methadone to other opioid agonists. The ratio between methadone and other opioid agonists may vary widely as a function of previous dose exposure. Methadone has a long half-life and can accumulate in the plasma.

2.3 Titration and Maintenance of Therapy

Individually titrate OPANA ER to a dose that provides adequate analgesia and minimizes adverse reactions. Continually reevaluate patients receiving OPANA ER to assess the maintenance of pain control and the relative incidence of adverse reactions, as well as monitoring for the development of addiction, abuse, and misuse. Frequent communication is important among the prescriber, other members of the healthcare team, the patient, and the caregiver/family during periods of changing analgesic requirements, including initial titration. During chronic therapy, periodically reassess the continued need for the use of opioid analgesics.

If the level of pain increases, attempt to identify the source of increased pain, while adjusting the OPANA ER dose to decrease the level of pain. Because steady-state plasma concentrations are approximated within 3 days, OPANA ER dosage adjustments, preferably at increments of 5-10 mg every 12 hours, may be done every 3 to 7 days.

Patients who experience breakthrough pain may require a dose increase of OPANA ER, or may need rescue medication with an appropriate dose of an immediate-release analgesic. If the level of pain increases after dose stabilization, attempt to identify the source

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.