



Drug Facts (continued)
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings) ■ drink a full glass of water with each dose
 ■ if taken with food, this product may take longer to work

adults and children 12 years and older: ■ take 1 capsule every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 capsules within the first hour ■ do not exceed 2 capsules in any 8- to 12- hour period ■ do not exceed 3 capsules in a 24 hour period

children under 12 years: ■ ask a doctor

Other information ■ each capsule contains: sodium 20 mg ■ contains color additives including FD&C Yellow No. 5 (tartrazine) ■ store at 20°-25°C (68°-77°F) avoid high humidity and excessive heat above 40°C (104°F) ■ do not use if carton is open or if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing. ■ read all directions and warnings before use. Keep carton.

NDC 00000-000-00

See new warnings information.

Naproxen Sodium Capsules, 220 mg

Pain Reliever/Fever Reducer (NSAID)

15 Capsules*

*Each Liquid-filled capsule contains 200 mg naproxen



Drug Facts

Active ingredient (in each capsule) Naproxen Sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Purposes

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache
 ■ the common cold ■ menstrual cramps
 ■ temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Lot #
 Expiration Date
 Rev # 2/06

Manufactured by:
 Banner Pharmaceuticals Inc.
 4725 Premier Drive
 High Point, NC 27265



Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if you have ■ problems or serious side effects from taking pain relievers or fever reducers
 ■ stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain ■ ulcers ■ bleeding problems ■ high blood pressure ■ heart or kidney disease ■ taken a diuretic ■ reached age 60 or older

Ask a doctor or pharmacist before use if you are ■ taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug
 ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs
 ■ long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days
 ■ stomach pain or upset gets worse or lasts ■ redness or swelling is present in the painful area ■ any new symptoms appear ■ you have difficulty swallowing ■ it feels like the capsule is stuck in your throat
 ■ you develop heartburn



#dd
 pp#
 12015

(b) (4) **Pharmalabel**

NDC 00000-000-00

See new warnings information.

Naproxen Sodium Capsules, 220 mg

Pain Reliever/Fever Reducer (NSAID)

15 Capsules*

*Each Liquid-filled capsule contains 200 mg naproxen

Active Ingredient Purposes

(Naproxen Sodium 220 mg) Pain reliever/
(naproxen 200 mg) fever reducer
(NSAID)* nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps. ■ temporarily reduces fever

Warnings
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
Allergy alert
Stomach bleeding warning

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ do not take longer than 10 days, unless directed by a doctor (see Warnings) ■ drink a full glass of water with each dose ■ if taken with food, this product may take longer to work

■ take 1 capsule every 8 to 12 hours while symptoms last
■ for the first dose you may take 2 capsules within the first hour
■ do not exceed 2 capsules in any 8- to 12-hour period
■ do not exceed 3 capsules in a 24-hour period
children under 12 years: ■ ask a doctor

Manufactured by: Banner Pharmaceuticals Inc.
4125 Pleasant Drive
High Point, NC 27265

Lot #
Expiration Date
Rev# 2.06

(b) (4)

IPharmaLabel



**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Andrea Segal
2/17/2006 02:39:32 PM