

.125" corner radius

2.375"



# Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

80 Capsules\*\*

\*\*Liquid-filled capsule

Strength to last 12 hours

Do not use if minimal safety seal under cap is torn or missing.  
**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**  
**Warnings Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives; facial swelling; asthma (wheezing); shock; skin reddening; rash; or blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer. **Ask a doctor or pharmacist before use if you are** • under a doctor's care for any serious condition • taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin • taking any other drug **Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), or if pain or fever gets worse. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** Adults and children 12 years and older: take 1 capsule with water every 8 to 12 hours while symptoms last. For the first dose, you may take 2 capsules within the first hour. Do not exceed 3 capsules in a 24-hour period. Children under 12 years: ask a doctor. **Questions or comments? 1-888-235-2466 (Mon - Fri 9AM - 5PM EST)**

Distributed by:  
**BIONPHARMA**  
 600 Alexander Road,  
 Princeton, NJ 08540  
 Lev/EXP

L000000  
 R0519

Non Varnish Area

6.00"

<b>BIONPHARMA</b>			
<b>Labeling Format Information:</b>			
<b>Fonts:</b> Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
<b>Drug Facts:</b>	N/A	<b>Leading:</b>	6.5pt
<b>Header:</b>	6pt	<b>Bullets:</b>	3.5pt
<b>Subheader:</b>	6pt	<b>Barlines:</b>	N/A
<b>Body Text:</b>	6pt	<b>Hairlines:</b>	N/A
<b>Drug Facts (continued):</b>	N/A		
Horizontal Scale: 100%		Average Kerning: 0%	

**Be Health**

# Naproxen Sodium Capsules, 220 mg (NSAID)

Do not use if printed safety seal under cap is torn or missing.

**Be Health**

NDC 69452-179-78

# Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

80 Capsules\*\*

\*\*Liquid-filled capsule

†Compare to the active ingredient in ALEVE® Liquid Gels

Strength to last 12 hours

**Be Health**

NDC 69452-179-78

†Compare to the active ingredient in ALEVE® Liquid Gels

# Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

80 Capsules\*\*

\*\*Liquid-filled capsule

Strength to last 12 hours

**Be Health**

NDC 69452-179-78

†Compare to the active ingredient in ALEVE® Liquid Gels

# Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

80 Capsules\*\*

\*\*Liquid-filled capsule

Strength to last 12 hours

**Be Health**

NDC 69452-179-78

†Compare to the active ingredient in ALEVE® Liquid Gels

# Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

80 Capsules\*\*

\*\*Liquid-filled capsule

Strength to last 12 hours

**Be Health**

NDC 69452-179-78

†Compare to the active ingredient in ALEVE® Liquid Gels

# Naproxen Sodium Capsules, 220 mg (NSAID)

Pain reliever/fever reducer

80 Capsules\*\*

\*\*Liquid-filled capsule

Strength to last 12 hours

**Drug Facts (continued)**

**When using this product**

- take with food or milk if stomach upset occurs
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

**Other information**

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep caution.

**Inactive ingredients**

FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polyvinyl alcohol, purified water, sodium benzoate, sodium bicarbonate, sodium chloride, sodium phosphate, sodium stearate, sodium sulfate, sodium sulfite, sodium tartrate, titanium dioxide, xanthan gum.

**Questions or comments?**

1-866-235-2466 (Mon-Fri 9AM-5PM EST)

**Drug Facts**

**Active ingredient (in each capsule)**

Naproxen sodium 220 mg (Naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - backache
  - headache
  - toothache
- minor pain of arthritis
- menstrual cramps
- the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away. Do not take naproxen sodium if you are taking aspirin for a long time, or if you have a severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription ibuprofen, naproxen, or others
- have 3 or more alcoholic drinks every day while using this drug

**Stomach bleeding warning:**

This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription ibuprofen, naproxen, or others
- have 3 or more alcoholic drinks every day while using this drug

**Heart attack and stroke warning:**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk may be higher if you take naproxen sodium for a longer time than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you are taking aspirin for a long time
- if you are taking a diuretic
- if you have problems or serious side effects from taking pain relievers

**Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers

**Ask a doctor or pharmacist before use if you are**


- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may interfere with the benefit of aspirin
- taking any other drug

\*\*\*This product is not manufactured or distributed by the owners of ALEVE® Liquid Gels.


Distributed by:  
**BOXPHARMA**  
600 Alexander Road,  
Princeton, NJ 08540

L0000000  
R0519

# Front



# Back



BIONPHARMA			
Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	9pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	2pt

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

KAREN M MAHONEY  
02/27/2020 06:13:32 PM