

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OXYCONTIN® safely and effectively. See full prescribing information for OXYCONTIN.

OxyContin® (oxycodone hydrochloride controlled-release) Tablets CII
Initial U.S. Approval: 1982

WARNING: IMPORTANCE OF PROPER PATIENT SELECTION AND POTENTIAL FOR ABUSE

See full prescribing information for complete boxed warning.

- OxyContin contains oxycodone which is an opioid agonist and a Schedule II controlled substance with an abuse liability similar to morphine. (9)
- OxyContin is indicated for the management of moderate to severe pain when a continuous, around-the-clock opioid analgesic is needed for an extended period of time. (1)
- OxyContin is NOT intended for use on an as-needed basis. (1)
- OxyContin 60 mg and 80 mg Tablets, a single dose greater than 40 mg, or a total daily dose greater than 80 mg are only for use in opioid-tolerant patients to avoid fatal respiratory depression. (2.7)
- Patients should be assessed for their clinical risks for opioid abuse or addiction prior to being prescribed opioids. (2.2)
- OxyContin tablets must be swallowed whole and must not be cut, broken, chewed, crushed, or dissolved which can lead to rapid release and absorption of a potentially fatal dose of oxycodone. (2.1)
- The concomitant use with cytochrome P450 3A4 inhibitors such as macrolide antibiotics and protease inhibitors may result in an increase in oxycodone plasma concentrations and may cause potentially fatal respiratory depression. (7.2)

INDICATIONS AND USAGE

OxyContin is an opioid agonist indicated for:

- Management of moderate to severe pain when a continuous, around-the-clock opioid analgesic is needed for an extended period of time. (1)
- Not for use on an as-needed basis or in the immediate post-operative period. (1)

DOSAGE AND ADMINISTRATION

- Use low initial doses in patients who are not already opioid-tolerant, especially those who are receiving concurrent treatment with muscle relaxants, sedatives, or other central nervous system (CNS) active medications. (2.2)
- For patients already receiving opioids, use standard conversion ratio estimates. (2.2)
- Tablets must be swallowed whole and are not to be cut, broken, chewed, crushed, or dissolved (risk of potentially fatal dose). (2.1)

DOSAGE FORMS AND STRENGTHS

- Controlled-Release Tablets: 10 mg, 15 mg, 20 mg, 30 mg, 40 mg, 60 mg and 80 mg (3)

CONTRAINDICATIONS

- in patients who have significant respiratory depression (4)
- in patients who have or are suspected of having paralytic ileus (4)
- in patients who have acute or severe bronchial asthma (4)
- in patients with known hypersensitivity to oxycodone (4)

WARNINGS AND PRECAUTIONS

- Must be swallowed whole (5.1)
- May cause somnolence, dizziness, alterations in judgment and alterations in levels of consciousness, including coma. (5.2)
- Additive CNS effects are expected when used with alcohol, other opioids, or illicit drugs. (5.1, 5.3, 7.3)
- Use with caution in patients who are receiving other CNS depressants. (5.1, 5.3, 7.3)

- May cause respiratory depression, use with extreme caution in patients at risk of respiratory depression, elderly and debilitated patients (5.4)
- May aggravate convulsions in patients with convulsive disorders, and may induce or aggravate seizures in some clinical settings. (5.5)
- May worsen increased intracranial pressure and obscure its signs, such as level of consciousness or pupillary signs. (5.6)
- May cause hypotension, use with caution in patients at increased risk of hypotension and in patients in circulatory shock. (5.7)
- Concomitant use of CYP3A4 inhibitors may increase opioid effects (5.8)
- Mixed agonist/antagonist analgesics may precipitate withdrawal symptoms. (5.9)
- Use with caution in patients with biliary tract disease, including acute pancreatitis. (5.10)
- Use with caution in patients at risk for ileus. Monitor for decreased bowel motility in postoperative patients. (5.10)
- Tolerance may develop. (5.11)
- Use with caution in alcoholism; adrenocortical insufficiency; hypothyroidism; prostatic hypertrophy or urethral stricture; severe impairment of hepatic, pulmonary or renal function; and toxic psychosis. (5.12)
- May impair the mental and physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. (5.13)
- No approved use in the treatment of addiction. (5.14)
- Not every urine drug test for "opioids" or "opiates" detects oxycodone reliably. (5.15)

ADVERSE REACTIONS

Most common adverse reactions (>5%) are constipation, nausea, somnolence, dizziness, vomiting, pruritus, headache, dry mouth, asthenia, and sweating.

To report Suspected Adverse Reactions, contact Purdue Pharma L.P. at 1-888-726-7535 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- OxyContin may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression. (7.1)
- The CYP3A4 isoenzyme plays a major role in the metabolism of OxyContin, drugs that inhibit CYP3A4 activity may cause decreased clearance of oxycodone which could lead to an increase in oxycodone plasma concentrations. (7.2)
- Concurrent use of other CNS depressants may cause respiratory depression, hypotension, and profound sedation or coma. (7.3)
- Mixed agonist/antagonist analgesics may reduce the analgesic effect of oxycodone and may precipitate withdrawal symptoms in these patients. (7.4)

USE IN SPECIFIC POPULATIONS

- Labor and Delivery: Not recommended for use in women immediately prior to and during labor and delivery; (8.2)
- Nursing Mothers: Nursing should not be undertaken while a patient is receiving OxyContin. (8.3)
- Pediatrics: Safety and effectiveness in pediatric patients below the age of 18 have not been established. (8.4)
- Geriatrics: The initial dose may need to be reduced to 1/3 to 1/2 of the usual doses. (8.5)
- Hepatic impairment: Initiate therapy at 1/3 to 1/2 the usual doses and titrate carefully. (8.6)
- Renal impairment: Dose initiation should follow a conservative approach. (8.7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: April 2010

FULL PRESCRIBING INFORMATION: CONTENTS*

BOX WARNING

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Safe Administration Instructions
- 2.2 Initiating Therapy with OxyContin
- 2.3 Conversion from Transdermal Fentanyl to OxyContin
- 2.4 Hepatic Impairment
- 2.5 Managing Expected Opioid Adverse Reactions
- 2.6 Individualization of Dosage
- 2.7 Special Instructions for Patients who are not Opioid Tolerant
- 2.8 Continuation of Therapy
- 2.9 Cessation of Therapy
- 2.10 Conversion from OxyContin to Parenteral Opioids

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Information Essential for Safe Administration
- 5.2 CNS Depression
- 5.3 Interactions with Alcohol, CNS Depressants and Illicit Drugs
- 5.4 Respiratory Depression
- 5.5 Seizures
- 5.6 Head Injury
- 5.7 Hypotensive Effect
- 5.8 Cytochrome P450 3A4 Inhibitors and Inducers
- 5.9 Interactions with Mixed Agonist/Antagonist Opioid Analgesics
- 5.10 Use in Pancreatic/Biliary Tract Disease and Other Gastrointestinal Conditions
- 5.11 Tolerance
- 5.12 Special Risk Groups
- 5.13 Driving and Operating Machinery
- 5.14 Use in Addiction Treatment
- 5.15 Laboratory Monitoring

6 ADVERSE REACTIONS

- 6.1 Clinical Trial Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Neuromuscular Junction Blocking Agents
- 7.2 Agents Affecting Cytochrome P450 Isoenzymes
- 7.3 CNS Depressants
- 7.4 Interactions with Mixed Agonist/Antagonist Opioid Analgesics

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Renal Impairment
- 8.8 Gender Differences

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Information for Patients and Caregivers

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: IMPORTANCE OF PROPER PATIENT SELECTION AND POTENTIAL FOR ABUSE

OxyContin contains oxycodone which is an opioid agonist and a Schedule II controlled substance with an abuse liability similar to morphine. (9)

OxyContin can be abused in a manner similar to other opioid agonists, legal or illicit. This should be considered when prescribing or dispensing OxyContin in situations where the physician or pharmacist is concerned about an increased risk of misuse, abuse, or diversion. (9.2)

OxyContin is a controlled-release oral formulation of oxycodone hydrochloride indicated for the management of moderate to severe pain when a continuous, around-the-clock opioid analgesic is needed for an extended period of time. (1)

OxyContin is not intended for use on an as-needed basis. (1)

Patients considered opioid tolerant are those who are taking at least 60 mg oral morphine/day, 25 mcg transdermal fentanyl/hour, 30 mg oral oxycodone/day, 8 mg oral hydromorphone/day, 25 mg oral oxymorphone/day, or an equianalgesic dose of another opioid for one week or longer.

OxyContin 60 mg and 80 mg tablets, a single dose greater than 40 mg, or a total daily dose greater than 80 mg are only for use in opioid-tolerant patients, as they may cause fatal respiratory depression when administered to patients who are not tolerant to the respiratory-depressant or sedating effects of opioids. (2.7)

Persons at increased risk for opioid abuse include those with a personal or family history of substance abuse (including drug or alcohol abuse or addiction) or mental illness (e.g., major depression). Patients should be assessed for their clinical risks for opioid abuse or addiction prior to being prescribed opioids. All patients receiving opioids should be routinely monitored for signs of misuse, abuse and addiction. (2.2)

OxyContin must be swallowed whole and must not be cut, broken, chewed, crushed, or dissolved. Taking cut, broken, chewed, crushed or dissolved OxyContin tablets leads to rapid release and absorption of a potentially fatal dose of oxycodone. (2.1)

The concomitant use of OxyContin with all cytochrome P450 3A4 inhibitors such as macrolide antibiotics (e.g., erythromycin), azole-antifungal agents (e.g., ketoconazole), and protease inhibitors (e.g., ritonavir) may result in an increase in oxycodone plasma concentrations, which could increase or prolong adverse effects and may cause potentially fatal respiratory depression. Patients receiving OxyContin and a CYP3A4 inhibitor should be carefully monitored for an extended period of time and dosage adjustments should be made if warranted (7.2).

1 INDICATIONS AND USAGE

OxyContin is a controlled-release oral formulation of oxycodone hydrochloride indicated for the management of moderate to severe pain when a continuous, around-the-clock opioid analgesic is needed for an extended period of time.

Limitations of Usage

OxyContin is not intended for use on an as-needed basis.

OxyContin is not indicated for the management of pain in the immediate postoperative period (the first 12-24 hours following surgery), or if the pain is mild, or not expected to persist for an extended period of time. OxyContin is indicated for postoperative use following the immediate post-operative period only if the patient is already receiving the drug prior to surgery or if the postoperative pain is expected to be moderate to severe and persist for an extended period of time. Physicians should individualize treatment, moving from parenteral to oral analgesics as appropriate. (See American Pain Society guidelines.)

OxyContin is not indicated for pre-emptive analgesia (preoperative administration for the management of postoperative pain).

OxyContin is not indicated for rectal administration.

2 DOSAGE AND ADMINISTRATION

2.1 Safe Administration Instructions

OxyContin-tablets must be swallowed whole and must not be cut, broken, chewed, crushed or dissolved. Taking cut, broken, chewed, crushed or dissolved OxyContin tablets leads to rapid release and absorption of a potentially fatal dose of oxycodone.

Selection of patients for treatment with OxyContin should be governed by the same principles that apply to the use of similar opioid analgesics. Physicians should individualize treatment using a progressive plan of pain management such as outlined by the World Health Organization, Federation of State Medical Boards Model Policy, and the American Pain Society. Healthcare professionals should follow appropriate pain management principles of careful assessment and ongoing monitoring.

2.2 Initiating Therapy with OxyContin

It is critical to initiate the dosing regimen for each patient individually. Attention should be given to:

- risk factors for abuse or addiction; including whether the patient has a previous or current substance abuse problem, a family history of substance abuse, or a history of mental illness or depression;
- the age, general condition and medical status of the patient;
- the patient's opioid exposure and opioid tolerance (if any);
- the daily dose, potency, and kind of the analgesic(s) the patient has been taking;
- the reliability of the conversion estimate used to calculate the dose of oxycodone;
- the special instructions for OxyContin 60 mg and 80 mg tablets, a single dose greater than 40 mg, **or total daily doses greater than 80 mg** [see *Dosage and Administration* (2.7)]; and
- the balance between pain control and adverse reactions.

Use low initial doses of OxyContin in patients who are not already opioid-tolerant [see *Dosage and Administration* (2.7)], especially those who are receiving concurrent treatment with muscle relaxants, sedatives, or other CNS active medications [see *Warnings and Precautions* (5.1, 5.3) and *Drug Interactions* (7.1, 7.3)].

Experience indicates a reasonable starting dose of OxyContin for patients who are taking non-opioid analgesics and require continuous around-the-clock therapy for an extended period of time is 10 mg every 12 hours. Individually titrate OxyContin to a dose that provides adequate analgesia and minimizes adverse reactions while maintaining an every-twelve-hour dosing regimen.

For initiation of OxyContin therapy for patients previously taking opioids, the conversion ratios found in Table 1 are a reasonable starting point, although not verified in well-controlled, multiple-dose trials. No fixed conversion ratio is likely to be satisfactory in all patients, especially patients receiving large opioid doses. A reasonable approach for converting from existing opioid therapy to OxyContin is as follows:

- Discontinue all other around-the-clock opioid drugs when OxyContin therapy is initiated.
- Using standard conversion ratio estimates (see Table 1), multiply the mg/day of each of the current opioids to be converted by their appropriate multiplication factor to obtain the equivalent total daily dose of oral oxycodone.
- Divide the calculated 24-hour oxycodone dose in half to approximate the every 12-hour dose of OxyContin.
- Round down, if necessary, to the appropriate OxyContin tablet strengths available.
- Close observation and frequent titration are indicated until patients are stable on the new therapy.

TABLE 1

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.