

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:
201655Orig1s000

LABELING

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OPANA® ER safely and effectively. See full prescribing information for OPANA® ER.

OPANA® ER (oxymorphone hydrochloride) Extended-Release tablets, CII

Initial U.S. Approval: 1959

WARNING: POTENTIAL FOR ABUSE, IMPORTANCE OF PROPER PATIENT SELECTION AND LIMITATIONS OF USE

See full prescribing information for complete boxed warning.

- OPANA ER contains oxymorphone which is an opioid agonist and a Schedule II controlled substance with an abuse liability similar to other opioid analgesics. (9)
- Oxymorphone can be abused in a manner similar to other opioid agonists, legal or illicit. This should be considered when prescribing or dispensing OPANA ER in situations where the physician or pharmacist is concerned about an increased risk of misuse, abuse, or diversion. (9)
- OPANA ER is NOT intended for use as an as needed analgesic. (1)
- OPANA ER tablets are to be swallowed whole and are not to be cut, broken, chewed, dissolved, or crushed as this leads to rapid release and absorption of a potentially fatal dose of oxymorphone. (2)
- Patients must not consume alcoholic beverages, prescription or non-prescription medications containing alcohol. Co-ingestion of alcohol with OPANA ER may result in a potentially fatal overdose of oxymorphone. (2)

INDICATIONS AND USAGE

- OPANA ER is an opioid agonist indicated for the relief of moderate to severe pain in patients requiring continuous around-the-clock opioid treatment for an extended period of time. (1)
- Not intended for use as an as needed analgesic. Not indicated in the immediate post-operative period or if the pain is mild or not expected to persist for an extended period of time. (1)

DOSAGE AND ADMINISTRATION

- Administer on an empty stomach, at least 1 hour prior to or 2 hours after eating. (2.2)
- Symmetrical, every 12 h dosing is appropriate for the majority of patients. (2.1)
- Opioid-Naïve Patients: Initiate treatment with 5 mg every 12 hours. (2.2)
- Opioid-Experienced Patients: Ratios as a guide to convert only from other opioids to OPANA ER. (2.2)
- Individualize treatment; titrate to effective and tolerable dose. (2.1)
- Don't stop abruptly (9.3); taper gradually to stop treatment. (2.8)
- OPANA ER tablets should be taken one tablet at a time, with enough water to ensure complete swallowing immediately after placing in the mouth (2.1, 17)

DOSAGE FORMS AND STRENGTHS

- Extended-Release Tablets, 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, and 40 mg (3)

CONTRAINDICATIONS

- Known hypersensitivity to oxymorphone, any other ingredients in OPANA ER, or morphine analogs. (4)
- Respiratory depression (4)
- Acute or severe bronchial asthma or hypercarbia (4)
- Paralytic ileus (4)
- Moderate or severe hepatic impairment (4)

WARNINGS AND PRECAUTIONS

See Boxed WARNINGS

- Respiratory depression: Increased risk in elderly, debilitated patients, and those suffering from conditions accompanied by hypoxia, hypercapnia, or decreased respiratory reserve. (5.2)
- Misuse, abuse, and diversion: OPANA ER is an opioid agonist and a Schedule II controlled substance with an abuse liability similar to morphine. (5.3)
- CNS effects: Additive CNS-depressive effects when used in conjunction with alcohol, other opioids, or illicit drugs. (5.4)
- Head Injury: Effects may be markedly exaggerated. Administer OPANA ER with extreme caution. (5.5)
- Hypotensive effect: Increased risk with compromised ability to maintain blood pressure. Administer with caution to patients in circulatory shock. (5.6)
- Mild hepatic impairment: Use with caution and at lower doses due to higher plasma concentrations than in patients with normal hepatic function. (5.7)
- Prolonged gastric obstruction: May occur in patients with gastrointestinal obstruction. (5.9)
- Sphincter of Oddi: Administer with caution in patients with biliary tract disease. (5.11)
- Impaired mental/physical abilities: Caution must be used with potentially hazardous activities (5.12)

ADVERSE REACTIONS

Adverse reactions in $\geq 2\%$ of oxymorphone ER-treated patients in placebo-controlled trials: nausea, constipation, dizziness, somnolence, vomiting, pruritus, headache, sweating increased, dry mouth, sedation, diarrhea, insomnia, fatigue, appetite decreased, and abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Endo Pharmaceuticals Inc. at (1-800-462-3636) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- CNS depressants: Increased risk of respiratory depression, hypotension, profound sedation, coma or death when combined with OPANA ER. When combined therapy with CNS depressant is contemplated, the dose of one or both agents should be reduced. (7.2)
- Mixed agonist/antagonist opioids (i.e., pentazocine, nalbuphine, and butorphanol): May reduce analgesic effect and/or precipitate withdrawal symptoms. (7.3)
- Cimetidine: Combination use with OPANA ER may precipitate confusion, disorientation, respiratory depression, apnea, seizures. (7.4)
- Anticholinergics: Concurrent use with OPANA ER may result in urinary retention and/or severe constipation, which may lead to paralytic ileus. (7.5)
- Monoamine oxidase inhibitors (MAOIs): Potentiate the action of opioids. OPANA ER should not be used in patients taking MAOIs or within 14 days of stopping such treatment. (7.6)

USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm. (8.1)
- Geriatric Patients – OPANA ER should be used with caution in elderly patients. (8.5)

See 17 for PATIENT COUNSELING INFORMATION and FDA – approved Medication Guide

Revised: 12/2011

FULL PRESCRIBING INFORMATION: CONTENTS*

BOXED WARNING

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Safe Administration Instructions
- 2.2 Initiating Therapy with OPANA ER
- 2.3 Patients with Hepatic Impairment
- 2.4 Patients with Renal Impairment
- 2.5 Use with Central Nervous System Depressants
- 2.6 Geriatric Patients
- 2.7 Maintenance of Therapy
- 2.8 Cessation of Therapy

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Information Essential for Safe Administration
- 5.2 Respiratory Depression

- 5.3 Misuse, Abuse and Diversion of Opioids
- 5.4 Interactions with Alcohol and other CNS Depressants
- 5.5 Use in Patients with Head Injury and Increased Intracranial Pressure
- 5.6 Hypotensive Effect
- 5.7 Hepatic Impairment
- 5.8 Special Risk Groups
- 5.9 Gastrointestinal Effects
- 5.10 Ambulatory Surgery and Post Operative Use
- 5.11 Use in Pancreatic/Biliary Tract Disease
- 5.12 Driving and Operating Machinery
- 6 ADVERSE REACTIONS
- 6.1 Clinical Trial Experience
- 7 DRUG INTERACTIONS
- 7.1 Drug-Drug Interactions
- 7.2 Use with CNS Depressants
- 7.3 Interactions with Mixed Agonist/Antagonist Opioid Analgesics
- 7.4 Cimetidine

- 7.5 Anticholinergics
- 7.6 MAO Inhibitors
- 8 USE IN SPECIFIC POPULATIONS**
 - 8.1 Pregnancy
 - 8.2 Labor and Delivery
 - 8.3 Nursing Mothers
 - 8.4 Pediatric Use
 - 8.5 Geriatric Use
 - 8.6 Hepatic Impairment
 - 8.7 Renal Impairment
- 9 DRUG ABUSE AND DEPENDENCE**
 - 9.1 Controlled Substance
 - 9.2 Abuse
 - 9.3 Dependence
- 10 OVERDOSAGE**
 - 10.1 Symptoms

- 10.2 Treatment
- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.2 Pharmacodynamics
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
 - 14.1 12-Week Study in Opioid-Naïve Patients with Low Back Pain
 - 14.2 12-Week Study in Opioid-Experienced Patients with Low Back Pain
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
- 17 PATIENT COUNSELING INFORMATION**

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: POTENTIAL FOR ABUSE, IMPORTANCE OF PROPER PATIENT SELECTION AND LIMITATIONS OF USE

Potential for Abuse

OPANA ER contains oxymorphone, which is a morphine-like opioid agonist and a Schedule II controlled substance, with an abuse liability similar to other opioid analgesics. (9)

Oxymorphone can be abused in a manner similar to other opioid agonists, legal or illicit. This should be considered when prescribing or dispensing OPANA ER in situations where the physician or pharmacist is concerned about an increased risk of misuse, abuse, or diversion. (9.2)

Proper Patient Selection

OPANA ER is an extended-release oral formulation of oxymorphone indicated for the management of moderate to severe pain when a continuous, around-the-clock opioid analgesic is needed for an extended period of time. (1)

Limitations of Use

OPANA ER is NOT intended for use as an as needed analgesic. (1)

OPANA ER tablets are to be swallowed whole and are not to be cut, broken, chewed, dissolved, or crushed. Taking cut, broken, chewed, dissolved, or crushed OPANA ER tablets leads to rapid release and absorption of a potentially fatal dose of oxymorphone. (2)

Patients must not consume alcoholic beverages, or prescription or non-prescription medications containing alcohol, while on OPANA ER therapy. The co-ingestion of alcohol with OPANA ER may result in increased plasma levels and a potentially fatal overdose of oxymorphone. (2)

1 INDICATIONS AND USAGE

OPANA ER is indicated for the relief of moderate to severe pain in patients requiring continuous, around-the-clock opioid treatment for an extended period of time.

Limitations of Usage

OPANA ER is not intended for use as an as needed analgesic.

OPANA ER is not indicated for pain in the immediate post-operative period if the pain is mild, or not expected to persist for an extended period of time.

OPANA ER is only indicated for post-operative use if the patient is already receiving the drug prior to surgery or if the post-operative pain is expected to be moderate or severe and persist for an extended period of time. Physicians should individualize treatment, moving from parenteral to oral analgesics as appropriate. (See American Pain Society guidelines).

2 DOSAGE AND ADMINISTRATION

2.1 Safe Administration Instructions

OPANA ER tablets must be swallowed whole and are not to be cut, broken, chewed, dissolved, or crushed. Taking cut, broken, chewed, dissolved, or crushed OPANA ER tablets leads to rapid release and absorption of a potentially fatal dose of oxymorphone.

Patients must not consume alcoholic beverages, or prescription or non-prescription medications containing alcohol, while on OPANA ER therapy. The co-ingestion of alcohol with OPANA ER may result in increased plasma levels and a potentially fatal overdose of oxymorphone.

OPANA ER tablets must be taken whole, one tablet at a time, with enough water to ensure complete swallowing immediately after placing in the mouth [see Patient Counseling Information (17)].

OPANA ER must be taken on an empty stomach, at least one hour prior to or two hours after eating [see Clinical Pharmacology (12.3)].

While symmetric (same dose AM and PM), around-the-clock, every 12 hours dosing is appropriate for the majority of patients, some patients may benefit from asymmetric (different dose given in AM than in PM) dosing, tailored to their pain pattern. It is usually appropriate to treat a patient with only one extended-release opioid for around-the-clock therapy.

Selection of patients for treatment with OPANA ER should be governed by the same principles that apply to the use of other extended-release opioid analgesics [see *Indications and Usage (1)*]. Physicians should individualize treatment in every case, using non-opioid analgesics, opioids on an as needed basis, combination products, and chronic opioid therapy in a progressive plan of pain management such as outlined by the World Health Organization, the American Pain Society and the Federation of State Medical Boards Model Guidelines. Healthcare professionals should follow appropriate pain management principles of careful assessment and ongoing monitoring [see *Boxed Warning*].

2.2 Initiating Therapy with OPANA ER

It is necessary to adjust the dosing regimen for each patient individually, taking into account the patient's prior analgesic treatment experience. In the selection of the initial dose of OPANA ER, attention should be given to the following:

- total daily dose, potency and specific characteristics of the opioid the patient has been taking previously;
- relative potency estimate used to calculate the equivalent oxymorphone dose needed;
- patient's degree of opioid tolerance;
- age, general condition, and medical status of the patient;
- concurrent non-opioid analgesics and other medications;
- type and severity of the patient's pain;
- balance between pain control and adverse experiences;
- risk factors for abuse or addiction, including a prior history of abuse or addiction.

Once therapy is initiated, frequently assess pain relief and other opioid effects. Base the titration of the total daily OPANA ER dose upon the amount of supplemental opioid utilization, severity of the patient's pain, and the patient's ability to tolerate the opioid. Titrate dose to generally mild or no pain with the regular use of no more than two doses of supplemental analgesia, i.e. "rescue," per 24 hours, and tolerable side effects. Patients who experience breakthrough pain may require dosage adjustment.

If signs of excessive opioid-related adverse experiences are observed, the next dose may be reduced. If this adjustment leads to inadequate analgesia, a supplemental dose of immediate-release opioid, or a non-opioid analgesic may be administered. Adjust dosing to obtain an appropriate balance between pain relief and opioid-related adverse experiences. If significant adverse events occur before the therapeutic goal of mild or no pain is achieved, the events should be treated aggressively. If adverse events are adequately managed, continue upward titration to an acceptable level of pain control.

During periods of changing analgesic requirements, including initial titration, frequent contact is recommended between physician, other members of the healthcare team, the patient and the caregiver/family. Advise patients and caregivers/family members of the potential adverse reactions.

The dosing recommendations below, therefore, can only be considered as suggested approaches to what is actually a series of clinical decisions over time in the management of the pain of each individual patient.

Titrate dose to adequate pain relief (generally mild or no pain).

Opioid-Naïve Patients

The initial dose for patients who are not opioid-experienced and who are being initiated on chronic around-the-clock opioid therapy with OPANA ER is 5 mg every 12 hours. Thereafter, titrate the dose individually at increments of 5-10 mg every 12 hours every 3-7 days, to a level that provides adequate analgesia and tolerable side effects under the close supervision of the prescribing physician.

Opioid-Experienced Patients

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.