

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

202895Orig1s000

LABELING

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PREZISTA safely and effectively. See Full Prescribing Information for PREZISTA.

**PREZISTA (darunavir) Oral Suspension, for Oral use
PREZISTA (darunavir) Tablet, Film Coated for Oral use**

Initial U.S. Approval – 2006

RECENT MAJOR CHANGES

- Indications and Usage
 - Pediatric Patients (1.2) 12/2011
- Dosage and Administration
 - Adult Patients (2.1) 12/2011
 - Pediatric Patients (2.2) 12/2011
- Warnings and Precautions
 - Severe Skin Reactions (5.3) 10/2011
 - Pediatric Patients (5.11) 12/2011

INDICATIONS AND USAGE

PREZISTA is a human immunodeficiency virus (HIV-1) protease inhibitor indicated for the treatment of HIV-1 infection in adult patients. PREZISTA is also indicated for the treatment of HIV-1 infection in pediatric patients 3 years of age and older. PREZISTA must be co-administered with ritonavir (PREZISTA/ritonavir) and with other antiretroviral agents. (1)

DOSAGE AND ADMINISTRATION

- Treatment-naïve adult patients and treatment-experienced adult patients with no darunavir resistance associated substitutions: 800 mg (two 400 mg tablets) taken with ritonavir 100 mg once daily and with food. (2.1)
- Treatment-experienced adult patients with at least one darunavir resistance associated substitution: 600 mg (one 600 mg tablet) taken with ritonavir 100 mg twice daily and with food. (2.1)
- Pediatric patients (3 to less than 18 years of age and weighing at least 10 kg): dosage of PREZISTA and ritonavir is based on body weight and should not exceed the treatment-experienced adult dose. Do not use once daily dosing in pediatric patients. PREZISTA should be taken with ritonavir twice daily and with food. (2.2)
- PREZISTA/ritonavir is not recommended for use in patients with severe hepatic impairment. (2.3)

DOSAGE FORMS AND STRENGTHS

- 100 mg/mL oral suspension (3)
- 75 mg tablets, 150 mg tablets, 400 mg tablets, and 600 mg tablets (3)

CONTRAINDICATIONS

Co-administration with alfuzosin, dihydroergotamine, ergonovine, ergotamine, methylergonovine, cisapride, pimozide, oral midazolam, triazolam, St. John's Wort, lovastatin, simvastatin, rifampin and sildenafil (for treatment of pulmonary arterial hypertension). (4)

- Due to the need for co-administration of PREZISTA with ritonavir, please refer to ritonavir prescribing information for a description of ritonavir contraindications. (4)

WARNINGS AND PRECAUTIONS

- Drug-induced hepatitis (e.g., acute hepatitis, cytolytic hepatitis) has been reported with PREZISTA/ritonavir. Monitor liver function before and during therapy, especially in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases. Post-marketing cases of liver injury, including some fatalities, have been reported. (5.2, 6)
- Skin reactions ranging from mild to severe, including Stevens-Johnson Syndrome and toxic epidermal necrolysis, have been reported. Discontinue treatment if severe reaction develops. (5.3, 6)
- Use with caution in patients with a known sulfonamide allergy. (5.4)
- Patients may develop new onset diabetes mellitus or hyperglycemia. Initiation or dose adjustments of insulin or oral hypoglycemic agents may be required. (5.6)
- Patients may develop redistribution/accumulation of body fat (5.7) or immune reconstitution syndrome. (5.8)
- Patients with hemophilia may develop increased bleeding events. (5.9)
- PREZISTA/ritonavir should not be used in pediatric patients below 3 years of age in view of toxicity and mortality observed in juvenile rats dosed with darunavir up to days 23 to 26 of age. (5.11)

ADVERSE REACTIONS

- The most common clinical adverse drug reactions to PREZISTA/ritonavir (incidence greater than or equal to 5% of at least moderate intensity (greater than or equal to Grade 2) were diarrhea, nausea, rash, headache, abdominal pain and vomiting. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Tibotec Therapeutics at 1-877-REACH-TT or 1-877-732-2488 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Co-administration of PREZISTA/ritonavir with other drugs can alter the concentration of other drugs and other drugs may alter the concentrations of darunavir. The potential drug-drug concentrations must be considered prior to and during therapy. (4, 5.5, 7, 12.3).

USE IN SPECIFIC POPULATIONS

- Use during pregnancy only if the potential benefit justifies the potential risk. (8.1)
 - Pregnancy Registry available. (8.1)
- Mothers should be instructed not to breastfeed due to the potential for HIV transmission and the potential for serious adverse reactions in nursing infants. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA approved patient labeling.

Revised: 12/2011

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE		
1.1	Adult Patients		7.1 Potential for PREZISTA/ritonavir to Affect Other Drugs
1.2	Pediatric Patients		7.2 Potential for Other Drugs to Affect Darunavir
2	DOSAGE AND ADMINISTRATION		7.3 Established and Other Potentially Significant Drug Interactions
2.1	Adult Patients		
2.2	Pediatric Patients (age 3 to less than 18 years)	8	USE IN SPECIFIC POPULATIONS
2.3	Patients with Hepatic Impairment	8.1	Pregnancy
3	DOSAGE FORMS AND STRENGTHS	8.3	Nursing Mothers
3.1	PREZISTA 100 mg/mL Oral Suspension	8.4	Pediatric Use
3.2	PREZISTA 75 mg Tablets	8.5	Geriatric Use
3.3	PREZISTA 150 mg Tablets	8.6	Hepatic Impairment
3.4	PREZISTA 400 mg Tablets	8.7	Renal Impairment
3.5	PREZISTA 600 mg Tablets	10	OVERDOSAGE
4	CONTRAINDICATIONS	11	DESCRIPTION
5	WARNINGS AND PRECAUTIONS	12	CLINICAL PHARMACOLOGY
5.1	General	12.1	Mechanism of Action
5.2	Hepatotoxicity	12.2	Pharmacodynamics
5.3	Severe Skin Reactions	12.3	Pharmacokinetics
5.4	Sulfa Allergy	12.4	Microbiology
5.5	Drug Interactions	13	NONCLINICAL TOXICOLOGY
5.6	Diabetes Mellitus/ Hyperglycemia	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
5.7	Fat Redistribution	13.2	Animal Toxicology and/or Pharmacology
5.8	Immune Reconstitution Syndrome	14	CLINICAL STUDIES
5.9	Hemophilia	14.1	Description of Adult Clinical Studies
5.10	Resistance / Cross-Resistance	14.2	Treatment-Naïve Adult Subjects
5.11	Pediatric Patients	14.3	Treatment-Experienced Adult Subjects
6	ADVERSE REACTIONS	14.4	Pediatric Patients
6.1	Clinical Trials Experience: Treatment-Naïve Adults	16	HOW SUPPLIED/STORAGE AND HANDLING
6.2	Clinical Trials Experience: Treatment-Experienced Adults	17	PATIENT COUNSELING INFORMATION
6.3	Serious ADRs	17.1	Information About Therapy with PREZISTA
6.4	Patients co-infected with hepatitis B and/or hepatitis C virus	17.2	Instructions for Use
6.5	Clinical Trials Experience: Pediatric Patients	17.3	Hepatotoxicity
6.6	Postmarketing Experience	17.4	Severe Skin Reactions
7	DRUG INTERACTIONS	17.5	Drug Interactions
		17.6	Fat Redistribution

[*Sections or subsections omitted from the Full Prescribing Information are not listed]

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Adult Patients

PREZISTA[®], co-administered with ritonavir (PREZISTA/ritonavir), and with other antiretroviral agents, is indicated for the treatment of human immunodeficiency virus (HIV-1) infection.

This indication is based on analyses of plasma HIV-1 RNA levels and CD4+ cell counts from 2 controlled Phase 3 trials of 48 weeks duration in antiretroviral treatment-naïve and treatment-experienced patients and 2 controlled Phase 2 trials of 96 weeks duration in clinically advanced, treatment-experienced adult patients.

1.2 Pediatric Patients

PREZISTA, co-administered with ritonavir (PREZISTA/ritonavir), and with other antiretroviral agents, is indicated for the treatment of HIV-1 infection in pediatric patients 3 years of age and older [*see Use in Specific Populations (8.4)*].

This indication is based on 24-week analyses of plasma HIV-1 RNA levels and CD4+ cell counts from 2 open-label Phase 2 trials in antiretroviral treatment-experienced pediatric patients (one trial in patients 6 to less than 18 years of age and one trial in patients 3 to less than 6 years of age).

In treatment-experienced adult and pediatric patients, the following points should be considered when initiating therapy with PREZISTA/ritonavir:

- Treatment history and, when available, genotypic or phenotypic testing should guide the use of PREZISTA/ritonavir [*see Clinical Pharmacology (12.4)*].
- The use of other active agents with PREZISTA/ritonavir is associated with a greater likelihood of treatment response [*see Clinical Pharmacology (12.4) and Clinical Studies (14.3)*].

2 DOSAGE AND ADMINISTRATION

2.1 Adult Patients

PREZISTA must be co-administered with ritonavir to exert its therapeutic effect. Failure to correctly co-administer PREZISTA with ritonavir will result in plasma levels of darunavir that will be insufficient to achieve the desired antiviral effect and will alter some drug interactions.

Patients who have difficulty swallowing PREZISTA tablets can use the 100 mg/mL PREZISTA oral suspension.

Treatment-Naïve Adult Patients

The recommended oral dose of PREZISTA is 800 mg (two 400 mg tablets or 8 mL of the oral suspension) taken with ritonavir 100 mg (one 100 mg tablet/capsule or 1.25 mL of a 80 mg/mL ritonavir oral solution) once daily and with food.

Treatment-Experienced Adult Patients

Treatment-Experienced Adult Patients	
With no darunavir resistance associated substitutions*	With at least one darunavir resistance associated substitution*
PREZISTA 800 mg (two 400 mg tablets or 8 mL [†]) once daily with ritonavir 100 mg (one 100 mg tablet/capsule or 1.25 mL) once daily and with food	PREZISTA 600 mg (e.g. one 600 mg tablet or 6 mL) twice daily with ritonavir 100 mg (one 100 mg tablet/capsule or 1.25 mL) twice daily and with food

*	V11I, V32I, L33F, I47V, I50V, I54L, I54M, T74P, L76V, I84V and L89V
†	An 8 mL dose should be taken as two 4 mL administrations with the included oral dosing syringe

For antiretroviral treatment-experienced patients genotypic testing is recommended. However, when genotypic testing is not feasible, PREZISTA/ritonavir 600/100 mg twice daily dosing is recommended.

2.2 Pediatric Patients (age 3 to less than 18 years)

Do not use once daily dosing in pediatric patients.

Healthcare professionals should pay special attention to accurate dose selection of PREZISTA, transcription of the medication order, dispensing information and dosing instruction to minimize risk for medication errors, overdose, and underdose.

Prescribers should select the appropriate dose of PREZISTA/ritonavir for each individual child based on body weight (kg) and should not exceed the recommended dose for treatment-experienced adults.

Before prescribing PREZISTA, children weighing greater than or equal to 15 kg should be assessed for the ability to swallow tablets. If a child is unable to reliably swallow a tablet, the use of PREZISTA oral suspension should be considered.

The recommended dose of PREZISTA/ritonavir for pediatric patients (3 to less than 18 years of age and weighing at least 10 kg is based on body weight (see Tables 1, 2 and 3) and should not exceed the recommended treatment-experienced adult dose (PREZISTA/ritonavir 600/100 mg twice daily). PREZISTA should be taken with ritonavir twice daily and with food.

Dosing recommendations for pediatric patients weighing at least 10 kg but less than 15 kg

The weight-based dose in pediatric patients weighing less than 15 kg is PREZISTA 20 mg/kg with ritonavir 3 mg/kg which can be dosed using the following table:

Table 1: Recommended Dose for Pediatric Patients with PREZISTA Oral Suspension (100 mg/mL) and Ritonavir Oral Solution* for Pediatric Patients Weighing 10 kg to Less Than 15 kg	
Body weight (kg)	Dose (twice daily with food)
Greater than or equal to 10 kg to less than 11 kg	PREZISTA 200 mg (2 mL) with ritonavir 32 mg (0.4 mL)
Greater than or equal to 11 kg to less than 12 kg	PREZISTA 220 mg (2.2 mL) with ritonavir 32 mg (0.4 mL)
Greater than or equal to 12 kg to less than 13 kg	PREZISTA 240 mg (2.4 mL) with ritonavir 40 mg (0.5 mL)
Greater than or equal to 13 kg to less than 14 kg	PREZISTA 260 mg (2.6 mL) with ritonavir 40 mg (0.5 mL)
Greater than or equal to 14 kg to less than 15 kg	PREZISTA 280 mg (2.8 mL) with ritonavir 48 mg (0.6 mL)

*with ritonavir oral solution: 80 mg/mL

Dosing recommendations for pediatric patients weighing at least 15 kg

Pediatric patients who weigh at least 15 kg and are able to swallow tablets can be dosed using the following table:

Table 2: Recommended Dose for Pediatric Patients with PREZISTA Tablets and Ritonavir Oral Solution or Tablets/Capsules for Pediatric Patients Weighing At Least 15 kg
--

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.