

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ZORVOLEX® safely and effectively. See full prescribing information for ZORVOLEX.

ZORVOLEX (diclofenac) capsules, for oral use

Initial U.S. Approval: 1988

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

See full prescribing information for complete boxed warning.

Cardiovascular Risk

- Nonsteroidal anti-inflammatory drugs (NSAIDs) may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. This risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk. (5.1)
- ZORVOLEX is contraindicated for the treatment of perioperative pain in the setting of coronary artery bypass graft (CABG) surgery. (4)

Gastrointestinal Risk

- NSAIDs cause an increased risk of serious gastrointestinal adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious gastrointestinal events. (5.2)

RECENT MAJOR CHANGES

- Indications and Usage, Osteoarthritis Pain (1.2) 8/2014

INDICATIONS AND USAGE

ZORVOLEX is an NSAID indicated for management of mild to moderate acute pain and osteoarthritis pain. (1)

DOSAGE AND ADMINISTRATION

- The dosage for acute pain is 18 mg or 35 mg orally three times a day. (1.2)
- The dosage for osteoarthritis pain is 35 mg orally three times a day. (1.2)
- Use lowest effective dosage for shortest duration consistent with individual patient treatment goals. (1.2)
- ZORVOLEX capsules are not interchangeable with other formulations of oral diclofenac even if the milligram strength is the same. (2.3)

DOSAGE FORMS AND STRENGTHS

Capsules: 18 mg or 35 mg (3)

CONTRAINDICATIONS

- Known hypersensitivity to diclofenac or any components of the drug product. (4)
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs. (4)
- Perioperative pain in the setting of coronary artery bypass graft (CABG) surgery. (4)

WARNINGS AND PRECAUTIONS

- Serious and potentially fatal cardiovascular (CV) thrombotic events, myocardial infarction, and stroke. Patients with known CV disease or risk

factors for CV disease may be at greater risk. Use the lowest effective dose for the shortest duration possible. (5.1)

- Serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation, which can be fatal. Prescribe ZORVOLEX with caution in patients with a prior history of ulcer disease or GI bleeding. Use the lowest effective dose for the shortest duration possible. (5.2)
- Elevation of one or more liver tests and severe hepatic reactions. Measure transaminases (ALT and AST) periodically in patients receiving long-term therapy with ZORVOLEX. Discontinue ZORVOLEX immediately if abnormal liver tests persist or worsen. (5.3)
- New onset or worsening of hypertension. Monitor blood pressure closely during treatment with ZORVOLEX. (5.4)
- Fluid retention and edema. Use ZORVOLEX with caution in patients with fluid retention or heart failure. (5.5)
- Renal papillary necrosis and other renal injury with long-term use. Use ZORVOLEX with caution in patients at greatest risk of this reaction, including the elderly, those with impaired renal function, heart failure, liver dysfunction, and those taking diuretics and ACE inhibitors. (5.6)
- Anaphylactoid reactions in patients with the aspirin triad or in patients without prior exposure to ZORVOLEX. Discontinue immediately if an anaphylactoid reaction occurs. (5.7)
- Serious skin adverse events such as exfoliative dermatitis, Stevens - Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. Discontinue ZORVOLEX if rash or other signs of local skin reaction occur. (5.8)

ADVERSE REACTIONS

Most common adverse reactions in controlled clinical trials (incidence $\geq 2\%$ in ZORVOLEX 18 mg or 35 mg group) include, edema, nausea, headache, dizziness, vomiting, constipation, pruritus, diarrhea, flatulence, pain in extremity, abdominal pain, sinusitis, alanine aminotransferase increased, blood creatinine increased, hypertension, and dyspepsia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Iroko Pharmaceuticals, LLC at 1-877-757-0676 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Concomitant use of ZORVOLEX and aspirin is not generally recommended because of the potential of increased adverse effects including increased GI bleeding. (7.1)
- Concomitant use of ZORVOLEX and anticoagulants have a risk of serious GI bleeding higher than users of either drug alone. (7.2)

USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm. Starting at 30 weeks gestation, ZORVOLEX should be avoided as premature closure of the ductus arteriosus in the fetus may occur. (5.9, 8.1)
- Nursing Mothers: Based on available data, diclofenac may be present in human milk. Exercise caution when ZORVOLEX is administered to a nursing woman. (8.3)
- Hepatic insufficiency: Patients with hepatic disease may require reduced doses of ZORVOLEX. Start treatment at the lowest dose. Discontinue use if efficacy is not achieved with the lowest dose. (2.2, 12.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 08/2014

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

1 INDICATIONS AND USAGE

- 1.1 Acute Pain
- 1.2 Osteoarthritis Pain

2 DOSAGE AND ADMINISTRATION

- 2.1 Initial Dosing
- 2.2 Dosage Adjustments in Patients with Hepatic Impairment
- 2.3 Non-Interchangeability with Other Formulations of Diclofenac

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Cardiovascular Thrombotic Events
- 5.2 Gastrointestinal (GI) Effects – Risk of GI Ulceration, Bleeding, and Perforation
- 5.3 Hepatic Effects
- 5.4 Hypertension
- 5.5 Congestive Heart Failure and Edema
- 5.6 Renal Effects
- 5.7 Anaphylactoid Reactions
- 5.8 Adverse Skin Reactions
- 5.9 Fetal Toxicity
- 5.10 Corticosteroid-Responsive Illness
- 5.11 Masking of Inflammation and Fever
- 5.12 Hematological Effects
- 5.13 Use in Patients with Preexisting Asthma
- 5.14 Monitoring

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Aspirin
- 7.2 Anticoagulants
- 7.3 ACE-inhibitors
- 7.4 Diuretics
- 7.5 Lithium
- 7.6 Methotrexate
- 7.7 Cyclosporine
- 7.8 Inhibitors or Substrates of Cytochrome P450 2C9 Other Considerations

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, and Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS

Cardiovascular Risk

- NSAIDs may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction, and stroke, which can be fatal. This risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk. [see *Warnings and Precautions* (5.1)]
- ZORVOLEX is contraindicated for the treatment of perioperative pain in the setting of coronary artery bypass graft (CABG) surgery. [see *Contraindications* (4)]

Gastrointestinal Risk

- NSAIDs cause an increased risk of serious gastrointestinal adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious gastrointestinal events. [see *Warnings and Precautions* (5.2)]

1 INDICATIONS AND USAGE

1.1 Acute Pain

ZORVOLEX is indicated for management of mild to moderate acute pain.

1.2 Osteoarthritis Pain

ZORVOLEX is indicated for management of osteoarthritis pain.

2 DOSAGE AND ADMINISTRATION

2.1 Initial Dosing

The effectiveness of ZORVOLEX when taken with food has not been studied in clinical studies. Taking ZORVOLEX with food may cause a reduction in effectiveness compared to taking ZORVOLEX on an empty stomach [see *Clinical Pharmacology* (12.3)]. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals [see *Warnings and Precautions* (5.1, 5.2)].

Acute Pain

For management of mild to moderate acute pain, the dosage is 18 mg or 35 mg orally three times daily.

Osteoarthritis Pain

For management of osteoarthritis pain, the dosage is 35 mg orally three times daily.

2.2 Dosage Adjustments in Patients with Hepatic Impairment

Patients with hepatic disease may require reduced doses of ZORVOLEX compared to patients with normal hepatic function [*see Clinical Pharmacology (12.3)*]. As with other diclofenac products, start treatment at the lowest dose. If efficacy is not achieved with the lowest dose, discontinue use.

2.3 Non-Interchangeability with Other Formulations of Diclofenac

ZORVOLEX capsules are not interchangeable with other formulations of oral diclofenac even if the milligram strength is the same. ZORVOLEX capsules contain diclofenac free acid whereas other diclofenac products contain a salt of diclofenac, i.e., diclofenac potassium or sodium. A 35 mg dose of ZORVOLEX is approximately equal to 37.6 mg of sodium diclofenac or 39.5 mg of potassium diclofenac. Therefore, do not substitute similar dosing strengths of other diclofenac products without taking this into consideration.

3 DOSAGE FORMS AND STRENGTHS

ZORVOLEX (diclofenac) capsules 18 mg - blue body and light green cap (imprinted IP-203 on the body and 18 mg on the cap in white ink).

ZORVOLEX (diclofenac) capsules 35 mg - blue body and green cap (imprinted IP-204 on the body and 35 mg on the cap in white ink).

4 CONTRAINDICATIONS

ZORVOLEX is contraindicated in patients with:

- known hypersensitivity (e.g., anaphylactoid reactions and serious skin reactions) to diclofenac or any components of the drug product [*see Warnings and Precautions (5.7, 5.8)*].
- a history of asthma, urticaria, or other allergic-type reactions after taking aspirin or other NSAIDs. Severe, rarely fatal, anaphylactic-like reactions to NSAIDs have been reported in such patients [*see Warnings and Precautions (5.7, 5.13)*].
- perioperative pain in the setting of coronary artery bypass graft (CABG) surgery [*see Warnings and Precautions (5.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Cardiovascular Thrombotic Events

Clinical studies of several COX-2 selective and nonselective NSAIDs of up to three years duration have shown an increased risk of serious cardiovascular (CV) thrombotic events, myocardial infarction, and stroke, which can be fatal. All NSAIDs, both COX-2 selective and nonselective, may have a similar risk. Patients with known CV disease or risk factors for

CV disease may be at greater risk. To minimize the potential risk for an adverse CV event in patients treated with an NSAID, use the lowest effective dose for the shortest duration possible. Physicians and patients should remain alert for the development of such events, even in the absence of previous CV symptoms. Inform patients about the signs and/or symptoms of serious CV events and the steps to take if they occur.

Two large, controlled, clinical studies of a COX-2 selective NSAID for the treatment of pain in the first 10-14 days following CABG surgery found an increased incidence of myocardial infarction and stroke [*see Contraindications (4)*].

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concurrent use of aspirin and an NSAID, such as diclofenac, does increase the risk of serious GI events [*see Warnings and Precautions (5.2)*, *Drug Interactions (7.1)*].

5.2 Gastrointestinal (GI) Effects – Risk of GI Ulceration, Bleeding, and Perforation

NSAIDs, including ZORVOLEX, can cause serious gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration, and perforation of the stomach, small intestine, or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one in five patients, who develop a serious upper GI adverse event on NSAID therapy, is symptomatic. Upper GI ulcers, gross bleeding or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3-6 months, and in about 2%-4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of developing a serious GI event at some time during the course of therapy. However, even short-term NSAID therapy is not without risk.

Prescribe NSAIDs, including ZORVOLEX, with extreme caution in patients with a prior history of ulcer disease or gastrointestinal bleeding. Patients with a prior history of peptic ulcer disease and/or gastrointestinal bleeding who use NSAIDs have a greater than 10-fold increased risk for developing a GI bleed compared to patients with neither of these risk factors. Other factors that increase the risk for GI bleeding in patients treated with NSAIDs include concomitant use of oral corticosteroids or anticoagulants, longer duration of NSAID therapy, smoking, use of alcohol, older age, and poor general health status. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and therefore, special care should be taken in treating this population.

To minimize the potential risk for an adverse GI event in patients treated with an NSAID, use the lowest effective dose for the shortest possible duration. Patients and physicians should remain alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy and promptly initiate additional evaluation and treatment if a serious GI adverse event is suspected. This should include discontinuation of the NSAID until a serious GI adverse event is ruled out. For high risk patients, alternative therapies that do not include NSAIDs should be considered.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.