

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use IMBRUVICA safely and effectively. See full prescribing information for IMBRUVICA.

IMBRUVICA® (ibrutinib) capsules, for oral use
IMBRUVICA® (ibrutinib) tablets, for oral use
Initial U.S. Approval: 2013

RECENT MAJOR CHANGES

Dosage and Administration (2.1, 2.2, 2.3, 2.4, 2.5, 2.6) 08/2018
Warnings and Precautions (5) 08/2018

INDICATIONS AND USAGE

IMBRUVICA is a kinase inhibitor indicated for the treatment of adult patients with:

- Mantle cell lymphoma (MCL) who have received at least one prior therapy (1.1). Accelerated approval was granted for this indication based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.
- Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) (1.2).
- Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion (1.3).
- Waldenström’s macroglobulinemia (WM) (1.4).
- Marginal zone lymphoma (MZL) who require systemic therapy and have received at least one prior anti-CD20-based therapy (1.5). Accelerated approval was granted for this indication based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.
- Chronic graft versus host disease (cGVHD) after failure of one or more lines of systemic therapy (1.6).

DOSAGE AND ADMINISTRATION

- MCL and MZL: 560 mg taken orally once daily (2.2).
- CLL/SLL, WM, and cGVHD: 420 mg taken orally once daily (2.2).

Dose should be taken orally with a glass of water. Do not open, break, or chew the capsules. Do not cut, crush, or chew the tablets (2.1).

DOSAGE FORMS AND STRENGTHS

Capsules: 70 mg and 140 mg (3)
Tablets: 140 mg, 280 mg, 420 mg, and 560 mg (3)

CONTRAINDICATIONS

None (4)

WARNINGS AND PRECAUTIONS

- Hemorrhage: Monitor for bleeding and manage (5.1).
- Infections: Monitor patients for fever and infections, evaluate promptly, and treat (5.2).
- Cytopenias: Check complete blood counts monthly (5.3).
- Cardiac arrhythmias: Monitor for symptoms of arrhythmias and manage (5.4).
- Hypertension: Monitor blood pressure and treat (5.5).
- Second Primary Malignancies: Other malignancies have occurred in patients, including skin cancers, and other carcinomas (5.6).
- Tumor Lysis Syndrome (TLS): Assess baseline risk and take precautions. Monitor and treat for TLS (5.7).
- Embryo-Fetal Toxicity: Can cause fetal harm. Advise women of the potential risk to a fetus and to avoid pregnancy while taking the drug and for 1 month after cessation of therapy. Advise men to avoid fathering a child during the same time period (5.8, 8.3).

ADVERSE REACTIONS

The most common adverse reactions (≥20%) in patients with B-cell malignancies (MCL, CLL/SLL, WM and MZL) were neutropenia, thrombocytopenia, diarrhea, anemia, musculoskeletal pain, rash, nausea, bruising, fatigue, hemorrhage, and pyrexia (6).

The most common adverse reactions (≥20%) in patients with cGVHD were fatigue, bruising, diarrhea, thrombocytopenia, muscle spasms, stomatitis, nausea, hemorrhage, anemia, and pneumonia (6).

To report SUSPECTED ADVERSE REACTIONS, contact Pharmacovigilance at 1-877-877-3536 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- CYP3A Inhibitors: Dose adjustments may be recommended (2.4, 7.1).
- CYP3A Inducers: Avoid coadministration with strong CYP3A inducers (7.2).

USE IN SPECIFIC POPULATIONS

Hepatic Impairment (based on Child-Pugh criteria): Avoid use of IMBRUVICA in patients with severe baseline hepatic impairment. In patients with mild or moderate impairment, reduce IMBRUVICA dose (2.5, 8.6).

See 17 for PATIENT COUNSELING INFORMATION and FDA approved patient labeling.

Revised: 08/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Mantle Cell Lymphoma
- 1.2 Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma
- 1.3 Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma with 17p deletion
- 1.4 Waldenström’s Macroglobulinemia
- 1.5 Marginal Zone Lymphoma
- 1.6 Chronic Graft versus Host Disease

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing Guidelines
- 2.2 Recommended Dosage
- 2.3 Dose Modifications for Adverse Reactions
- 2.4 Dose Modifications for Use with CYP3A Inhibitors
- 2.5 Dose Modifications for Use in Hepatic Impairment
- 2.6 Missed Dose

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hemorrhage
- 5.2 Infections
- 5.3 Cytopenias
- 5.4 Cardiac Arrhythmias
- 5.5 Hypertension
- 5.6 Second Primary Malignancies
- 5.7 Tumor Lysis Syndrome
- 5.8 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Effect of CYP3A Inhibitors on Ibrutinib
- 7.2 Effect of CYP3A Inducers on Ibrutinib

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment
- 8.7 Plasmapheresis

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Mantle Cell Lymphoma
- 14.2 Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma
- 14.3 Waldenström’s Macroglobulinemia
- 14.4 Marginal Zone Lymphoma
- 14.5 Chronic Graft versus Host Disease

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Mantle Cell Lymphoma

IMBRUVICA is indicated for the treatment of adult patients with mantle cell lymphoma (MCL) who have received at least one prior therapy.

Accelerated approval was granted for this indication based on overall response rate. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial [*see Clinical Studies (14.1)*].

1.2 Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma

IMBRUVICA is indicated for the treatment of adult patients with chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL).

1.3 Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma with 17p deletion

IMBRUVICA is indicated for the treatment of adult patients with chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL) with 17p deletion.

1.4 Waldenström's Macroglobulinemia

IMBRUVICA is indicated for the treatment of adult patients with Waldenström's macroglobulinemia (WM).

1.5 Marginal Zone Lymphoma

IMBRUVICA is indicated for the treatment of adult patients with marginal zone lymphoma (MZL) who require systemic therapy and have received at least one prior anti-CD20-based therapy.

Accelerated approval was granted for this indication based on overall response rate [*see Clinical Studies (14.4)*]. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

1.6 Chronic Graft versus Host Disease

IMBRUVICA is indicated for the treatment of adult patients with chronic graft-versus-host disease (cGVHD) after failure of one or more lines of systemic therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Guidelines

Administer IMBRUVICA orally once daily at approximately the same time each day. The dose should be taken orally with a glass of water. Do not open, break, or chew the capsules. Do not cut, crush, or chew the tablets.

2.2 Recommended Dosage

Mantle Cell Lymphoma and Marginal Zone Lymphoma

The recommended dose of IMBRUVICA for MCL and MZL is 560 mg orally once daily until disease progression or unacceptable toxicity.

Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma and Waldenström's Macroglobulinemia

The recommended dose of IMBRUVICA for CLL/SLL and WM as a single agent, in combination with rituximab for WM, or in combination with bendamustine and rituximab for CLL/SLL is 420 mg orally once daily until disease progression or unacceptable toxicity.

When administering IMBRUVICA in combination with rituximab, consider administering IMBRUVICA prior to rituximab when given on the same day.

Chronic Graft versus Host Disease

The recommended dose of IMBRUVICA for cGVHD is 420 mg orally once daily until cGVHD progression, recurrence of an underlying malignancy, or unacceptable toxicity. When a patient no longer requires therapy for the treatment of cGVHD, IMBRUVICA should be discontinued considering the medical assessment of the individual patient.

2.3 Dose Modifications for Adverse Reactions

Interrupt IMBRUVICA therapy for any Grade 3 or greater non-hematological toxicities, Grade 3 or greater neutropenia with infection or fever, or Grade 4 hematological toxicities. Once the symptoms of the toxicity have resolved to Grade 1 or baseline (recovery), IMBRUVICA therapy may be reinitiated at the starting dose. If the toxicity reoccurs, reduce dose by 140 mg per day. A second reduction of dose by 140 mg may be considered as needed. If these toxicities persist or recur following two dose reductions, discontinue IMBRUVICA.

Recommended dose modifications are described below:

Toxicity Occurrence	Dose Modification for MCL and MZL After Recovery Starting Dose = 560 mg	Dose Modification for CLL/SLL, WM, and cGVHD After Recovery Starting Dose = 420 mg
First	Restart at 560 mg daily	Restart at 420 mg daily
Second	Restart at 420 mg daily	Restart at 280 mg daily
Third	Restart at 280 mg daily	Restart at 140 mg daily
Fourth	Discontinue IMBRUVICA	Discontinue IMBRUVICA

2.4 Dose Modifications for Use with CYP3A Inhibitors

Recommended dose modifications are described below [see *Drug Interactions (7.1)*]:

Patient Population	Coadministered Drug	Recommended IMBRUVICA Dose
B-Cell Malignancies	<ul style="list-style-type: none"> Moderate CYP3A inhibitor 	280 mg once daily Modify dose as recommended [see <i>Dosage and Administration (2.3)</i>].
	<ul style="list-style-type: none"> Voriconazole 200 mg twice daily Posaconazole suspension 100 mg once daily, 100 mg twice daily, or 200 mg twice daily 	140 mg once daily Modify dose as recommended [see <i>Dosage and Administration (2.3)</i>].
	<ul style="list-style-type: none"> Posaconazole suspension 200 mg three times daily or 400 mg twice daily Posaconazole IV injection 300 mg once daily Posaconazole delayed-release tablets 300 mg once daily 	70 mg once daily Interrupt dose as recommended [see <i>Dosage and Administration (2.3)</i>].
	<ul style="list-style-type: none"> Other strong CYP3A inhibitors 	Avoid concomitant use. If these inhibitors will be used short-term (such as anti-infectives for seven days or less), interrupt IMBRUVICA.
Chronic Graft versus Host Disease	<ul style="list-style-type: none"> Moderate CYP3A inhibitor 	420 mg once daily Modify dose as recommended [see <i>Dosage and Administration (2.3)</i>].
	<ul style="list-style-type: none"> Voriconazole 200 mg twice daily Posaconazole suspension 100 mg once daily, 100 mg twice daily, or 200 mg twice daily 	280 mg once daily Modify dose as recommended [see <i>Dosage and Administration (2.3)</i>].
	<ul style="list-style-type: none"> Posaconazole suspension 200 mg three times daily or 400 mg twice daily Posaconazole IV injection 300 mg once daily Posaconazole delayed-release tablets 300 mg once daily 	140 mg once daily Interrupt dose as recommended [see <i>Dosage and Administration (2.3)</i>].
	<ul style="list-style-type: none"> Other strong CYP3A inhibitors 	Avoid concomitant use. If these inhibitors will be used short-term (such as anti-infectives for seven days or less), interrupt IMBRUVICA.

After discontinuation of a CYP3A inhibitor, resume previous dose of IMBRUVICA [see *Dosage and Administration (2.2)* and *Drug Interactions (7.1)*].

2.5 Dose Modifications for Use in Hepatic Impairment

The recommended dose is 140 mg daily for patients with mild hepatic impairment (Child-Pugh class A).

The recommended dose is 70 mg daily for patients with moderate hepatic impairment (Child-Pugh class B).

Avoid the use of IMBRUVICA in patients with severe hepatic impairment (Child-Pugh class C) [see *Use in Specific Populations (8.6) and Clinical Pharmacology (12.3)*].

2.6 Missed Dose

If a dose of IMBRUVICA is not taken at the scheduled time, it can be taken as soon as possible on the same day with a return to the normal schedule the following day. Extra doses of IMBRUVICA should not be taken to make up for the missed dose.

3 DOSAGE FORMS AND STRENGTHS

Capsules:

Each 70 mg capsule is a yellow, opaque capsule marked with “ibr 70 mg” in black ink.

Each 140 mg capsule is a white, opaque capsule marked with “ibr 140 mg” in black ink.

Tablets:

Each 140 mg tablet is a yellow green to green round tablet debossed with “ibr” on one side and “140” on the other side.

Each 280 mg tablet is a purple oblong tablet debossed with “ibr” on one side and “280” on the other side.

Each 420 mg tablet is a yellow green to green oblong tablet debossed with “ibr” on one side and “420” on the other side.

Each 560 mg tablet is a yellow to orange oblong tablet debossed with “ibr” on one side and “560” on the other side.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Hemorrhage

Fatal bleeding events have occurred in patients treated with IMBRUVICA. Grade 3 or higher bleeding events (intracranial hemorrhage [including subdural hematoma], gastrointestinal bleeding, hematuria, and post procedural hemorrhage) have occurred in 3% of patients, with fatalities occurring in 0.3% of 1,011 patients exposed to IMBRUVICA in clinical trials. Bleeding events of any grade, including bruising and petechiae, occurred in 44% of patients treated with IMBRUVICA.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.