

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

214120Orig1s000

LABELING

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ONUREG safely and effectively. See full prescribing information for ONUREG.

ONUREG (azacitidine) tablets, for oral use
Initial U.S. Approval: 2004

INDICATIONS AND USAGE

ONUREG is a nucleoside metabolic inhibitor indicated for continued treatment of adult patients with acute myeloid leukemia who achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and are not able to complete intensive curative therapy (1).

DOSAGE AND ADMINISTRATION

- Do not substitute ONUREG for intravenous or subcutaneous azacitidine. The indications and dosing regimen for ONUREG differ from that of intravenous or subcutaneous azacitidine (2.1, 5.1).
- Administer ONUREG 300 mg orally once daily on Days 1 through 14 of each 28-day cycle (2.2).
- Administer an antiemetic before each dose for at least the first 2 cycles (2.2).

DOSAGE FORMS AND STRENGTHS

Tablets: 200 mg and 300 mg (3).

CONTRAINDICATIONS

History of severe hypersensitivity to azacitidine or its components (4).

WARNINGS AND PRECAUTIONS

- **Risks of Substitution with Other Azacitidine Products:** Do not substitute ONUREG for intravenous or subcutaneous azacitidine (2.1, 5.1).
- **Myelosuppression:** Monitor complete blood counts every other week for the first 2 cycles and prior to the start of each cycle thereafter. Increase monitoring to every other week for the 2 cycles after any dose reduction. Withhold and then resume at same or reduced dose or discontinue ONUREG based on severity (2.3, 5.2).
- **Embryo-Fetal Toxicity:** Can cause fetal harm. Advise patients of the potential risk to a fetus and use of effective contraception (5.4, 8.1, 8.3).

ADVERSE REACTIONS

The most common adverse reactions ($\geq 10\%$) are nausea, vomiting, diarrhea, fatigue/asthenia, constipation, pneumonia, abdominal pain, arthralgia, decreased appetite, febrile neutropenia, dizziness, and pain in extremity.

To report SUSPECTED ADVERSE REACTIONS, contact Celgene Corporation at 1-888-423-5436 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

Lactation: Advise not to breastfeed (8.2).

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling

Revised: 9/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE		
2	DOSAGE AND ADMINISTRATION		
	2.1	Important Administration Information	
	2.2	Recommended Dosage	
	2.3	Monitoring and Dosage Modifications for Adverse Reactions	
3	DOSAGE FORMS AND STRENGTHS		
4	CONTRAINDICATIONS		
5	WARNINGS AND PRECAUTIONS		
	5.1	Risks of Substitution with Other Azacitidine Products	
	5.2	Myelosuppression	
	5.3	Increased Early Mortality in Patients with Myelodysplastic Syndromes	
	5.4	Embryo-Fetal Toxicity	
6	ADVERSE REACTIONS		
	6.1	Clinical Trials Experience	
	6.2	Postmarketing Experience	
8	USE IN SPECIFIC POPULATIONS		
	8.1	Pregnancy	
	8.2	Lactation	
	8.3	Females and Males of Reproductive Potential	
	8.4	Pediatric Use	
	8.5	Geriatric Use	
	8.6	Renal Impairment	
	8.7	Hepatic Impairment	
11	DESCRIPTION		
12	CLINICAL PHARMACOLOGY		
	12.1	Mechanism of Action	
	12.2	Pharmacodynamics	
	12.3	Pharmacokinetics	
13	NONCLINICAL TOXICOLOGY		
	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility	
14	CLINICAL STUDIES		
15	REFERENCES		
16	HOW SUPPLIED/STORAGE AND HANDLING		
17	PATIENT COUNSELING INFORMATION		

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ONUREG is indicated for continued treatment of adult patients with acute myeloid leukemia who achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and are not able to complete intensive curative therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Information

Do not substitute ONUREG for intravenous or subcutaneous azacitidine. The indications and dosing regimen for ONUREG differ from that of intravenous or subcutaneous azacitidine [see Warnings and Precautions (5.1)].

2.2 Recommended Dosage

The recommended dosage of ONUREG is 300 mg orally once daily with or without food on Days 1 through 14 of each 28-day cycle. Continue ONUREG until disease progression or unacceptable toxicity.

Administer an antiemetic 30 minutes prior to each dose of ONUREG for the first 2 cycles. Antiemetic prophylaxis may be omitted after 2 cycles if there has been no nausea and vomiting.

If the absolute neutrophil count (ANC) is less than 0.5 Gi/L on Day 1 of a cycle, do not administer ONUREG. Delay the start of the cycle until the ANC is 0.5 Gi/L or more.

Instruct patients on the following:

- Do not split, crush, or chew ONUREG tablets.
- Take a dose about the same time each day.
- If a dose of ONUREG is missed, or not taken at the usual time, take the dose as soon as possible on the same day, and resume the normal schedule the following day. Do not take 2 doses on the same day.
- If a dose is vomited, do not take another dose on the same day. Resume the normal schedule the following day.

ONUREG is a hazardous drug. Follow applicable special handling and disposal procedures.¹

2.3 Monitoring and Dosage Modifications for Adverse Reactions

Monitor complete blood count every other week for the first 2 cycles and prior to the start of each cycle thereafter. Increase monitoring to every other week for the 2 cycles after any dose reduction for myelosuppression.

The recommended dosage modifications for adverse reactions are provided in Table 1.

Table 1: Recommended Dosage Modifications for Adverse Reactions

Adverse Reaction	Severity	Recommended Dosage Modification
Myelosuppression <i>[see Warnings and Precautions (5.2)]</i>	Neutrophils less than 0.5 Gi/L on Cycle Day 1	<ul style="list-style-type: none"> Interrupt treatment. Resume at the same dose once neutrophils return to 0.5 Gi/L or higher.
	Neutrophils less than 1 Gi/L with fever at anytime	<p>First Occurrence</p> <ul style="list-style-type: none"> Interrupt treatment. Resume at the same dose once neutrophils return to 1 Gi/L or higher. <p>Occurrence in 2 Consecutive Cycles</p> <ul style="list-style-type: none"> Interrupt treatment. After neutrophils return to 1 Gi/L or higher, resume at reduced dose of 200 mg. If a patient continues to experience febrile neutropenia after dose reduction, reduce the treatment duration by 7 days. If febrile neutropenia reoccurs after dose and schedule reduction, discontinue ONUREG.
	Platelets less than 50 Gi/L with bleeding	<p>First Occurrence</p> <ul style="list-style-type: none"> Interrupt dose. Resume at the same dose once platelets return to 50 Gi/L or higher. <p>Occurrence in 2 Consecutive Cycles</p> <ul style="list-style-type: none"> Interrupt dose. After platelets return to 50 Gi/L or higher, resume at reduced dose of 200 mg. If a patient continues to experience thrombocytopenia with bleeding after dose reduction, reduce the treatment duration by 7 days. If thrombocytopenia with bleeding reoccurs after dose and schedule reduction, discontinue ONUREG.

Adverse Reaction	Severity	Recommended Dosage Modification
Gastrointestinal Toxicity [<i>see Adverse Reactions (6.1)</i>]	Grade 3 or 4 Nausea or Vomiting	<ul style="list-style-type: none"> • Interrupt dose. Resume at the same dose once toxicity has resolved to Grade 1 or lower. • If toxicity reoccurs, interrupt dose until resolved to Grade 1 or lower. Resume at reduced dose of 200 mg. • If a patient continues to experience the toxicity after dose reduction, reduce the treatment duration by 7 days. • If the toxicity continues or reoccurs after dose and schedule reduction, discontinue ONUREG.
	Grade 3 or 4 Diarrhea	<ul style="list-style-type: none"> • Interrupt dose. Resume at the same dose once toxicity has resolved to Grade 1 or lower. • If toxicity reoccurs, interrupt dose until resolved to Grade 1 or lower. Resume at reduced dose of 200 mg. • If a patient continues to experience the toxicity after dose reduction, reduce the treatment duration by 7 days. • If the toxicity continues or reoccurs after dose and schedule reduction, discontinue ONUREG.
Other Adverse Reactions [<i>see Adverse Reactions (6.1)</i>]	Grade 3 or 4	<ul style="list-style-type: none"> • Interrupt dose and provide medical support. Resume at the same dose once toxicity has resolved to Grade 1 or lower. • If toxicity re-occurs, interrupt dose until resolved to Grade 1 or lower. Resume at reduced dose of 200 mg. • If a patient continues to experience the toxicity after dose reduction, reduce the treatment duration by 7 days. • If the toxicity continues or reoccurs after dose and schedule reduction, discontinue ONUREG.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.