

EXHIBIT 3

MOLECULAR BIOLOGY OF THE CELL

fourth edition

Bruce Alberts

Alexander Johnson

Julian Lewis

Martin Raff

Keith Roberts

Peter Walter

 **Garland Science**
Taylor & Francis Group

APPX 0099

Garland

Vice President: Denise Schanck
Managing Editor: Sarah Gibbs
Senior Editorial Assistant: Kirsten Jenner
Managing Production Editor: Emma Hunt
Proofreader and Layout: Emma Hunt
Production Assistant: Angela Bennett
Text Editors: Marjorie Singer Anderson and Betsy Dilerenia
Copy Editor: Bruce Goatly
Word Processors: Fran Dependahl, Misty Landers and Carol Winter
Designer: Blink Studio, London
Illustrator: Nigel Orme
Indexer: Janine Ross and Sherry Granum
Manufacturing: Nigel Eyre and Marion Morrow

Cell Biology Interactive

Artistic and Scientific Direction: Peter Walter
Narrated by: Julie Theriot
Production, Design, and Development: Mike Morales

Bruce Alberts received his Ph.D. from Harvard University and is President of the National Academy of Sciences and Professor of Biochemistry and Biophysics at the University of California, San Francisco. **Alexander Johnson** received his Ph.D. from Harvard University and is a Professor of Microbiology and Immunology at the University of California, San Francisco. **Julian Lewis** received his D.Phil. from the University of Oxford and is a Principal Scientist at the Imperial Cancer Research Fund, London. **Martin Raff** received his M.D. from McGill University and is at the Medical Research Council Laboratory for Molecular Cell Biology and Cell Biology Unit and in the Biology Department at University College London. **Keith Roberts** received his Ph.D. from the University of Cambridge and is Associate Research Director at the John Innes Centre, Norwich. **Peter Walter** received his Ph.D. from The Rockefeller University in New York and is Professor and Chairman of the Department of Biochemistry and Biophysics at the University of California, San Francisco, and an Investigator of the Howard Hughes Medical Institute.

© 2002 by Bruce Alberts, Alexander Johnson, Julian Lewis, Martin Raff, Keith Roberts, and Peter Walter.
© 1983, 1989, 1994 by Bruce Alberts, Dennis Bray, Julian Lewis, Martin Raff, Keith Roberts, and James D. Watson.

All rights reserved. No part of this book covered by the copyright hereon may be reproduced or used in any format in any form or by any means—graphic, electronic, or mechanical, including photocopying, recording, taping, or information storage and retrieval systems—without permission of the publisher.

Library of Congress Cataloging-in-Publication Data

Molecular biology of the cell / Bruce Alberts ... [et al.].-- 4th ed.
p. cm
Includes bibliographical references and index.
ISBN 0-8153-3218-1 (hardbound) -- ISBN 0-8153-4072-9 (pbk.)
1. Cytology. 2. Molecular biology. I. Alberts, Bruce.
[DNLM: 1. Cells. 2. Molecular Biology.]
QH581.2 .M64 2002
571.6--dc21

2001054471 CIP

Published by Garland Science, a member of the Taylor & Francis Group,
29 West 35th Street, New York, NY 10001-2299

Printed in the United States of America

15 14 13 12 11 10 9 8 7 6 5 4 3 2 1

Front cover Human Genome: Reprinted by permission from *Nature*, International Human Genome Sequencing Consortium, 409:860–921, 2001 © Macmillan Magazines Ltd. Adapted from an image by Francis Collins, NHGRI; Jim Kent, UCSC; Ewan Birney, EBI; and Darryl Leja, NHGRI; showing a portion of Chromosome 1 from the initial sequencing of the human genome.

Back cover In 1967, the British artist Peter Blake created a design classic. Nearly 35 years later Nigel Orme (illustrator), Richard Denyer (photographer), and the authors have together produced an affectionate tribute to Mr Blake's image. With its gallery of icons and influences, its assembly created almost as much complexity, intrigue and mystery as the original. *Drosophila*, *Arabidopsis*, Dolly and the assembled company tempt you to dip inside where, as in the original, "a splendid time is guaranteed for all." (Gunter Blobel, courtesy of The Rockefeller University; Marie Curie, Keystone Press Agency Inc; Darwin bust, by permission of the President and Council of the Royal Society; Rosalind Franklin, courtesy of Cold Spring Harbor Laboratory Archives; Dorothy Hodgkin, © The Nobel Foundation, 1964; James Joyce, etching by Peter Blake; Robert Johnson, photo booth self-portrait early 1930s, © 1986 Delta Haze Corporation all rights reserved, used by permission; Albert L. Lehninger, (unidentified photographer) courtesy of The Alan Mason Chesney Medical Archives of The Johns Hopkins Medical Institutions; Linus Pauling, from Ava Helen and Linus Pauling Papers, Special Collections, Oregon State University; Nicholas Poussin, courtesy of ArtToday.com; Barbara McClintock, © David Micklos, 1983; Andrei Sakharov, courtesy of Elena Bonner; Frederick Sanger, © The Nobel Foundation, 1958.)

3

PROTEINS

THE SHAPE AND STRUCTURE OF
PROTEINS

PROTEIN FUNCTION

When we look at a cell through a microscope or analyze its electrical or biochemical activity, we are, in essence, observing proteins. Proteins constitute most of a cell's dry mass. They are not only the building blocks from which cells are built; they also execute nearly all cell functions. Thus, enzymes provide the intricate molecular surfaces in a cell that promote its many chemical reactions. Proteins embedded in the plasma membrane form channels and pumps that control the passage of small molecules into and out of the cell. Other proteins carry messages from one cell to another, or act as signal integrators that relay sets of signals inward from the plasma membrane to the cell nucleus. Yet others serve as tiny molecular machines with moving parts: *kinesin*, for example, propels organelles through the cytoplasm; *topoisomerase* can untangle knotted DNA molecules. Other specialized proteins act as antibodies, toxins, hormones, antifreeze molecules, elastic fibers, ropes, or sources of luminescence. Before we can hope to understand how genes work, how muscles contract, how nerves conduct electricity, how embryos develop, or how our bodies function, we must attain a deep understanding of proteins.

THE SHAPE AND STRUCTURE OF PROTEINS

From a chemical point of view, proteins are by far the most structurally complex and functionally sophisticated molecules known. This is perhaps not surprising, once one realizes that the structure and chemistry of each protein has been developed and fine-tuned over billions of years of evolutionary history. We start this chapter by considering how the location of each amino acid in the long string of amino acids that forms a protein determines its three-dimensional shape. We will then use this understanding of protein structure at the atomic level to describe how the precise shape of each protein molecule determines its function in a cell.

The Shape of a Protein Is Specified by Its Amino Acid Sequence

Recall from Chapter 2 that there are 20 types of amino acids in proteins, each with different chemical properties. A **protein** molecule is made from a long chain of these amino acids, each linked to its neighbor through a covalent peptide bond (Figure 3-1). Proteins are therefore also known as *polypeptides*. Each type of protein has a unique sequence of amino acids, exactly the same from one molecule to the next. Many thousands of different proteins are known, each with its own particular amino acid sequence.




Figure 3-1 A peptide bond. This covalent bond forms when the carbon atom from the carboxyl group of one amino acid shares electrons with the nitrogen atom (blue) from the amino group of a second amino acid. As indicated, a molecule of water is lost in this condensation reaction.

The repeating sequence of atoms along the core of the polypeptide chain is referred to as the **polypeptide backbone**. Attached to this repetitive chain are those portions of the amino acids that are not involved in making a peptide bond and which give each amino acid its unique properties: the 20 different amino acid **side chains** (Figure 3-2). Some of these side chains are nonpolar and hydrophobic (“water-fearing”), others are negatively or positively charged, some are reactive, and so on. Their atomic structures are presented in Panel 3-1, and a brief list with abbreviations is provided in Figure 3-3.

As discussed in Chapter 2, atoms behave almost as if they were hard spheres with a definite radius (their *van der Waals radius*). The requirement that no two atoms overlap limits greatly the possible bond angles in a polypeptide chain (Figure 3-4). This constraint and other steric interactions severely restrict the variety of three-dimensional arrangements of atoms (or *conformations*) that are possible. Nevertheless, a long flexible chain, such as a protein, can still fold in an enormous number of ways.

The folding of a protein chain is, however, further constrained by many different sets of weak *noncovalent bonds* that form between one part of the chain and another. These involve atoms in the polypeptide backbone, as well as atoms in the amino acid side chains. The weak bonds are of three types: *hydrogen bonds*, *ionic bonds*, and *van der Waals attractions*, as explained in Chapter 2 (see p. 57). Individual noncovalent bonds are 30–300 times weaker than the typical covalent bonds that create biological molecules. But many weak bonds can act in parallel to hold two regions of a polypeptide chain tightly together. The stability of each folded shape is therefore determined by the combined strength of large numbers of such noncovalent bonds (Figure 3-5).

A fourth weak force also has a central role in determining the shape of a protein. As described in Chapter 2, hydrophobic molecules, including the nonpolar side chains of particular amino acids, tend to be forced together in an aqueous environment in order to minimize their disruptive effect on the hydrogen-bonded network of water molecules (see p. 58 and Panel 2-2, pp. 112–113). Therefore, an important factor governing the folding of any protein is the distribution of its polar and nonpolar amino acids. The nonpolar (hydrophobic) side chains in a protein—belonging to such amino acids as phenylalanine, leucine, valine, and tryptophan—tend to cluster in the interior of the molecule (just as hydrophobic oil droplets coalesce in water to form one large droplet). This enables them to




Figure 3-2 The structural components of a protein. A protein consists of a polypeptide backbone with attached side chains. Each type of protein differs in its sequence and number of amino acids; therefore, it is the sequence of the chemically different side chains that makes each protein distinct. The two ends of a polypeptide chain are chemically different: the end carrying the free amino group (NH_3^+ , also written NH_2) is the amino terminus, or N-terminus, and that carrying the free carboxyl group (COO^- , also written COOH) is the carboxyl terminus or C-terminus. The amino acid sequence of a protein is always presented in the N-to-C direction, reading from left to right.

AMINO ACID	SIDE CHAIN	AMINO ACID	SIDE CHAIN
Aspartic acid	Asp D negative	Alanine	Ala A nonpolar
Glutamic acid	Glu E negative	Glycine	Gly G nonpolar
Arginine	Arg R positive	Valine	Val V nonpolar
Lysine	Lys K positive	Leucine	Leu L nonpolar
Histidine	His H positive	Isoleucine	Ile I nonpolar
Asparagine	Asn N uncharged polar	Proline	Pro P nonpolar
Glutamine	Gln Q uncharged polar	Phenylalanine	Phe F nonpolar
Serine	Ser S uncharged polar	Methionine	Met M nonpolar
Threonine	Thr T uncharged polar	Tryptophan	Trp W nonpolar
Tyrosine	Tyr Y uncharged polar	Cysteine	Cys C nonpolar


┌────────── POLAR AMINO ACIDS ─────────┐ ┌────────── NONPOLAR AMINO ACIDS ─────────┐

Figure 3-3 The 20 amino acids found in proteins. Both three-letter and one-letter abbreviations are listed. As shown, there are equal numbers of polar and nonpolar side chains. For their atomic structures, see Panel 3-1 (pp. 132-133).


PANEL 3-1 The 20 Amino Acids Found in Proteins

THE AMINO ACID

The general formula of an amino acid is




R is commonly one of 20 different side chains. At pH 7 both the amino and carboxyl groups are ionized.



OPTICAL ISOMERS

The α -carbon atom is asymmetric, which allows for two mirror image (or stereo-) isomers, L and D.



Proteins consist exclusively of L-amino acids.

FAMILIES OF AMINO ACIDS

The common amino acids are grouped according to whether their side chains are


- acidic
- basic
- uncharged polar
- nonpolar

These 20 amino acids are given both three-letter and one-letter abbreviations.

Thus: alanine = Ala = A


BASIC SIDE CHAINS

lysine
(Lys, or K)




This group is very basic because its positive charge is stabilized by resonance.

arginine
(Arg, or R)



histidine
(His, or H)




These nitrogens have a relatively weak affinity for an H+ and are only partly positive at neutral pH.


PEPTIDE BONDS

Amino acids are commonly joined together by an amide linkage, called a peptide bond.

Peptide bond: The four atoms in each gray box form a rigid planar unit. There is no rotation around the C-N bond.



Proteins are long polymers of amino acids linked by peptide bonds, and they are always written with the N-terminus toward the left. The sequence of this tripeptide is histidine-cysteine-valine.




These two single bonds allow rotation, so that long chains of amino acids are very flexible.

ACIDIC SIDE CHAINS


aspartic acid

(Asp, or D)



glutamic acid


(Glu, or E)



UNCHARGED POLAR SIDE CHAINS


asparagine

(Asn, or N)



glutamine


(Gln, or Q)



Although the amide N is not charged at neutral pH, it is polar.


serine

(Ser, or S)




threonine

(Thr, or T)



tyrosine

(Tyr, or Y)




The -OH group is polar.

NONPOLAR SIDE CHAINS


alanine

(Ala, or A)




valine

(Val, or V)




leucine

(Leu, or L)




isoleucine

(Ile, or I)




proline

(Pro, or P)




phenylalanine

(Phe, or F)




methionine

(Met, or M)




tryptophan

(Trp, or W)




glycine

(Gly, or G)




cysteine

(Cys, or C)



Disulfide bonds can form between two cysteine side chains in proteins.







Figure 3-4 Steric limitations on the bond angles in a polypeptide chain. (A) Each amino acid contributes three bonds (red) to the backbone of the chain. The peptide bond is planar (gray shading) and does not permit rotation. By contrast, rotation can occur about the C_{α} -C bond, whose angle of rotation is called psi (ψ), and about the N- C_{α} bond, whose angle of rotation is called phi (ϕ). By convention, an R group is often used to denote an amino acid side chain (green circles). (B) The conformation of the main-chain atoms in a protein is determined by one pair of ϕ and ψ angles for each amino acid; because of steric collisions between atoms within each amino acid, most pairs of ϕ and ψ angles do not occur. In this so-called Ramachandran plot, each dot represents an observed pair of angles in a protein. (B, from J. Richardson, *Adv. Prot. Chem.* 34:174-175, 1981. © Academic Press.)

avoid contact with the water that surrounds them inside a cell. In contrast, polar side chains—such as those belonging to arginine, glutamine, and histidine—tend to arrange themselves near the outside of the molecule, where they can form hydrogen bonds with water and with other polar molecules (Figure 3-6). When polar amino acids are buried within the protein, they are usually hydrogen-bonded to other polar amino acids or to the polypeptide backbone (Figure 3-7).

Proteins Fold into a Conformation of Lowest Energy

As a result of all of these interactions, each type of protein has a particular three-dimensional structure, which is determined by the order of the amino acids in its chain. The final folded structure, or **conformation**, adopted by any polypeptide chain is generally the one in which the free energy is minimized. Protein folding has been studied in a test tube by using highly purified proteins. A protein can be unfolded, or *denatured*, by treatment with certain solvents, which disrupt the noncovalent interactions holding the folded chain together. This treatment converts the protein into a flexible polypeptide chain that has lost its

Figure 3-5 Three types of noncovalent bonds that help proteins fold. Although a single one of these bonds is quite weak, many of them often form together to create a strong bonding arrangement, as in the example shown. As in the previous figure, R is used as a general designation for an amino acid side chain.






Figure 3-6 How a protein folds into a compact conformation. The polar amino acid side chains tend to gather on the outside of the protein, where they can interact with water; the nonpolar amino acid side chains are buried on the inside to form a tightly packed hydrophobic core of atoms that are hidden from water. In this schematic drawing, the protein contains only about 30 amino acids.

natural shape. When the denaturing solvent is removed, the protein often refolds spontaneously, or *renatures*, into its original conformation (Figure 3-8), indicating that all the information needed for specifying the three-dimensional shape of a protein is contained in its amino acid sequence.

Each protein normally folds up into a single stable conformation. However, the conformation often changes slightly when the protein interacts with other molecules in the cell. This change in shape is often crucial to the function of the protein, as we see later.

Although a protein chain can fold into its correct conformation without outside help, protein folding in a living cell is often assisted by special proteins called *molecular chaperones*. These proteins bind to partly folded polypeptide chains and help them progress along the most energetically favorable folding pathway. Chaperones are vital in the crowded conditions of the cytoplasm, since they prevent the temporarily exposed hydrophobic regions in newly synthesized protein chains from associating with each other to form protein aggregates (see p. 357). However, the final three-dimensional shape of the protein is still specified by its amino acid sequence: chaperones simply make the folding process more reliable.

Proteins come in a wide variety of shapes, and they are generally between 50 and 2000 amino acids long. Large proteins generally consist of several distinct *protein domains*—structural units that fold more or less independently of each other, as we discuss below. The detailed structure of any protein is complicated; for simplicity a protein's structure can be depicted in several different ways, each emphasizing different features of the protein.




Figure 3-7 Hydrogen bonds in a protein molecule. Large numbers of hydrogen bonds form between adjacent regions of the folded polypeptide chain and help stabilize its three-dimensional shape. The protein depicted is a portion of the enzyme lysozyme, and the hydrogen bonds between the three possible pairs of partners have been differently colored, as indicated. (After C.K. Matthews and K.E. van Holde, *Biochemistry*, Redwood City, CA: Benjamin/Cummings, 1996.)




Figure 3–8 The refolding of a denatured protein. (A) This experiment demonstrates that the conformation of a protein is determined solely by its amino acid sequence. (B) The structure of urea. Urea is very soluble in water and unfolds proteins at high concentrations, where there is about one urea molecule for every six water molecules.

Panel 3–2 (pp. 138–139) presents four different depictions of a protein domain called SH2, which has important functions in eucaryotic cells. Constructed from a string of 100 amino acids, the structure is displayed as (A) a polypeptide backbone model, (B) a ribbon model, (C) a wire model that includes the amino acid side chains, and (D) a space-filling model. Each of the three horizontal rows shows the protein in a different orientation, and the image is colored in a way that allows the polypeptide chain to be followed from its N-terminus (*purple*) to its C-terminus (*red*).

Panel 3–2 shows that a protein's conformation is amazingly complex, even for a structure as small as the SH2 domain. But the description of protein structures can be simplified by the recognition that they are built up from several common structural motifs, as we discuss next.

The α Helix and the β Sheet Are Common Folding Patterns

When the three-dimensional structures of many different protein molecules are compared, it becomes clear that, although the overall conformation of each protein is unique, two regular folding patterns are often found in parts of them. Both patterns were discovered about 50 years ago from studies of hair and silk. The first folding pattern to be discovered, called the α helix, was found in the protein α -keratin, which is abundant in skin and its derivatives—such as hair, nails, and horns. Within a year of the discovery of the α helix, a second folded structure, called a β sheet, was found in the protein fibroin, the major constituent of silk. These two patterns are particularly common because they result from hydrogen-bonding between the N–H and C=O groups in the polypeptide backbone, without involving the side chains of the amino acids. Thus, they can be formed by many different amino acid sequences. In each case, the protein chain adopts a regular, repeating conformation. These two conformations, as well as the abbreviations that are used to denote them in ribbon models of proteins, are shown in Figure 3–9.

The core of many proteins contains extensive regions of β sheet. As shown in Figure 3–10, these β sheets can form either from neighboring polypeptide chains that run in the same orientation (parallel chains) or from a polypeptide chain that folds back and forth upon itself, with each section of the chain running in the direction opposite to that of its immediate neighbors (antiparallel chains). Both types of β sheet produce a very rigid structure, held together by hydrogen bonds that connect the peptide bonds in neighboring chains (see Figure 3–9D).

An α helix is generated when a single polypeptide chain twists around on itself to form a rigid cylinder. A hydrogen bond is made between every fourth peptide bond, linking the C=O of one peptide bond to the N–H of another (see Figure 3–9A). This gives rise to a regular helix with a complete turn every 3.6 amino acids. Note that the protein domain illustrated in Panel 3–2 contains two α helices, as well as β sheet structures.

Short regions of α helix are especially abundant in proteins located in cell membranes, such as transport proteins and receptors. As we discuss in Chapter 10, those portions of a transmembrane protein that cross the lipid bilayer usually cross as an α helix composed largely of amino acids with nonpolar side chains. The polypeptide backbone, which is hydrophilic, is hydrogen-bonded to itself in the α helix and shielded from the hydrophobic lipid environment of the membrane by its protruding nonpolar side chains (see also Figure 3–77).










Figure 3-9 The regular conformation of the polypeptide backbone observed in the α helix and the β sheet. (A, B, and C) The α helix. The N-H of every peptide bond is hydrogen-bonded to the C=O of a neighboring peptide bond located four peptide bonds away in the same chain. (D, E, and F) The β sheet. In this example, adjacent peptide chains run in opposite (antiparallel) directions. The individual polypeptide chains (strands) in a β sheet are held together by hydrogen-bonding between peptide bonds in different strands, and the amino acid side chains in each strand alternately project above and below the plane of the sheet. (A) and (D) show all the atoms in the polypeptide backbone, but the amino acid side chains are truncated and denoted by R. In contrast, (B) and (E) show the backbone atoms only, while (C) and (F) display the shorthand symbols that are used to represent the α helix and the β sheet in ribbon drawings of proteins (see Panel 3-2B).






PANEL 3-2 Four Different Ways of Depicting a Small Protein Domain: the SH2 Domain.
(Courtesy of David Lawson.)




(A) Backbone



(B) Ribbon



(C) Wire



(D) Space-filling

In other proteins, α helices wrap around each other to form a particularly stable structure, known as a **coiled-coil**. This structure can form when the two (or in some cases three) α helices have most of their nonpolar (hydrophobic) side chains on one side, so that they can twist around each other with these side chains facing inward (Figure 3-11). Long rodlike coiled-coils provide the structural framework for many elongated proteins. Examples are α -keratin, which forms the intracellular fibers that reinforce the outer layer of the skin and its appendages, and the myosin molecules responsible for muscle contraction.

The Protein Domain Is a Fundamental Unit of Organization

Even a small protein molecule is built from thousands of atoms linked together by precisely oriented covalent and noncovalent bonds, and it is extremely difficult to visualize such a complicated structure without a three-dimensional display. For this reason, various graphic and computer-based aids are used. A CD-ROM produced to accompany this book contains computer-generated images of selected proteins, designed to be displayed and rotated on the screen in a variety of formats.

Biologists distinguish four levels of organization in the structure of a protein. The amino acid sequence is known as the **primary structure** of the protein. Stretches of polypeptide chain that form α helices and β sheets constitute the protein's **secondary structure**. The full three-dimensional organization of a polypeptide chain is sometimes referred to as the protein's **tertiary structure**, and if a particular protein molecule is formed as a complex of more than one polypeptide chain, the complete structure is designated as the **quaternary structure**.

Studies of the conformation, function, and evolution of proteins have also revealed the central importance of a unit of organization distinct from the four just described. This is the **protein domain**, a substructure produced by any part of a polypeptide chain that can fold independently into a compact, stable structure. A domain usually contains between 40 and 350 amino acids, and it is the




Figure 3-10 Two types of β sheet structures. (A) An antiparallel β sheet (see Figure 3-9D). (B) A parallel β sheet. Both of these structures are common in proteins.




Figure 3-11 The structure of a coiled-coil. (A) A single α helix, with successive amino acid side chains labeled in a sevenfold sequence, "abcdefg" (from bottom to top). Amino acids "a" and "d" in such a sequence lie close together on the cylinder surface, forming a "stripe" (red) that winds slowly around the α helix. Proteins that form coiled-coils typically have nonpolar amino acids at positions "a" and "d." Consequently, as shown in (B), the two α helices can wrap around each other with the nonpolar side chains of one α helix interacting with the nonpolar side chains of the other, while the more hydrophilic amino acid side chains are left exposed to the aqueous environment. (C) The atomic structure of a coiled-coil determined by x-ray crystallography. The red side chains are nonpolar.




Figure 3-12 A protein formed from four domains. In the Src protein shown, two of the domains form a protein kinase enzyme, while the SH2 and SH3 domains perform regulatory functions. (A) A ribbon model, with ATP substrate in red. (B) A spacing-filling model, with ATP substrate in red. Note that the site that binds ATP is positioned at the interface of the two domains that form the kinase. The detailed structure of the SH2 domain is illustrated in Panel 3-2 (pp. 138–139).

modular unit from which many larger proteins are constructed. The different domains of a protein are often associated with different functions. Figure 3-12 shows an example—the Src protein kinase, which functions in signaling pathways inside vertebrate cells (Src is pronounced “sarc”). This protein has four domains: the SH2 and SH3 domains have regulatory roles, while the two remaining domains are responsible for the kinase catalytic activity. Later in the chapter, we shall return to this protein, in order to explain how proteins can form molecular switches that transmit information throughout cells.

The smallest protein molecules contain only a single domain, whereas larger proteins can contain as many as several dozen domains, usually connected to each other by short, relatively unstructured lengths of polypeptide chain. Figure 3-13 presents ribbon models of three differently organized protein domains. As these examples illustrate, the central core of a domain can be constructed from α helices, from β sheets, or from various combinations of these two fundamental folding elements. Each different combination is known as a *protein fold*. So far, about 1000 different protein folds have been identified among the ten thousand proteins whose detailed conformations are known.

Few of the Many Possible Polypeptide Chains Will Be Useful

Since each of the 20 amino acids is chemically distinct and each can, in principle, occur at any position in a protein chain, there are $20 \times 20 \times 20 \times 20 = 160,000$ different possible polypeptide chains four amino acids long, or 20^n different possible polypeptide chains n amino acids long. For a typical protein length of about 300 amino acids, more than 10^{390} (20^{300}) different polypeptide chains could theoretically be made. This is such an enormous number that to produce just one molecule of each kind would require many more atoms than exist in the universe.

Only a very small fraction of this vast set of conceivable polypeptide chains would adopt a single, stable three-dimensional conformation—by some estimates, less than one in a billion. The vast majority of possible protein molecules could adopt many conformations of roughly equal stability, each conformation having different chemical properties. And yet virtually all proteins present in cells adopt unique and stable conformations. How is this possible? The answer lies in natural selection. A protein with an unpredictably variable structure and biochemical activity is unlikely to help the survival of a cell that contains it. Such proteins would therefore have been eliminated by natural selection through the enormously long trial-and-error process that underlies biological evolution.

Because of natural selection, not only is the amino acid sequence of a present-day protein such that a single conformation is extremely stable, but this conformation has its chemical properties finely tuned to enable the protein to




Figure 3-13 Ribbon models of three different protein domains.

(A) Cytochrome *b*₅₆₂, a single-domain protein involved in electron transport in mitochondria. This protein is composed almost entirely of α helices. (B) The NAD-binding domain of the enzyme lactic dehydrogenase, which is composed of a mixture of α helices and β sheets. (C) The variable domain of an immunoglobulin (antibody) light chain, composed of a sandwich of two β sheets. In these examples, the α helices are shown in green, while strands organized as β sheets are denoted by red arrows.

Note that the polypeptide chain generally traverses back and forth across the entire domain, making sharp turns only at the protein surface. It is the protruding loop regions (yellow) that often form the binding sites for other molecules. (Adapted from drawings courtesy of Jane Richardson.)

perform a particular catalytic or structural function in the cell. Proteins are so precisely built that the change of even a few atoms in one amino acid can sometimes disrupt the structure of the whole molecule so severely that all function is lost.

Proteins Can Be Classified into Many Families

Once a protein had evolved that folded up into a stable conformation with useful properties, its structure could be modified during evolution to enable it to perform new functions. This process has been greatly accelerated by genetic mechanisms that occasionally produce duplicate copies of genes, allowing one gene copy to evolve independently to perform a new function (discussed in Chapter 7). This type of event has occurred quite often in the past; as a result, many present-day proteins can be grouped into protein families, each family member having an amino acid sequence and a three-dimensional conformation that resemble those of the other family members.

Consider, for example, the *serine proteases*, a large family of protein-cleaving (proteolytic) enzymes that includes the digestive enzymes chymotrypsin, trypsin, and elastase, and several proteases involved in blood clotting. When the protease portions of any two of these enzymes are compared, parts of their amino acid sequences are found to match. The similarity of their three-dimensional conformations is even more striking: most of the detailed twists and turns in their polypeptide chains, which are several hundred amino acids long, are virtually identical (Figure 3-14). The many different serine proteases nevertheless have distinct enzymatic activities, each cleaving different proteins or the peptide bonds between different types of amino acids. Each therefore performs a distinct function in an organism.

The story we have told for the serine proteases could be repeated for hundreds of other protein families. In many cases the amino acid sequences have diverged much further than for the serine proteases, so that one cannot be sure of a family relationship between two proteins without determining their three-dimensional structures. The yeast α 2 protein and the *Drosophila* engrailed protein, for example, are both gene regulatory proteins in the homeodomain family. Because they are identical in only 17 of their 60 amino acid residues, their relationship became certain only when their three-dimensional structures were compared (Figure 3-15).

The various members of a large protein family often have distinct functions. Some of the amino acid changes that make family members different were no doubt selected in the course of evolution because they resulted in useful changes in biological activity, giving the individual family members the different functional properties they have today. But many other amino acid changes are effectively “neutral,” having neither a beneficial nor a damaging effect on the




Figure 3-16 Percentage of total genes containing one or more copies of the indicated protein domain, as derived from complete genome sequences. Note that one of the three domains selected, the immunoglobulin domain, has been a relatively late addition, and its relative abundance has increased in the vertebrate lineage. The estimates of human gene numbers are approximate.

thousands of protein sequences from sequencing the genes that encode them, what technical developments can we look forward to next?

It is no longer a big step to progress from a gene sequence to the production of large amounts of the pure protein encoded by that gene. Thanks to DNA cloning and genetic engineering techniques (discussed in Chapter 8), this step is often routine. But there is still nothing routine about determining the complete three-dimensional structure of a protein. The standard technique based on x-ray diffraction requires that the protein be subjected to conditions that cause the molecules to aggregate into a large, perfectly ordered crystalline array—that is, a protein crystal. Each protein behaves quite differently in this respect, and protein crystals can be generated only through exhaustive trial-and-error methods that often take many years to succeed—if they succeed at all.

Membrane proteins and large protein complexes with many moving parts have generally been the most difficult to crystallize, which is why only a few such protein structures are displayed in this book. Increasingly, therefore, large proteins have been analyzed through determination of the structures of their individual domains: either by crystallizing isolated domains and then bombarding the crystals with x-rays, or by studying the conformations of isolated domains in concentrated aqueous solutions with powerful nuclear magnetic resonance (NMR) techniques (discussed in Chapter 8). From a combination of x-ray and NMR studies, we now know the three-dimensional shapes, or conformations, of thousands of different proteins.

By carefully comparing the conformations of known proteins, structural biologists (that is, experts on the structure of biological molecules) have concluded that there are a limited number of ways in which protein domains fold up—maybe as few as 2000. As we saw, the structures for about 1000 of these protein folds have thus far been determined; we may, therefore, already know half of the total number of possible structures for a protein domain. A complete catalog of all of the protein folds that exist in living organisms would therefore seem to be within our reach.

Sequence Homology Searches Can Identify Close Relatives

The present database of known protein sequences contains more than 500,000 entries, and it is growing very rapidly as more and more genomes are sequenced—revealing huge numbers of new genes that encode proteins. Powerful computer search programs are available that allow one to compare each newly discovered protein with this entire database, looking for possible relatives. Homologous proteins are defined as those whose genes have evolved from a common ancestral gene, and these are identified by the discovery of statistically significant similarities in amino acid sequences.

With such a large number of proteins in the database, the search programs find many nonsignificant matches, resulting in a background noise level that makes it very difficult to pick out all but the closest relatives. Generally speaking, a 30% identity in the sequence of two proteins is needed to be certain that a match has been found. However, many short signature sequences ("fingerprints") indicative of particular protein functions are known, and these are widely used to find more distant homologies (Figure 3-17).

These protein comparisons are important because related structures often imply related functions. Many years of experimentation can be saved by discovering that a new protein has an amino acid sequence homology with a protein of known function. Such sequence homologies, for example, first indicated that certain genes that cause mammalian cells to become cancerous are protein kinases. In the same way, many of the proteins that control pattern formation during the embryonic development of the fruit fly *Drosophila* were quickly recognized to be gene regulatory proteins.

Computational Methods Allow Amino Acid Sequences to Be Threaded into Known Protein Folds

We know that there are an enormous number of ways to make proteins with the same three-dimensional structure, and that—over evolutionary time—random mutations can cause amino acid sequences to change without a major change in the conformation of a protein. For this reason, one current goal of structural biologists is to determine all the different protein folds that proteins have in nature, and to devise computer-based methods to test the amino acid sequence of a domain to identify which one of these previously determined conformations the domain is likely to adopt.


A computational technique called threading can be used to fit an amino acid sequence to a particular protein fold. For each possible fold known, the computer searches for the best fit of the particular amino acid sequence to that structure. Are the hydrophobic residues on the inside? Are the sequences with a strong propensity to form an α helix in an α helix? And so on. The best fit gets a numerical score reflecting the estimated stability of the structure.

In many cases, one particular three-dimensional structure will stand out as a good fit for the amino acid sequence, suggesting an approximate conformation for the protein domain. In other cases, none of the known folds will seem possible. By applying x-ray and NMR studies to the latter class of proteins, structural biologists hope to be able to expand the number of known folds rapidly, aiming for a database that contains the complete library of protein folds that exist in nature. With such a library, plus expected improvements in the computational methods used for threading, it may eventually become possible to obtain an approximate three-dimensional structure for a protein as soon as its amino acid sequence is known.

Some Protein Domains, Called Modules, Form Parts of Many Different Proteins

As previously stated, most proteins are composed of a series of protein domains, in which different regions of the polypeptide chain have folded independently to form compact structures. Such multidomain proteins are believed to have originated when the DNA sequences that encode each domain accidentally became joined, creating a new gene. Novel binding surfaces have often been created at the juxtaposition of domains, and many of the functional sites where

Figure 3-17 The use of short signature sequences to find homologous protein domains. The two short sequences of 15 and 9 amino acids shown (green) can be used to search large databases for a protein domain that is found in many proteins, the SH2 domain. Here, the first 50 amino acids of the SH2 domain of 100 amino acids is compared for the human and *Drosophila* Src protein (see Figure 3-12). In the computer-generated sequence comparison (yellow row), exact matches between the human and *Drosophila* proteins are noted by the one-letter abbreviation for the amino acid; the positions with a similar but nonidentical amino acid are denoted by +, and nonmatches are blank. In this diagram, wherever one or both proteins contain an exact match to a position in the green sequences, both aligned sequences are colored red.



proteins bind to small molecules are found to be located there (for an example see Figure 3-12). Many large proteins show clear signs of having evolved by the joining of preexisting domains in new combinations, an evolutionary process called *domain shuffling* (Figure 3-18).

A subset of protein domains have been especially mobile during evolution; these so-called **protein modules** are generally somewhat smaller (40–200 amino acids) than an average domain, and they seem to have particularly versatile structures. The structure of one such module, the SH2 domain, was illustrated in Panel 3-2 (pp. 138–139). The structures of some additional protein modules are illustrated in Figure 3-19.

Each of the modules shown has a stable core structure formed from strands of β sheet, from which less-ordered loops of polypeptide chain protrude (*green*). The loops are ideally situated to form binding sites for other molecules, as most flagrantly demonstrated for the immunoglobulin fold, which forms the basis for antibody molecules (see Figure 3-42). The evolutionary success of such β -sheet-based modules is likely to have been due to their providing a convenient framework for the generation of new binding sites for ligands through small changes to these protruding loops.

A second feature of protein modules that explains their utility is the ease with which they can be integrated into other proteins. Five of the six modules illustrated in Figure 3-19 have their N- and C-terminal ends at opposite poles of the module. This “in-line” arrangement means that when the DNA encoding such a module undergoes tandem duplication, which is not unusual in the evolution of genomes (discussed in Chapter 7), the duplicated modules can be readily linked in series to form extended structures—either with themselves or with other in-line modules (Figure 3-20). Stiff extended structures composed of a series of modules are especially common in extracellular matrix molecules and in the extracellular portions of cell-surface receptor proteins. Other modules, including the SH2 domain and the kringle module illustrated in Figure 3-19, are




Figure 3-18 Domain shuffling. An extensive shuffling of blocks of protein sequence (protein domains) has occurred during protein evolution. Those portions of a protein denoted by the same shape and color in this diagram are evolutionarily related. Serine proteases like chymotrypsin are formed from two domains (*brown*). In the three other proteases shown, which are highly regulated and more specialized, these two protease domains are connected to one or more domains homologous to domains found in epidermal growth factor (EGF; *green*), to a calcium-binding protein (*yellow*), or to a “kringle” domain (*blue*) that contains three internal disulfide bridges. Chymotrypsin is illustrated in Figure 3-14.




Figure 3-19 The three-dimensional structures of some protein modules. In these ribbon diagrams, β -sheet strands are shown as *arrows*, and the N- and C-termini are indicated by *red spheres*. (Adapted from M. Baron, D.G. Norman, and I.D. Campbell, *Trends Biochem. Sci.* 16:13–17, 1991, and D.J. Leahy et al., *Science* 258:987–991, 1992.)

of a "plug-in" type. After genomic rearrangements, such modules are usually accommodated as an insertion into a loop region of a second protein.

The Human Genome Encodes a Complex Set of Proteins, Revealing Much That Remains Unknown

The result of sequencing the human genome has been surprising, because it reveals that our chromosomes contain only 30,000 to 35,000 genes. With regard to gene number, we would appear to be no more than 1.4-fold more complex than the tiny mustard weed, *Arabidopsis*, and less than 2-fold more complex than a nematode worm. The genome sequences also reveal that vertebrates have inherited nearly all of their protein domains from invertebrates—with only 7 percent of identified human domains being vertebrate-specific.

Each of our proteins is on average more complicated, however. A process of domain shuffling during vertebrate evolution has given rise to many novel combinations of protein domains, with the result that there are nearly twice as many combinations of domains found in human proteins as in a worm or a fly. Thus, for example, the trypsinlike serine protease domain is linked to at least 18 other types of protein domains in human proteins, whereas it is found covalently joined to only 5 different domains in the worm. This extra variety in our proteins greatly increases the range of protein-protein interactions possible (see Figure 3-78), but how it contributes to making us human is not known.

The complexity of living organisms is staggering, and it is quite sobering to note that we currently lack even the tiniest hint of what the function might be for more than 10,000 of the proteins that have thus far been identified in the human genome. There are certainly enormous challenges ahead for the next generation of cell biologists, with no shortage of fascinating mysteries to solve.

Larger Protein Molecules Often Contain More Than One Polypeptide Chain

The same weak noncovalent bonds that enable a protein chain to fold into a specific conformation also allow proteins to bind to each other to produce larger structures in the cell. Any region of a protein's surface that can interact with another molecule through sets of noncovalent bonds is called a **binding site**. A protein can contain binding sites for a variety of molecules, both large and small. If a binding site recognizes the surface of a second protein, the tight binding of two folded polypeptide chains at this site creates a larger protein molecule with a precisely defined geometry. Each polypeptide chain in such a protein is called a **protein subunit**.

In the simplest case, two identical folded polypeptide chains bind to each other in a "head-to-head" arrangement, forming a symmetric complex of two protein subunits (a *dimer*) held together by interactions between two identical binding sites. The *Cro repressor protein*—a gene regulatory protein that binds to DNA to turn genes off in a bacterial cell—provides an example (Figure 3-21). Many other types of symmetric protein complexes, formed from multiple copies of a single polypeptide chain, are commonly found in cells. The enzyme




Figure 3-20 An extended structure formed from a series of in-line protein modules. Four fibronectin type 3 modules (see Figure 3-19) from the extracellular matrix molecule fibronectin are illustrated in (A) ribbon and (B) space-filling models. (Adapted from D.J. Leahy, I. Aukhil, and H.P. Erickson, *Cell* 84:155-164, 1996.)




Figure 3-21 Two identical protein subunits binding together to form a symmetric protein dimer. The *Cro* repressor protein from bacteriophage lambda binds to DNA to turn off viral genes. Its two identical subunits bind head-to-head, held together by a combination of hydrophobic forces (blue) and a set of hydrogen bonds (yellow region). (Adapted from D.H. Ohlendorf, D.E. Tronrud, and B.W. Matthews, *J. Mol. Biol.* 280:129-136, 1998.)




Figure 3-22 A protein molecule containing multiple copies of a single protein subunit. The enzyme neuraminidase exists as a ring of four identical polypeptide chains. The small diagram shows how the repeated use of the same binding interaction forms the structure.

neuraminidase, for example, consists of four identical protein subunits, each bound to the next in a “head-to-tail” arrangement that forms a closed ring (Figure 3-22).

Many of the proteins in cells contain two or more types of polypeptide chains. *Hemoglobin*, the protein that carries oxygen in red blood cells, is a particularly well-studied example (Figure 3-23). It contains two identical α -globin subunits and two identical β -globin subunits, symmetrically arranged. Such multisubunit proteins are very common in cells, and they can be very large. Figure 3-24 provides a sampling of proteins whose exact structures are known, allowing the sizes and shapes of a few larger proteins to be compared with the relatively small proteins that we have thus far presented as models.

Some Proteins Form Long Helical Filaments

Some protein molecules can assemble to form filaments that may span the entire length of a cell. Most simply, a long chain of identical protein molecules can be constructed if each molecule has a binding site complementary to another region of the surface of the same molecule (Figure 3-25). An actin filament, for example, is a long helical structure produced from many molecules of the protein *actin* (Figure 3-26). Actin is very abundant in eucaryotic cells, where it constitutes one of the major filament systems of the cytoskeleton (discussed in Chapter 16).

Why is a helix such a common structure in biology? As we have seen, biological structures are often formed by linking subunits that are very similar to each other—such as amino acids or protein molecules—into long, repetitive chains. If all the subunits are identical, the neighboring subunits in the chain can often fit together in only one way, adjusting their relative positions to minimize the free energy of the contact between them. As a result, each subunit is positioned in exactly the same way in relation to the next, so that subunit 3 fits onto subunit 2 in the same way that subunit 2 fits onto subunit 1, and so on. Because it is very rare for subunits to join up in a straight line, this arrangement generally results in a helix—a regular structure that resembles a spiral staircase,




Figure 3-23 A protein formed as a symmetric assembly of two different subunits. Hemoglobin is an abundant protein in red blood cells that contains two copies of α globin and two copies of β globin. Each of these four polypeptide chains contains a heme molecule (red), which is the site where oxygen (O_2) is bound. Thus, each molecule of hemoglobin in the blood carries four molecules of oxygen.




Figure 3-24 A collection of protein molecules, shown at the same scale. For comparison, a DNA molecule bound to a protein is also illustrated. These space-filling models represent a range of sizes and shapes. Hemoglobin, catalase, porin, alcohol dehydrogenase, and aspartate transcarbamoylase are formed from multiple copies of subunits. The SH2 domain (*top left*) is presented in detail in Panel 3-2 (pp. 138-139). (After David S. Goodsell, *Our Molecular Nature*. New York: Springer-Verlag, 1996.)




Figure 3-25 Protein assemblies. (A) A protein with just one binding site can form a dimer with another identical protein. (B) Identical proteins with two different binding sites often form a long helical filament. (C) If the two binding sites are disposed appropriately in relation to each other, the protein subunits may form a closed ring instead of a helix. (For an example of A, see Figure 3-21; for an example of C, see Figure 3-22.)

as illustrated in Figure 3-27. Depending on the twist of the staircase, a helix is said to be either right-handed or left-handed (Figure 3-27E). Handedness is not affected by turning the helix upside down, but it is reversed if the helix is reflected in the mirror.

Helices occur commonly in biological structures, whether the subunits are small molecules linked together by covalent bonds (for example, the amino acids in an α helix) or large protein molecules that are linked by noncovalent forces (for example, the actin molecules in actin filaments). This is not surprising. A helix is an unexceptional structure, and it is generated simply by placing many similar subunits next to each other, each in the same strictly repeated relationship to the one before.

A Protein Molecule Can Have an Elongated, Fibrous Shape

Most of the proteins we have discussed so far are *globular proteins*, in which the polypeptide chain folds up into a compact shape like a ball with an irregular surface. Enzymes tend to be globular proteins: even though many are large and complicated, with multiple subunits, most have an overall rounded shape (see Figure 3-24). In contrast, other proteins have roles in the cell requiring each individual protein molecule to span a large distance. These proteins generally have a relatively simple, elongated three-dimensional structure and are commonly referred to as *fibrous proteins*.

One large family of intracellular fibrous proteins consists of α -keratin, introduced earlier, and its relatives. Keratin filaments are extremely stable and are the main component in long-lived structures such as hair, horn, and nails. An α -keratin molecule is a dimer of two identical subunits, with the long α helices of each subunit forming a coiled-coil (see Figure 3-11). The coiled-coil regions are capped at each end by globular domains containing binding sites. This enables this class of protein to assemble into ropelike *intermediate filaments*—an important component of the cytoskeleton that creates the cell's internal structural scaffold (see Figure 16-16).




Figure 3-26 Actin filaments. (A) Transmission electron micrographs of negatively stained actin filaments. (B) The helical arrangement of actin molecules in an actin filament. (A, courtesy of Roger Craig.)






Figure 3-27 Some properties of a helix. (A–D) A helix forms when a series of subunits bind to each other in a regular way. At the bottom, the interaction between two subunits is shown; behind them are the helices that result. These helices have two (A), three (B), and six (C and D) subunits per helical turn. At the top, the arrangement of subunits has been photographed from directly above the helix. Note that the helix in (D) has a wider path than that in (C), but the same number of subunits per turn. (E) A helix can be either right-handed or left-handed. As a reference, it is useful to remember that standard metal screws, which insert when turned clockwise, are right-handed. Note that a helix retains the same handedness when it is turned upside down.

Fibrous proteins are especially abundant outside the cell, where they are a main component of the gel-like *extracellular matrix* that helps to bind collections of cells together to form tissues. Extracellular matrix proteins are secreted by the cells into their surroundings, where they often assemble into sheets or long fibrils. *Collagen* is the most abundant of these proteins in animal tissues. A collagen molecule consists of three long polypeptide chains, each containing the nonpolar amino acid glycine at every third position. This regular structure allows the chains to wind around one another to generate a long regular triple helix (Figure 3-28A). Many collagen molecules then bind to one another side-by-side and end-to-end to create long overlapping arrays—thereby generating the extremely tough collagen fibrils that give connective tissues their tensile strength, as described in Chapter 19.

In complete contrast to collagen is another protein in the extracellular matrix, *elastin*. Elastin molecules are formed from relatively loose and unstructured polypeptide chains that are covalently cross-linked into a rubberlike elastic meshwork: unlike most proteins, they do not have a uniquely defined stable structure, but can be reversibly pulled from one conformation to another, as illustrated in Figure 3-28B. The resulting elastic fibers enable skin and other tissues, such as arteries and lungs, to stretch and recoil without tearing.

Extracellular Proteins Are Often Stabilized by Covalent Cross-Linkages

Many protein molecules are either attached to the outside of a cell's plasma membrane or secreted as part of the extracellular matrix. All such proteins are directly exposed to extracellular conditions. To help maintain their structures, the polypeptide chains in such proteins are often stabilized by covalent cross-linkages. These linkages can either tie two amino acids in the same protein together, or connect different polypeptide chains in a multisubunit protein. The most common cross-linkages in proteins are covalent sulfur–sulfur bonds. These *disulfide bonds* (also called *S–S bonds*) form as proteins are being prepared for export from cells. As described in Chapter 12, their formation is catalyzed in the endoplasmic reticulum by an enzyme that links together two pairs of –SH groups of cysteine side chains that are adjacent in the folded protein (Figure 3-29). Disulfide bonds do not change the conformation of a protein but



instead act as atomic staples to reinforce its most favored conformation. For example, lysozyme—an enzyme in tears that dissolves bacterial cell walls—retains its antibacterial activity for a long time because it is stabilized by such cross-linkages.

Disulfide bonds generally fail to form in the cell cytosol, where a high concentration of reducing agents converts S-S bonds back to cysteine -SH groups. Apparently, proteins do not require this type of reinforcement in the relatively mild environment inside the cell.

Protein Molecules Often Serve as Subunits for the Assembly of Large Structures

The same principles that enable a protein molecule to associate with itself to form rings or filaments operate to generate much larger structures in the cell—supramolecular structures such as enzyme complexes, ribosomes, protein filaments, viruses, and membranes. These large objects are not made as single, giant, covalently linked molecules. Instead they are formed by the noncovalent assembly of many separately manufactured molecules, which serve as the subunits of the final structure.

The use of smaller subunits to build larger structures has several advantages:

1. A large structure built from one or a few repeating smaller subunits requires only a small amount of genetic information.

Figure 3-28 Collagen and elastin. (A) Collagen is a triple helix formed by three extended protein chains that wrap around one another (*bottom*). Many rodlike collagen molecules are cross-linked together in the extracellular space to form unextendable collagen fibrils (*top*) that have the tensile strength of steel. The striping on the collagen fibril is caused by the regular repeating arrangement of the collagen molecules within the fibril. (B) Elastin polypeptide chains are cross-linked together to form rubberlike, elastic fibers. Each elastin molecule uncoils into a more extended conformation when the fiber is stretched and recoils spontaneously as soon as the stretching force is relaxed.




Figure 3-29 Disulfide bonds. This diagram illustrates how covalent disulfide bonds form between adjacent cysteine side chains. As indicated, these cross-linkages can join either two parts of the same polypeptide chain or two different polypeptide chains. Since the energy required to break one covalent bond is much larger than the energy required to break even a whole set of noncovalent bonds (see Table 2-2, p. 57), a disulfide bond can have a major stabilizing effect on a protein.





Figure 3-30 An example of single protein subunit assembly requiring multiple protein-protein contacts. Hexagonally packed globular protein subunits can form either a flat sheet or a tube.

2. Both assembly and disassembly can be readily controlled, reversible processes, since the subunits associate through multiple bonds of relatively low energy.
3. Errors in the synthesis of the structure can be more easily avoided, since correction mechanisms can operate during the course of assembly to exclude malformed subunits.

Some protein subunits assemble into flat sheets in which the subunits are arranged in hexagonal patterns. Specialized membrane proteins are sometimes arranged this way in lipid bilayers. With a slight change in the geometry of the individual subunits, a hexagonal sheet can be converted into a tube (Figure 3-30) or, with more changes, into a hollow sphere. Protein tubes and spheres that bind specific RNA and DNA molecules form the coats of viruses.

The formation of closed structures, such as rings, tubes, or spheres, provides additional stability because it increases the number of bonds between the protein subunits. Moreover, because such a structure is created by mutually dependent, cooperative interactions between subunits, it can be driven to assemble or disassemble by a relatively small change that affects each subunit individually. These principles are dramatically illustrated in the protein coat or *capsid* of many simple viruses, which takes the form of a hollow sphere (Figure 3-31).

Figure 3-31 The capsids of some viruses, all shown at the same scale. (A) Tomato bushy stunt virus; (B) poliovirus; (C) simian virus 40 (SV40); (D) satellite tobacco necrosis virus. The structures of all of these capsids have been determined by x-ray crystallography and are known in atomic detail. (Courtesy of Robert Grant, Stephan Crainic, and James M. Hogle.)






Figure 3–32 The structure of a spherical virus. In many viruses, identical protein subunits pack together to create a spherical shell (a capsid) that encloses the viral genome, composed of either RNA or DNA (see also Figure 3–31). For geometric reasons, no more than 60 identical subunits can pack together in a precisely symmetric way. If slight irregularities are allowed, however, more subunits can be used to produce a larger capsid. The tomato bushy stunt virus (TBSV) shown here, for example, is a spherical virus about 33 nm in diameter formed from 180 identical copies of a 386 amino acid capsid protein plus an RNA genome of 4500 nucleotides. To construct such a large capsid, the protein must be able to fit into three somewhat different environments, each of which is differently colored in the virus particle shown here. The postulated pathway of assembly is shown; the precise three-dimensional structure has been determined by x-ray diffraction. (Courtesy of Steve Harrison.)

Capsids are often made of hundreds of identical protein subunits that enclose and protect the viral nucleic acid (Figure 3–32). The protein in such a capsid must have a particularly adaptable structure: it must not only make several different kinds of contacts to create the sphere, it must also change this arrangement to let the nucleic acid out to initiate viral replication once the virus has entered a cell.

Many Structures in Cells Are Capable of Self-Assembly

The information for forming many of the complex assemblies of macromolecules in cells must be contained in the subunits themselves, because purified subunits can spontaneously assemble into the final structure under the appropriate conditions. The first large macromolecular aggregate shown to be capable of self-assembly from its component parts was *tobacco mosaic virus (TMV)*. This virus is a long rod in which a cylinder of protein is arranged around a helical RNA core (Figure 3–33). If the dissociated RNA and protein subunits are mixed together in solution, they recombine to form fully active viral particles. The assembly process is unexpectedly complex and includes the formation of double rings of protein, which serve as intermediates that add to the growing viral coat.

Another complex macromolecular aggregate that can reassemble from its component parts is the bacterial ribosome. This structure is composed of about





Figure 3-33 The structure of tobacco mosaic virus (TMV). (A) An electron micrograph of the viral particle, which consists of a single long RNA molecule enclosed in a cylindrical protein coat composed of identical protein subunits. (B) A model showing part of the structure of TMV. A single-stranded RNA molecule of 6000 nucleotides is packaged in a helical coat constructed from 2130 copies of a coat protein 158 amino acids long. Fully infective viral particles can self-assemble in a test tube from purified RNA and protein molecules. (A, courtesy of Robley Williams; B, courtesy of Richard J. Feldmann.)

55 different protein molecules and 3 different rRNA molecules. If the individual components are incubated under appropriate conditions in a test tube, they spontaneously re-form the original structure. Most importantly, such reconstituted ribosomes are able to perform protein synthesis. As might be expected, the reassembly of ribosomes follows a specific pathway: after certain proteins have bound to the RNA, this complex is then recognized by other proteins, and so on, until the structure is complete.

It is still not clear how some of the more elaborate self-assembly processes are regulated. Many structures in the cell, for example, seem to have a precisely defined length that is many times greater than that of their component macromolecules. How such length determination is achieved is in many cases a mystery. Three possible mechanisms are illustrated in Figure 3-34. In the simplest case, a long core protein or other macromolecule provides a scaffold that determines the extent of the final assembly. This is the mechanism that determines the length of the TMV particle, where the RNA chain provides the core. Similarly, a core protein is thought to determine the length of the thin filaments in muscle, as well as the length of the long tails of some bacterial viruses (Figure 3-35).




Figure 3-34 Three mechanisms of length determination for large protein assemblies. (A) Coassembly along an elongated core protein or other macromolecule that acts as a measuring device. (B) Termination of assembly because of strain that accumulates in the polymeric structure as additional subunits are added, so that beyond a certain length the energy required to fit another subunit onto the chain becomes excessively large. (C) A vernier type of assembly, in which two sets of rodlike molecules differing in length form a staggered complex that grows until their ends exactly match. The name derives from a measuring device based on the same principle, used in mechanical instruments.




Figure 3-35 An electron micrograph of bacteriophage lambda. The tip of the virus tail attaches to a specific protein on the surface of a bacterial cell, after which the tightly packaged DNA in the head is injected through the tail into the cell. The tail has a precise length, determined by the mechanism shown in Figure 3-34A.

The Formation of Complex Biological Structures Is Often Aided by Assembly Factors

Not all cellular structures held together by noncovalent bonds are capable of self-assembly. A mitochondrion, a cilium, or a myofibril of a muscle cell, for example, cannot form spontaneously from a solution of its component macromolecules. In these cases, part of the assembly information is provided by special enzymes and other cellular proteins that perform the function of templates, guiding construction but taking no part in the final assembled structure.

Even relatively simple structures may lack some of the ingredients necessary for their own assembly. In the formation of certain bacterial viruses, for example, the head, which is composed of many copies of a single protein subunit, is assembled on a temporary scaffold composed of a second protein. Because the second protein is absent from the final viral particle, the head structure cannot spontaneously reassemble once it has been taken apart. Other examples are known in which proteolytic cleavage is an essential and irreversible step in the normal assembly process. This is even the case for some small protein assemblies, including the structural protein collagen and the hormone insulin (Figure 3-36). From these relatively simple examples, it seems very likely that the assembly of a structure as complex as a mitochondrion or a cilium will involve temporal and spatial ordering imparted by numerous other cell components.

Summary

The three-dimensional conformation of a protein molecule is determined by its amino acid sequence. The folded structure is stabilized by noncovalent interactions between different parts of the polypeptide chain. The amino acids with hydrophobic side chains tend to cluster in the interior of the molecule, and local hydrogen-bond interactions between neighboring peptide bonds give rise to α helices and β sheets.

Globular regions, known as domains, are the modular units from which many proteins are constructed; such domains generally contain 40–350 amino acids. Small proteins typically consist of only a single domain, while large proteins are formed from several domains linked together by short lengths of polypeptide chain. As proteins have evolved, domains have been modified and combined with other domains to construct new proteins. Domains that participate in the formation of large numbers of proteins are known as protein modules. Thus far, about 1000 different ways of folding up a domain have been observed, among more than about 10,000 known protein structures.

Proteins are brought together into larger structures by the same noncovalent forces that determine protein folding. Proteins with binding sites for their own surface can assemble into dimers, closed rings, spherical shells, or helical polymers. Although mixtures of proteins and nucleic acids can assemble spontaneously into complex structures in a test tube, many biological assembly processes involve irreversible steps. Consequently, not all structures in the cell are capable of spontaneous reassembly after they have been dissociated into their component parts.

PROTEIN FUNCTION

We have seen that each type of protein consists of a precise sequence of amino acids that allows it to fold up into a particular three-dimensional shape, or conformation. But proteins are not rigid lumps of material. They can have precisely engineered moving parts whose mechanical actions are coupled to chemical events. It is this coupling of chemistry and movement that gives proteins the extraordinary capabilities that underlie the dynamic processes in living cells.

In this section, we explain how proteins bind to other selected molecules and how their activity depends on such binding. We show that the ability to bind to other molecules enables proteins to act as catalysts, signal receptors, switches, motors, or tiny pumps. The examples we discuss in this chapter by no means exhaust the vast functional repertoire of proteins. However, the specialized




Figure 3-36 Proteolytic cleavage in insulin assembly. The polypeptide hormone insulin cannot spontaneously re-form efficiently if its disulfide bonds are disrupted. It is synthesized as a larger protein (proinsulin) that is cleaved by a proteolytic enzyme after the protein chain has folded into a specific shape. Excision of part of the proinsulin polypeptide chain removes some of the information needed for the protein to fold spontaneously into its normal conformation once it has been denatured and its two polypeptide chains separated.




Figure 3-37 The selective binding of a protein to another molecule. Many weak bonds are needed to enable a protein to bind tightly to a second molecule, which is called a *ligand* for the protein. A ligand must therefore fit precisely into a protein's binding site, like a hand into a glove, so that a large number of noncovalent bonds can be formed between the protein and the ligand.

functions of many of the proteins you will encounter elsewhere in this book are based on similar principles.

All Proteins Bind to Other Molecules

The biological properties of a protein molecule depend on its physical interaction with other molecules. Thus, antibodies attach to viruses or bacteria to mark them for destruction, the enzyme hexokinase binds glucose and ATP so as to catalyze a reaction between them, actin molecules bind to each other to assemble into actin filaments, and so on. Indeed, all proteins stick, or *bind*, to other molecules. In some cases, this binding is very tight; in others, it is weak and short-lived. But the binding always shows great *specificity*, in the sense that each protein molecule can usually bind just one or a few molecules out of the many thousands of different types it encounters. The substance that is bound by the protein—no matter whether it is an ion, a small molecule, or a macromolecule—is referred to as a **ligand** for that protein (from the Latin word *ligare*, meaning “to bind”).

The ability of a protein to bind selectively and with high affinity to a ligand depends on the formation of a set of weak, noncovalent bonds—hydrogen bonds, ionic bonds, and van der Waals attractions—plus favorable hydrophobic interactions (see Panel 2-3, pp. 114–115). Because each individual bond is weak, an effective binding interaction requires that many weak bonds be formed simultaneously. This is possible only if the surface contours of the ligand molecule fit very closely to the protein, matching it like a hand in a glove (Figure 3-37).

The region of a protein that associates with a ligand, known as the ligand's *binding site*, usually consists of a cavity in the protein surface formed by a particular arrangement of amino acids. These amino acids can belong to different portions of the polypeptide chain that are brought together when the protein folds (Figure 3-38). Separate regions of the protein surface generally provide binding sites for different ligands, allowing the protein's activity to be regulated, as we shall see later. And other parts of the protein can serve as a handle to place the protein in a particular location in the cell—an example is the SH2 domain discussed previously, which is often used to move a protein containing it to sites in the plasma membrane in response to particular signals.

Although the atoms buried in the interior of the protein have no direct contact with the ligand, they provide an essential scaffold that gives the surface its contours and chemical properties. Even small changes to the amino acids in the interior of a protein molecule can change its three-dimensional shape enough to destroy a binding site on the surface.

The Details of a Protein's Conformation Determine Its Chemistry

Proteins have impressive chemical capabilities because the neighboring chemical groups on their surface often interact in ways that enhance the chemical reactivity of amino acid side chains. These interactions fall into two main categories.





Figure 3-38 The binding site of a protein. (A) The folding of the polypeptide chain typically creates a crevice or cavity on the protein surface. This crevice contains a set of amino acid side chains disposed in such a way that they can make noncovalent bonds only with certain ligands. (B) A close-up of an actual binding site showing the hydrogen bonds and ionic interactions formed between a protein and its ligand (in this example, cyclic AMP is the bound ligand).

First, neighboring parts of the polypeptide chain may interact in a way that restricts the access of water molecules to a ligand binding site. Because water molecules tend to form hydrogen bonds, they can compete with ligands for sites on the protein surface. The tightness of hydrogen bonds (and ionic interactions) between proteins and their ligands is therefore greatly increased if water molecules are excluded. Initially, it is hard to imagine a mechanism that would exclude a molecule as small as water from a protein surface without affecting the access of the ligand itself. Because of the strong tendency of water molecules to form water-water hydrogen bonds, however, water molecules exist in a large hydrogen-bonded network (see Panel 2-2, pp. 112-113). In effect, a ligand binding site can be kept dry because it is energetically unfavorable for individual water molecules to break away from this network, as they must do to reach into a crevice on a protein's surface.

Second, the clustering of neighboring polar amino acid side chains can alter their reactivity. If a number of negatively charged side chains are forced together against their mutual repulsion by the way the protein folds, for example, the affinity of the site for a positively charged ion is greatly increased. In addition, when amino acid side chains interact with one another through hydrogen bonds, normally unreactive side groups (such as the $-CH_2OH$ on the serine shown in Figure 3-39) can become reactive, enabling them to enter into reactions that make or break selected covalent bonds.

The surface of each protein molecule therefore has a unique chemical reactivity that depends not only on which amino acid side chains are exposed, but also on their exact orientation relative to one another. For this reason, even two slightly different conformations of the same protein molecule may differ greatly in their chemistry.

Figure 3-39 An unusually reactive amino acid at the active site of an enzyme. This example is the "catalytic triad" found in chymotrypsin, elastase, and other serine proteases (see Figure 3-14). The aspartic acid side chain (Asp 102) induces the histidine (His 57) to remove the proton from serine 195. This activates the serine to form a covalent bond with the enzyme substrate, hydrolyzing a peptide bond.



Sequence Comparisons Between Protein Family Members Highlight Crucial Ligand Binding Sites

As we have described previously, many of the domains in proteins can be grouped into families that show clear evidence of their evolution from a common ancestor, and genome sequences reveal large numbers of proteins that contain one or more common domains. The three-dimensional structures of the members of the same domain family are remarkably similar. For example, even when the amino acid sequence identity falls to 25%, the backbone atoms in a domain have been found to follow a common protein fold within 0.2 nanometers (2 Å).

These facts allow a method called “evolutionary tracing” to be used to identify those sites in a protein domain that are the most crucial to the domain’s function. For this purpose, those amino acids that are unchanged, or nearly unchanged, in all of the known protein family members are mapped onto a structural model of the three-dimensional structure of one family member. When this is done, the most invariant positions often form one or more clusters on the protein surface, as illustrated in Figure 3–40A for the SH2 domain described previously (see Panel 3–2, pp. 138–139). These clusters generally correspond to ligand binding sites.

The SH2 domain is a module that functions in protein–protein interactions. It binds the protein containing it to a second protein that contains a phosphorylated tyrosine side chain in a specific amino acid sequence context, as shown in Figure 3–40B. The amino acids located at the binding site for the phosphorylated polypeptide have been the slowest to change during the long evolutionary process that produced the large SH2 family of peptide recognition domains. Because mutation is a random process, this result is attributed to the preferential elimination during evolution of all organisms whose SH2 domains became altered in a way that inactivated the SH2-binding site, thereby destroying the function of the SH2 domain.


In this era of extensive genome sequencing, many new protein families have been discovered whose functions are unknown. By identifying the critical binding sites on a three-dimensional structure determined for one family member, the above method of evolutionary tracing is being used to help determine the functions of such proteins.

Proteins Bind to Other Proteins Through Several Types of Interfaces

Proteins can bind to other proteins in at least three ways. In many cases, a portion of the surface of one protein contacts an extended loop of polypeptide chain (a “string”) on a second protein (Figure 3–41A). Such a surface–string interaction, for example, allows the SH2 domain to recognize a phosphorylated polypeptide as a loop on a second protein, as just described, and it also enables a protein kinase to recognize the proteins that it will phosphorylate (see below).

Figure 3–40 The evolutionary trace method applied to the SH2 domain.

(A) Front and back views of a space-filling model of the SH2 domain, with evolutionarily conserved amino acids on the protein surface colored yellow, and those more toward the protein interior colored red. (B) The structure of the SH2 domain with its bound polypeptide. Here, those amino acids located within 0.4 nm of the bound ligand are colored blue. The two key amino acids of the ligand are yellow, and the others are purple. (Adapted from O. Lichtarge, H.R. Bourne, and F.E. Cohen, *J. Mol. Biol.* 257:342–358, 1996.)






Figure 3-41 Three ways in which two proteins can bind to each other. Only the interacting parts of the two proteins are shown. (A) A rigid surface on one protein can bind to an extended loop of polypeptide chain (a “string”) on a second protein. (B) Two α helices can bind together to form a coiled-coil. (C) Two complementary rigid surfaces often link two proteins together.

A second type of protein–protein interface is formed when two α helices, one from each protein, pair together to form a coiled-coil (Figure 3-41B). This type of protein interface is found in several families of gene regulatory proteins, as discussed in Chapter 7.

The most common way for proteins to interact, however, is by the precise matching of one rigid surface with that of another (Figure 3-41C). Such interactions can be very tight, since a large number of weak bonds can form between two surfaces that match well. For the same reason, such surface–surface interactions can be extremely specific, enabling a protein to select just one partner from the many thousands of different proteins found in a cell.

The Binding Sites of Antibodies Are Especially Versatile

All proteins must bind to particular ligands to carry out their various functions. This capacity for tight selective binding is displayed to an extraordinary degree by the antibody family, as discussed in detail in Chapter 24.

Antibodies, or immunoglobulins, are proteins produced by the immune system in response to foreign molecules, such as those on the surface of an invading microorganism. Each antibody binds to a particular target molecule extremely tightly, thereby either inactivating the target directly or marking it for destruction. An antibody recognizes its target (called an **antigen**) with remarkable specificity. Because there are potentially billions of different antigens we might encounter, we have to be able to produce billions of different antibodies.

Antibodies are Y-shaped molecules with two identical binding sites that are complementary to a small portion of the surface of the antigen molecule. A detailed examination of the antigen-binding sites of antibodies reveals that they are formed from several loops of polypeptide chain that protrude from the ends of a pair of closely juxtaposed protein domains (Figure 3-42). The enormous diversity of antigen-binding sites possessed by different antibodies is generated by changing only the length and amino acid sequence of these loops, without altering the basic protein structure.

Loops of this kind are ideal for grasping other molecules. They allow a large number of chemical groups to surround a ligand so that the protein can link to it with many weak bonds. For this reason, loops are often used to form the ligand-binding sites in proteins.

Binding Strength Is Measured by the Equilibrium Constant

Molecules in the cell encounter each other very frequently because of their continual random thermal movements. When colliding molecules have poorly matching surfaces, few noncovalent bonds form, and the two molecules dissociate as rapidly as they come together. At the other extreme, when many noncovalent bonds form, the association can persist for a very long time (Figure 3-43). Strong interactions occur in cells whenever a biological function requires that molecules remain associated for a long time—for example, when a group of RNA and protein molecules come together to make a subcellular structure such as a ribosome.

The strength with which any two molecules bind to each other can be measured directly. As an example, imagine a situation in which a population of





Figure 3-42 An antibody molecule.

(A) A typical antibody molecule is Y-shaped and has two identical binding sites for its antigen, one on each arm of the Y. The protein is composed of four polypeptide chains (two identical heavy chains and two identical and smaller light chains) held together by disulfide bonds. Each chain is made up of several different immunoglobulin domains, here shaded either blue or gray. The antigen-binding site is formed where a heavy-chain variable domain (V_H) and a light-chain variable domain (V_L) come close together. These are the domains that differ most in their sequence and structure in different antibodies. (B) This ribbon model of a light chain shows the parts of the V_L domain most closely involved in binding to the antigen in red. They contribute half of the fingerlike loops that fold around each of the antigen molecules in (A).

identical antibody molecules suddenly encounters a population of ligands diffusing in the fluid surrounding them. At frequent intervals, one of the ligand molecules will bump into the binding site of an antibody and form an antibody–ligand complex. The population of antibody–ligand complexes will therefore increase, but not without limit: over time, a second process, in which individual complexes break apart because of thermally induced motion, will become increasingly important. Eventually, any population of antibody molecules and ligands will reach a steady state, or equilibrium, in which the number of binding (association) events per second is precisely equal to the number of “unbinding” (dissociation) events (see Figure 2-52).


From the concentrations of the ligand, antibody, and antibody–ligand complex at equilibrium, one can calculate a convenient measure—termed the **equilibrium constant (K)**—of the strength of binding (Figure 3-44A). The equilibrium constant is greater the greater the binding strength, and it is a direct measure of the free-energy difference between the bound and free states (Figure 3-44B). Even a change of a few noncovalent bonds can have a striking effect on a binding interaction, as shown by the example in Figure 3-44C.

Figure 3-43 How noncovalent bonds mediate interactions between macromolecules.



the surfaces of molecules A and B, and A and C, are a poor match and are capable of forming only a few weak bonds; thermal motion rapidly breaks them apart

the surfaces of molecules A and D match well and therefore can form enough weak bonds to withstand thermal jolting; they therefore stay bound to each other



We have used the case of an antibody binding to its ligand to illustrate the effect of binding strength on the equilibrium state, but the same principles apply to any molecule and its ligand. Many proteins are enzymes, which, as we now discuss, first bind to their ligands and then catalyze the breakage or formation of covalent bonds in these molecules.

Enzymes Are Powerful and Highly Specific Catalysts

Many proteins can perform their function simply by binding to another molecule. An actin molecule, for example, need only associate with other actin molecules to form a filament. There are other proteins, however, for which ligand binding is only a necessary first step in their function. This is the case for the large and very important class of proteins called **enzymes**. As described in Chapter 2, enzymes are remarkable molecules that determine all the chemical transformations that make and break covalent bonds in cells. They bind to one or more ligands, called **substrates**, and convert them into one or more chemically modified **products**, doing this over and over again with amazing rapidity. Enzymes speed up reactions, often by a factor of a million or more, without themselves being changed—that is, they act as **catalysts** that permit cells to make or break covalent bonds in a controlled way. It is the catalysis of organized sets of chemical reactions by enzymes that creates and maintains the cell, making life possible.

Enzymes can be grouped into functional classes that perform similar chemical reactions (Table 3-1). Each type of enzyme within such a class is highly specific, catalyzing only a single type of reaction. Thus, *hexokinase* adds a phosphate group to D-glucose but ignores its optical isomer L-glucose; the blood-clotting enzyme *thrombin* cuts one type of blood protein between a particular arginine and its adjacent glycine and nowhere else, and so on. As discussed in detail in Chapter 2, enzymes work in teams, with the product of one enzyme becoming the substrate for the next. The result is an elaborate network of metabolic pathways that provides the cell with energy and generates the many large and small molecules that the cell needs (see Figure 2-35).

Substrate Binding Is the First Step in Enzyme Catalysis

For a protein that catalyzes a chemical reaction (an enzyme), the binding of each substrate molecule to the protein is an essential prelude. In the simplest case, if we denote the enzyme by E, the substrate by S, and the product by P, the basic reaction path is $E + S \rightarrow ES \rightarrow EP \rightarrow E + P$. From this reaction path, we see that there is a limit to the amount of substrate that a single enzyme molecule can process in a given time. If the concentration of substrate is increased, the rate at

Figure 3-44 Relating binding energies to the equilibrium constant.

(A) The equilibrium between molecules A and B and the complex AB is maintained by a balance between the two opposing reactions shown in panels 1 and 2. Molecules A and B must collide if they are to react, and the association rate is therefore proportional to the product of their individual concentrations $[A] \times [B]$. (Square brackets indicate concentration.) As shown in panel 3, the ratio of the rate constants for the association and the dissociation reactions is equal to the equilibrium constant (K) for the reaction. (B) The equilibrium constant in panel 3 is that for the association reaction $A + B \leftrightarrow AB$, and the larger its value, the stronger the binding between A and B. Note that, for every 1.4 kcal/mole of free-energy drop, the equilibrium constant increases by a factor of 10. (C) An example of the dramatic effect that the presence or absence of a few weak bonds can have in a biological context.

The equilibrium constant here has units of liters/mole: for simple binding interactions it is also called the *affinity constant* or *association constant*, denoted K_a . The reciprocal of K_a is called the *dissociation constant*, K_d (in units of moles/liter).

TABLE 3-1 Some Common Types of Enzymes

ENZYME	REACTION CATALYZED
Hydrolases	general term for enzymes that catalyze a hydrolytic cleavage reaction.
Nucleases	break down nucleic acids by hydrolyzing bonds between nucleotides.
Proteases	break down proteins by hydrolyzing bonds between amino acids.
Synthases	general name used for enzymes that synthesize molecules in anabolic reactions by condensing two smaller molecules together.
Isomerases	catalyze the rearrangement of bonds within a single molecule.
Polymerases	catalyze polymerization reactions such as the synthesis of DNA and RNA.
Kinases	catalyze the addition of phosphate groups to molecules. Protein kinases are an important group of kinases that attach phosphate groups to proteins.
Phosphatases	catalyze the hydrolytic removal of a phosphate group from a molecule.
Oxido-Reductases	general name for enzymes that catalyze reactions in which one molecule is oxidized while the other is reduced. Enzymes of this type are often called <i>oxidases</i> , <i>reductases</i> , and <i>dehydrogenases</i> .
ATPases	hydrolyze ATP. Many proteins with a wide range of roles have an energy-harnessing ATPase activity as part of their function, for example, motor proteins such as <i>myosin</i> and membrane transport proteins such as the <i>sodium-potassium pump</i> .

Enzyme names typically end in "-ase," with the exception of some enzymes, such as pepsin, trypsin, thrombin and lysozyme that were discovered and named before the convention became generally accepted at the end of the nineteenth century. The common name of an enzyme usually indicates the substrate and the nature of the reaction catalyzed. For example, citrate synthase catalyzes the synthesis of citrate by a reaction between acetyl CoA and oxaloacetate.

which product is formed also increases, up to a maximum value (Figure 3-45). At that point the enzyme molecule is saturated with substrate, and the rate of reaction (V_{\max}) depends only on how rapidly the enzyme can process the substrate molecule. This maximum rate divided by the enzyme concentration is called the *turnover number*. The turnover number is often about 1000 substrate molecules processed per second per enzyme molecule, although turnover numbers between 1 and 10,000 are known.

The other kinetic parameter frequently used to characterize an enzyme is its K_m , the concentration of substrate that allows the reaction to proceed at one-half its maximum rate ($0.5 V_{\max}$) (see Figure 3-45). A *low* K_m value means that the enzyme reaches its maximum catalytic rate at a *low concentration* of substrate and generally indicates that the enzyme binds to its substrate very tightly, whereas a *high* K_m value corresponds to weak binding. The methods used to characterize enzymes in this way are explained in Panel 3-3 (pp. 164-165).

Enzymes Speed Reactions by Selectively Stabilizing Transition States

Extremely high rates of chemical reaction are achieved by enzymes—far higher than for any synthetic catalysts. This efficiency is attributable to several factors. The enzyme serves, first, to increase the local concentration of substrate molecules at the catalytic site and to hold all the appropriate atoms in the correct orientation for the reaction that is to follow. More importantly, however, some of the binding energy contributes directly to the catalysis. Substrate molecules must pass through a series of intermediate states of altered geometry




Figure 3-45 Enzyme kinetics. The rate of an enzyme reaction (V) increases as the substrate concentration increases until a maximum value (V_{\max}) is reached. At this point all substrate-binding sites on the enzyme molecules are fully occupied, and the rate of reaction is limited by the rate of the catalytic process on the enzyme surface. For most enzymes, the concentration of substrate at which the reaction rate is half-maximal (K_m) is a measure of how tightly the substrate is bound, with a large value of K_m corresponding to weak binding.

PANEL 3-3 Some of the Methods Used to Study Enzymes

WHY ANALYZE THE KINETICS OF ENZYMES?

Enzymes are the most selective and powerful catalysts known. An understanding of their detailed mechanisms provides a critical tool for the discovery of new drugs, for the large-scale industrial synthesis of useful chemicals, and for appreciating the chemistry of cells and organisms. A detailed study of the rates of the chemical reactions that are catalyzed by a purified enzyme—more specifically how these rates change with changes in conditions such as the concentrations of substrates, products, inhibitors, and

regulatory ligands—allows biochemists to figure out exactly how each enzyme works. For example, this is the way that the ATP-producing reactions of glycolysis, shown previously in Figure 2-73, were deciphered—allowing us to appreciate the rationale for this critical enzymatic pathway.

In this Panel, we introduce the important field of **enzyme kinetics**, which has been indispensable for deriving much of the detailed knowledge that we now have about cellular chemistry.

STEADY STATE ENZYME KINETICS

Many enzymes have only one substrate, which they bind and then process to produce products according to the scheme outlined in Figure 3-50A. In this case, the reaction is written as




Here we have assumed that the reverse reaction, in which E + P recombine to form EP and then ES, occurs so rarely that we can ignore it. In this case, we can express the rate of the reaction—known as its velocity, *V*, as

$$V = k_{cat} [ES]$$

where [ES] is the concentration of the enzyme substrate complex, and *k_{cat}* is the **turnover number**: a rate constant that is equal to the number of substrate molecules processed per enzyme molecule each second.

But how does the value of [ES] relate to the concentrations that we know directly, which are the total concentration of the enzyme, [E₀], and the concentration of the substrate, [S]? When enzyme and substrate are first mixed, the concentration [ES] will rise rapidly from zero to a so-called steady state level, as illustrated below



At this steady state, [ES] is nearly constant, so that

$$\text{rate of ES breakdown} = \text{rate of ES formation}$$

$$k_{-1} [ES] + k_{cat} [ES] = k_1 [E][S]$$

or, since the concentration of the free enzyme, [E], is equal to [E₀] - [ES]

$$[ES] = \left(\frac{k_1}{k_{-1} + k_{cat}} \right) [E][S] = \left(\frac{k_1}{k_{-1} + k_{cat}} \right) ([E_0] - [ES])[S]$$

Rearranging, and defining the constant *K_m* as

$$\frac{k_{-1} + k_{cat}}{k_1}$$

we get

$$[ES] = \frac{[E_0][S]}{K_m + [S]}$$

or, remembering that *V* = *k_{cat}* [ES], we obtain the famous Michaelis-Menton equation


$$V = \frac{k_{cat} [E_0][S]}{K_m + [S]}$$

As [S] is increased to higher and higher levels, essentially all of the enzyme will be bound to substrate at steady state; at this point, a maximum rate of reaction, *V_{max}*, will be reached where *V* = *V_{max}* = *k_{cat}* [E₀]. Thus, it is convenient to rewrite the Michaelis-Menton equation as

$$V = \frac{V_{max} [S]}{K_m + [S]}$$


THE DOUBLE RECIPROCAL PLOT

A typical plot of V versus $[S]$ for an enzyme that follows Michaelis-Menten kinetics is shown below. From this plot, neither the value of V_{max} nor of K_m is immediately clear.



To obtain V_{max} and K_m from such data, a double-reciprocal plot is often used, in which the Michaelis-Menten equation has merely been rearranged, so that $1/V$ can be plotted versus $1/[S]$.

$$1/V = \left(\frac{K_m}{V_{max}} \right) \left(\frac{1}{[S]} \right) + 1/V_{max}$$



THE SIGNIFICANCE OF K_m , k_{cat} , and k_{cat}/K_m

As described in the text, K_m is an approximate measure of substrate affinity for the enzyme: it is numerically equal to the concentration of $[S]$ at $V = 0.5 V_{max}$. In general, a lower value of K_m means tighter substrate binding.

We have seen that k_{cat} is the turnover number for the enzyme. At very low substrate concentrations, where $[S] \ll K_m$, most of the enzyme is free. Thus we can think of $[E] = [E_0]$, so that the Michaelis-Menten equation becomes $V = k_{cat}/K_m [E][S]$. Thus, the ratio k_{cat}/K_m is equivalent to the rate constant for the reaction between free enzyme and free substrate.

A comparison of k_{cat}/K_m for the same enzyme with different substrates, or for two enzymes with their different substrates, is widely used as a measure of enzyme effectiveness.

For simplicity, in this Panel we have discussed enzymes that have only one substrate, such as the lysozyme enzyme described in the text (see p. 167). Most enzymes have two substrates, one of which is often an active carrier molecule—such as NADH or ATP.

A similar, but more complex analysis is used to determine the kinetics of such enzymes—allowing the order of substrate binding and the presence of covalent intermediates along the pathway to be revealed (see, for example, Figure 2-73).

SOME ENZYMES ARE DIFFUSION LIMITED

The values of k_{cat} , K_m , and k_{cat}/K_m for some selected enzymes are given below:

enzyme	substrate	k_{cat} (sec ⁻¹)	K_m (M)	k_{cat}/K_m (sec ⁻¹ M ⁻¹)
acetylcholinesterase	acetylcholine	1.4×10^4	9×10^{-5}	1.6×10^8
catalase	H ₂ O ₂	4×10^7	1	4×10^7
fumarase	fumarate	8×10^2	5×10^{-6}	1.6×10^8

Because an enzyme and its substrate must collide before they can react, k_{cat}/K_m has a maximum possible value that is limited by collision rates. If every collision forms an enzyme-substrate complex, one can calculate from diffusion theory that k_{cat}/K_m will be between 10^8 and 10^9 sec⁻¹M⁻¹, in the case where all subsequent steps proceed immediately. Thus, it is claimed that enzymes like acetylcholinesterase and fumarase are "perfect enzymes", each enzyme having evolved to the point where nearly every collision with its substrate converts the substrate to a product.

and electron distribution before they form the ultimate products of the reaction. The free energy required to attain the most unstable **transition state** is called the *activation energy* for the reaction, and it is the major determinant of the reaction rate. Enzymes have a much higher affinity for the transition state of the substrate than they have for the stable form. Because this tight binding greatly lowers the energies of the transition state, the enzyme greatly accelerates a particular reaction by lowering the activation energy that is required (Figure 3–46).

A dramatic proof that stabilizing a transition state can greatly increase a reaction rate is provided by the intentional production of antibodies that act like enzymes. Consider, for example, the hydrolysis of an amide bond, which is similar to the peptide bond that joins two adjacent amino acids in a protein. In an aqueous solution, an amide bond hydrolyzes very slowly by the mechanism shown in Figure 3–47A. In the central intermediate, or transition state, the carbonyl carbon is bonded to four atoms arranged at the corners of a tetrahedron. By generating monoclonal antibodies that bind tightly to a stable analog of this very unstable *tetrahedral intermediate*, an antibody that functions like an enzyme can be obtained (Figure 3–47B). Because this *catalytic antibody* binds to and stabilizes the tetrahedral intermediate, it increases the spontaneous rate of amide-bond hydrolysis by more than 10,000-fold.

Enzymes Can Use Simultaneous Acid and Base Catalysis

Figure 3–48 compares the spontaneous reaction rates and the corresponding enzyme-catalyzed rates for five enzymes. Rate accelerations of 10^9 – 10^{23} are observed. Clearly, enzymes are much better catalysts than catalytic antibodies. Enzymes not only bind tightly to a transition state, they also contain precisely positioned atoms that alter the electron distributions in those atoms that participate directly in the making and breaking of covalent bonds. Peptide bonds, for example, can be hydrolyzed in the absence of an enzyme by exposing a polypeptide to either a strong acid or a strong base, as explained in Figure 3–49. Enzymes are unique, however, in being able to use acid and base catalysis simultaneously, since the acidic and basic residues required are prevented from combining with each other (as they would do in solution) by being tied to the rigid framework of the protein itself (Figure 3–49D).

The fit between an enzyme and its substrate needs to be precise. A small change introduced by genetic engineering in the active site of an enzyme can have a profound effect. Replacing a glutamic acid with an aspartic acid in one enzyme, for example, shifts the position of the catalytic carboxylate ion by only 1 Å (about the radius of a hydrogen atom); yet this is enough to decrease the activity of the enzyme a thousandfold.





Figure 3–46 Enzymatic acceleration of chemical reactions by decreasing the activation energy. Often both the uncatalyzed reaction (A) and the enzyme-catalyzed reaction (B) can go through several transition states. It is the transition state with the highest energy (S^T and ES^T) that determines the activation energy and limits the rate of the reaction. (S = substrate; P = product of the reaction.)

(A) HYDROLYSIS OF AN AMIDE BOND



(B) TRANSITION-STATE ANALOG FOR AMIDE HYDROLYSIS




Figure 3–47 Catalytic antibodies. The stabilization of a transition state by an antibody creates an enzyme. (A) The reaction path for the hydrolysis of an amide bond goes through a tetrahedral intermediate, the high-energy transition state for the reaction. (B) The molecule on the left was covalently linked to a protein and used as an antigen to generate an antibody that binds tightly to the region of the molecule shown in yellow. Because this antibody also bound tightly to the transition state in (A), it was found to function as an enzyme that efficiently catalyzed the hydrolysis of the amide bond in the molecule on the right.




Figure 3-48 The rate accelerations caused by five different enzymes. (Modified from A. Radzicka and R. Wolfenden, *Science* 267:90-93, 1995.)


Lysozyme Illustrates How an Enzyme Works


To demonstrate how enzymes catalyze chemical reactions, we shall use the example of an enzyme that acts as a natural antibiotic in egg white, saliva, tears, and other secretions. **Lysozyme** is an enzyme that catalyzes the cutting of polysaccharide chains in the cell walls of bacteria. Because the bacterial cell is under pressure from osmotic forces, cutting even a small number of polysaccharide chains causes the cell wall to rupture and the cell to burst. Lysozyme is a relatively small and stable protein that can be easily isolated in large quantities. For these reasons, it has been intensively studied, and it was the first enzyme to have its structure worked out in atomic detail by x-ray crystallography.

The reaction catalyzed by lysozyme is a hydrolysis: a molecule of water is added to a single bond between two adjacent sugar groups in the polysaccharide chain, thereby causing the bond to break (see Figure 2-19). The reaction is energetically favorable because the free energy of the severed polysaccharide chain is lower than the free energy of the intact chain. However, the pure polysaccharide can sit for years in water without being hydrolyzed to any detectable degree. This is because there is an energy barrier to the reaction, as discussed in Chapter 2 (see Figure 2-46). For a colliding water molecule to break a bond linking two sugars, the polysaccharide molecule has to be distorted into a particular shape—the transition state—in which the atoms around the bond have an altered geometry and electron distribution. Because of this distortion, a large activation energy must be supplied through random collisions before the reaction can take place. In an aqueous solution at room temperature, the energy of collisions almost never exceeds the activation energy. Consequently, hydrolysis occurs extremely slowly, if at all.

This situation is drastically changed when the polysaccharide binds to lysozyme. The active site of lysozyme, because its substrate is a polymer, is a long groove that holds six linked sugars at the same time. As soon as the polysaccharide binds to form an enzyme-substrate complex, the enzyme cuts the polysaccharide by adding a water molecule across one of its sugar-sugar bonds. The product chains are then quickly released, freeing the enzyme for further cycles of reaction (Figure 3-50).

Figure 3-49 Acid catalysis and base catalysis. (A) The start of the uncatalyzed reaction shown in Figure 3-47A is diagrammed, with blue indicating electron distribution in the water and carbonyl bonds. (B) An acid likes to donate a proton (H^+) to other atoms. By pairing with the carbonyl oxygen, an acid causes electrons to move away from the carbonyl carbon, making this atom much more attractive to the electronegative oxygen of an attacking water molecule. (C) A base likes to take up H^+ . By pairing with a hydrogen of the attacking water molecule, a base causes electrons to move toward the water oxygen, making it a better attacking group for the carbonyl carbon. (D) By having appropriately positioned atoms on its surface, an enzyme can perform both acid catalysis and base catalysis at the same time.





The chemistry that underlies the binding of lysozyme to its substrate is the same as that for antibody binding to its antigen—the formation of multiple non-covalent bonds. However, lysozyme holds its polysaccharide substrate in a particular way, so that one of the two sugars involved in the bond to be broken is distorted from its normal, most stable conformation. The bond to be broken is also held close to two amino acids with acidic side chains (a glutamic acid and an aspartic acid) within the active site.

Conditions are thereby created in the microenvironment of the lysozyme active site that greatly reduce the activation energy necessary for the hydrolysis to take place. Figure 3–51 shows the stages proposed in 1967 for this enzymatically catalyzed reaction.


1. The enzyme stresses its bound substrate by bending some critical chemical bonds that will participate in the chemical reaction, so that the shape of the substrate more closely resembles the shape of the high-energy transition state formed halfway through the reaction.
2. A precisely positioned acidic side chain of the glutamic acid within the active site speeds up the hydrolysis by providing a high concentration of acidifying H^+ ions (acid catalysis), even though the solution surrounding the enzyme is at neutral pH.
3. The negatively charged aspartic acid further stabilizes the positively charged transition state (base catalysis).

As indicated in the Figure legend, this classical view of the lysozyme reaction is in need of modification. In particular, a tetrahedral intermediate forms in the transition state—created by the transient covalent addition of the aspartic acid to the carbon atom at the point of cleavage.

Similar mechanisms are used by other enzymes to lower activation energies and speed up the reactions they catalyze. In reactions involving two or more reactants, the active site also acts like a template, or mold, that brings the substrates together in the proper orientation for a reaction to occur between them (Figure 3–52A). As we saw for lysozyme, the active site of an enzyme contains precisely positioned atoms that speed up a reaction by using charged groups to alter the distribution of electrons in the substrates (Figure 3–52B). The binding to the enzyme also changes substrate shapes, bending bonds so as to drive a substrate toward a particular transition state (Figure 3–52C). Finally, like lysozyme, many enzymes participate intimately in the reaction by briefly forming a covalent bond between the substrate and a side chain of the enzyme. Subsequent steps in the reaction restore the side chain to its original state, so that the enzyme remains unchanged after the reaction (see Figure 2–73).


Tightly Bound Small Molecules Add Extra Functions to Proteins

Although we have emphasized the versatility of proteins as chains of amino acids that perform different functions, there are many instances in which the amino acids by themselves are not enough. Just as humans employ tools to enhance and extend the capabilities of their hands, proteins often use small nonprotein molecules to perform functions that would be difficult or impossible to do with amino acids alone. Thus, the signal receptor protein *rhodopsin*, which is made by the photoreceptor cells in the retina, detects light by means of a small molecule, *retinal*, embedded in the protein (Figure 3–53A). Retinal changes its shape when it absorbs a photon of light, and this change causes the protein to trigger a cascade of enzymatic reactions that eventually leads to an electrical signal being carried to the brain.



(B)

Figure 3–50 The reaction catalyzed by lysozyme. (A) The enzyme lysozyme (denoted as E) catalyzes the cutting of a polysaccharide chain, which is its substrate (S). The enzyme first binds to the chain to form an enzyme–substrate complex (ES) and then catalyzes the cleavage of a specific covalent bond in the backbone of the polysaccharide, forming an enzyme–product complex (EP) that rapidly dissociates. Release of the severed chain (the products P) leaves the enzyme free to act on another substrate molecule. (B) A space-filling model of the lysozyme molecule bound to a short length of polysaccharide chain before cleavage. (B, courtesy of Richard J. Feldmann.)



In the enzyme-substrate complex (ES), the enzyme forces sugar D into a strained conformation, with Glu 35 of lysozyme positioned to serve as an acid that attacks the adjacent sugar-sugar bond by donating a proton (H^+).

This is the unstable transition state, with a positive charge on sugar D. Both the strain on sugar D and the nearby negative charge on Asp 52 of lysozyme stabilize this intermediate, greatly lowering its energy on the enzyme surface.

The rapid addition of a water molecule (green) completes the hydrolysis and regenerates the protonated form of Glu 35, forming the enzyme-product complex (EP).


Another example of a protein that contains a nonprotein portion is hemoglobin (see Figure 3-23). A molecule of hemoglobin carries four *heme* groups, ring-shaped molecules each with a single central iron atom (Figure 3-53B). Heme gives hemoglobin (and blood) its red color. By binding reversibly to oxygen gas through its iron atom, heme enables hemoglobin to pick up oxygen in the lungs and release it in the tissues.

Sometimes these small molecules are attached covalently and permanently to their protein, thereby becoming an integral part of the protein molecule itself. We see in Chapter 10 that proteins are often anchored to cell membranes through covalently attached lipid molecules. And membrane proteins exposed on the surface of the cell, as well as proteins secreted outside the cell, are often modified by the covalent addition of sugars and oligosaccharides.

Enzymes frequently have a small molecule or metal atom tightly associated with their active site that assists with their catalytic function. *Carboxypeptidase*, for example, an enzyme that cuts polypeptide chains, carries a tightly bound zinc ion in its active site. During the cleavage of a peptide bond by carboxypeptidase, the zinc ion forms a transient bond with one of the substrate atoms, thereby assisting the hydrolysis reaction. In other enzymes, a small organic

Figure 3-51 The classical model for events at the active site of lysozyme.

The top left and top right drawings depict the free substrate and the free products, respectively, whereas the other three drawings depict the sequential events proposed at the enzyme active site. Note the change in the conformation of sugar D in the enzyme-substrate complex; this sugar has been postulated to acquire a positive charge in the unstable transition state, as shown in the central panel. However, very recent data has revealed that this standard textbook view of the reaction is incorrect. Instead, the Asp 52 forms a covalent bond to the carbon shown here with a positive charge. This covalent bond is rapidly broken in the next step, as water is added to complete the reaction.




(A) enzyme binds to two substrate molecules and orients them precisely to encourage a reaction to occur between them

(B) binding of substrate to enzyme rearranges electrons in the substrate, creating partial negative and positive charges that favor a reaction

(C) enzyme strains the bound substrate molecule, forcing it toward a transition state to favor a reaction

Figure 3-52 Some general strategies of enzyme catalysis. (A) Holding substrates together in a precise alignment. (B) Charge stabilization of reaction intermediates. (C) Altering bond angles in the substrate to increase the rate of a particular reaction.

**Figure 3-53 Retinal and heme.**

(A) The structure of retinal, the light-sensitive molecule attached to rhodopsin in the eye. (B) The structure of a heme group. The carbon-containing heme ring is red and the iron atom at its center is orange. A heme group is tightly bound to each of the four polypeptide chains in hemoglobin, the oxygen-carrying protein whose structure is shown in Figure 3-23.

molecule serves a similar purpose. Such organic molecules are often referred to as **coenzymes**. An example is *biotin*, which is found in enzymes that transfer a carboxylate group ($-\text{COO}^-$) from one molecule to another (see Figure 2-63). Biotin participates in these reactions by forming a transient covalent bond to the $-\text{COO}^-$ group to be transferred, being better suited to this function than any of the amino acids used to make proteins. Because it cannot be synthesized by humans, and must therefore be supplied in small quantities in our diet, biotin is a *vitamin*, as are many other coenzymes (Table 3-2). Other vitamins are needed to make other small molecules that are essential components of our proteins; vitamin A, for example, is needed in the diet to make retinal, the light-sensitive part of rhodopsin.

Multienzyme Complexes Help to Increase the Rate of Cell Metabolism

The efficiency of enzymes in accelerating chemical reactions is crucial to the maintenance of life. Cells, in effect, must race against the unavoidable processes of decay, which—if left unattended—cause macromolecules to run downhill toward greater and greater disorder. If the rates of desirable reactions were not greater than the rates of competing side reactions, a cell would soon die. Some idea of the rate at which cell metabolism proceeds can be obtained by measuring the rate of ATP utilization. A typical mammalian cell “turns over” (i.e., hydrolyzes and restores by phosphorylation) its entire ATP pool once every 1 or 2 minutes. For each cell this turnover represents the utilization of roughly 10^7 molecules of ATP per second (or, for the human body, about 1 gram of ATP every minute).

The rates of reactions in cells are rapid because of the effectiveness of enzyme catalysis. Many important enzymes have become so efficient that there

TABLE 3-2 Many Vitamins Provide Critical Coenzymes for Human Cells

VITAMIN	COENZYME	ENZYME-CATALYZED REACTIONS REQUIRING THESE COENZYMES
Thiamine (vitamin B ₁)	thiamine pyrophosphate	activation and transfer of aldehydes
Riboflavin (vitamin B ₂)	FADH	oxidation–reduction
Niacin	NADH, NADPH	oxidation–reduction
Pantothenic acid	coenzyme A	acyl group activation and transfer
Pyridoxine	pyridoxal phosphate	reactions involving amino acid activation
Biotin	biotin	CO ₂ activation and transfer
Lipoic acid	lipoamide	acyl group activation; oxidation–reduction
Folic acid	tetrahydrofolate	activation and transfer of single carbon groups
Vitamin B ₁₂	cobalamin coenzymes	isomerization and methyl group transfers

is no possibility of further useful improvement. The factor that limits the reaction rate is no longer the enzyme's intrinsic speed of action; rather, it is the frequency with which the enzyme collides with its substrate. Such a reaction is said to be *diffusion-limited*.

If an enzyme-catalyzed reaction is diffusion-limited, its rate depends on the concentration of both the enzyme and its substrate. If a sequence of reactions is to occur extremely rapidly, each metabolic intermediate and enzyme involved must be present in high concentration. However, given the enormous number of different reactions performed by a cell, there are limits to the concentrations of substrates that can be achieved. In fact, most metabolites are present in micromolar (10^{-6} M) concentrations, and most enzyme concentrations are much lower. How is it possible, therefore, to maintain very fast metabolic rates?

The answer lies in the spatial organization of cell components. Reaction rates can be increased without raising substrate concentrations by bringing the various enzymes involved in a reaction sequence together to form a large protein assembly known as a *multienzyme complex* (Figure 3-54). Because this allows the product of enzyme A to be passed directly to enzyme B, and so on, diffusion rates need not be limiting, even when the concentrations of the substrates in the cell as a whole are very low. It is perhaps not surprising, therefore, that such enzyme complexes are very common, and they are involved in nearly all aspects of metabolism—including the central genetic processes of DNA, RNA, and protein synthesis. In fact, few enzymes in eucaryotic cells may be left to diffuse freely in solution; instead, most seem to have evolved binding sites that concentrate them with other proteins of related function in particular regions of the cell, thereby increasing the rate and efficiency of the reactions that they catalyze.

Eucaryotic cells have yet another way of increasing the rate of metabolic reactions, using their intracellular membrane systems. These membranes can segregate particular substrates and the enzymes that act on them into the same membrane-enclosed compartment, such as the endoplasmic reticulum or the cell nucleus. If, for example, a compartment occupies a total of 10% of the volume of the cell, the concentration of reactants in the compartment may be increased as much as 10 times compared with the same cell with no compartmentalization. Reactions that would otherwise be limited by the speed of diffusion can thereby be speeded up by a factor of 10.

The Catalytic Activities of Enzymes Are Regulated

A living cell contains thousands of enzymes, many of which operate at the same time and in the same small volume of the cytosol. By their catalytic action, these enzymes generate a complex web of metabolic pathways, each composed of chains of chemical reactions in which the product of one enzyme becomes the substrate of the next. In this maze of pathways, there are many branch points where different enzymes compete for the same substrate. The system is so complex (see Figure 2-88) that elaborate controls are required to regulate when and how rapidly each reaction occurs.




Figure 3-54 The structure of pyruvate dehydrogenase. This enzyme complex catalyzes the conversion of pyruvate to acetyl CoA, as part of the pathway that oxidizes sugars to CO_2 and H_2O . It is an example of a large multienzyme complex in which reaction intermediates are passed directly from one enzyme to another.

Regulation occurs at many levels. At one level, the cell controls how many molecules of each enzyme it makes by regulating the expression of the gene that encodes that enzyme (discussed in Chapter 7). The cell also controls enzymatic activities by confining sets of enzymes to particular subcellular compartments, enclosed by distinct membranes (discussed in Chapters 12 and 14). The rate of protein destruction by targeted proteolysis represents yet another important regulatory mechanism (see p. 361). But the most rapid and general process that adjusts reaction rates operates through a direct, reversible change in the activity of an enzyme in response to specific molecules that it encounters.

The most common type of control occurs when a molecule other than one of the substrates binds to an enzyme at a special regulatory site outside the active site, thereby altering the rate at which the enzyme converts its substrates to products. In **feedback inhibition**, an enzyme acting early in a reaction pathway is inhibited by a late product of that pathway. Thus, whenever large quantities of the final product begin to accumulate, this product binds to the first enzyme and slows down its catalytic action, thereby limiting the further entry of substrates into that reaction pathway (Figure 3–55). Where pathways branch or intersect, there are usually multiple points of control by different final products, each of which works to regulate its own synthesis (Figure 3–56). Feedback inhibition can work almost instantaneously and is rapidly reversed when the level of the product falls.

Feedback inhibition is *negative regulation*: it prevents an enzyme from acting. Enzymes can also be subject to *positive regulation*, in which the enzyme's activity is stimulated by a regulatory molecule rather than being shut down. Positive regulation occurs when a product in one branch of the metabolic maze stimulates the activity of an enzyme in another pathway. As one example, the accumulation of ADP activates several enzymes involved in the oxidation of sugar molecules, thereby stimulating the cell to convert more ADP to ATP.

Allosteric Enzymes Have Two or More Binding Sites That Interact

One feature of feedback inhibition was initially puzzling to those who discovered it: the regulatory molecule often has a shape totally different from the shape of the substrate of the enzyme. This is why this form of regulation is termed *allostery* (from the Greek words *allos*, meaning “other,” and *stereos*, meaning “solid” or “three-dimensional”). As more was learned about feedback inhibition, it was recognized that many enzymes must have at least two different binding sites on their surface—an **active site** that recognizes the substrates, and a **regulatory site** that recognizes a regulatory molecule. These two sites must somehow communicate in a way that allows the catalytic events at the active site to be influenced by the binding of the regulatory molecule at its separate site on the protein's surface.

The interaction between separated sites on a protein molecule is now known to depend on a *conformational change* in the protein: binding at one of the sites causes a shift from one folded shape to a slightly different folded shape. During feedback inhibition, for example, the binding of an inhibitor at one site on the protein causes the protein to shift to a conformation in which its active site—located elsewhere in the protein—becomes incapacitated.

It is thought that most protein molecules are allosteric. They can adopt two or more slightly different conformations, and a shift from one to another caused by the binding of a ligand can alter their activity. This is true not only for enzymes but also for many other proteins—including receptors, structural proteins, and motor proteins. There is nothing mysterious about the allosteric regulation of these proteins: each conformation of the protein has somewhat different surface contours, and the protein's binding sites for ligands are altered when the protein changes shape. Moreover as we discuss next, each ligand stabilizes the conformation that it binds to most strongly, and thus—at high enough concentrations—tends to “switch” the protein toward the conformation it prefers.




Figure 3–55 Feedback inhibition of a single biosynthetic pathway. The end-product Z inhibits the first enzyme that is unique to its synthesis and thereby controls its own level in the cell. This is an example of negative regulation.




Figure 3-56 Multiple feedback inhibition. In this example, which shows the biosynthetic pathways for four different amino acids in bacteria, the red arrows indicate positions at which products feed back to inhibit enzymes. Each amino acid controls the first enzyme specific to its own synthesis, thereby controlling its own levels and avoiding a wasteful buildup of intermediates. The products can also separately inhibit the initial set of reactions common to all the syntheses; in this case, three different enzymes catalyze the initial reaction, each inhibited by a different product.

Two Ligands Whose Binding Sites Are Coupled Must Reciprocally Affect Each Other's Binding

The effects of ligand binding on a protein follow from a fundamental chemical principle known as **linkage**. Suppose, for example, that a protein that binds glucose also binds another molecule, X, at a distant site on the protein's surface. If the binding site for X changes shape as part of the conformational change induced by glucose binding, the binding sites for X and for glucose are said to be *coupled*. Whenever two ligands prefer to bind to the *same* conformation of an allosteric protein, it follows from basic thermodynamic principles that each ligand must increase the affinity of the protein for the other. Thus, if the shift of the protein in Figure 3-57 to the closed conformation that binds glucose best also causes the binding site for X to fit X better, then the protein will bind glucose more tightly when X is present than when X is absent.

Conversely, linkage operates in a negative way if two ligands prefer to bind to *different* conformations of the same protein. In this case, the binding of the first ligand discourages the binding of the second ligand. Thus, if a shape change caused by glucose binding decreases the affinity of a protein for molecule X, the binding of X must also decrease the protein's affinity for glucose (Figure 3-58). The linkage relationship is quantitatively reciprocal, so that, for example, if glucose has a very large effect on the binding of X, X has a very large effect on the binding of glucose.

The relationships shown in Figures 3-57 and 3-58 underlie all of cell biology. They seem so obvious in retrospect that we now take them for granted. But their discovery in the 1950s, followed by a general description of allostery in the early 1960s, was revolutionary at the time. Since molecule X in these examples




Figure 3-57 Positive regulation caused by conformational coupling between two distant binding sites. In this example, both glucose and molecule X bind best to the closed conformation of a protein with two domains. Because both glucose and molecule X drive the protein toward its closed conformation, each ligand helps the other to bind. Glucose and molecule X are therefore said to bind cooperatively to the protein.

binds at a site that is distinct from the site where catalysis occurs, it need have no chemical relationship to glucose or to any other ligand that binds at the active site. As we have just seen, for enzymes that are regulated in this way, molecule X can either turn the enzyme on (positive regulation) or turn it off (negative regulation). By such a mechanism, **allosteric proteins** serve as general switches that, in principle, allow one molecule in a cell to affect the fate of any other.

Symmetric Protein Assemblies Produce Cooperative Allosteric Transitions

A single subunit enzyme that is regulated by negative feedback can at most decrease from 90% to about 10% activity in response to a 100-fold increase in the concentration of an inhibitory ligand that it binds (Figure 3-59, red line). Responses of this type are apparently not sharp enough for optimal cell regulation, and most enzymes that are turned on or off by ligand binding consist of symmetric assemblies of identical subunits. With this arrangement, the binding of a molecule of ligand to a single site on one subunit can trigger an allosteric change in the subunit that can be transmitted to the neighboring subunits, helping them to bind the same ligand. As a result, a *cooperative allosteric transition* occurs (Figure 3-59, blue line), allowing a relatively small change in ligand concentration in the cell to switch the whole assembly from an almost fully active to an almost fully inactive conformation (or vice versa).

The principles involved in a cooperative “all-or-none” transition are easiest to visualize for an enzyme that forms a symmetric dimer. In the example shown in Figure 3-60, the first molecule of an inhibitory ligand binds with great difficulty since its binding destroys an energetically favorable interaction between the two identical monomers in the dimer. A second molecule of inhibitory ligand now binds more easily, however, because its binding restores the




Figure 3-58 Negative regulation caused by conformational coupling between two distant binding sites. The scheme here resembles that in the previous figure, but here molecule X prefers the open conformation, while glucose prefers the closed conformation. Because glucose and molecule X drive the protein toward opposite conformations (closed and open, respectively), the presence of either ligand interferes with the binding of the other.




Figure 3-59 Enzyme activity versus the concentration of inhibitory ligand for single-subunit and multi-subunit allosteric enzymes. For an enzyme with a single subunit (red line), a drop from 90% enzyme activity to 10% activity (indicated by the two dots on the curve) requires a 100-fold increase in the concentration of inhibitor. The enzyme activity is calculated from the simple equilibrium relationship $K = \frac{[I][P]}{[IP]}$, where P is active protein, I is inhibitor, and IP is the inactive protein bound to inhibitor. An identical curve applies to any simple binding interaction between two molecules, A and B. In contrast, a multisubunit allosteric enzyme can respond in a switchlike manner to a change in ligand concentration: the steep response is caused by a cooperative binding of the ligand molecules, as explained in Figure 3-60. Here, the green line represents the idealized result expected for the cooperative binding of two inhibitory ligand molecules to an allosteric enzyme with two subunits, and the blue line shows the idealized response of an enzyme with four subunits. As indicated by the two dots on each of these curves, the more complex enzymes drop from 90% to 10% activity over a much narrower range of inhibitor concentration than does the enzyme composed of a single subunit.

monomer–monomer contacts of a symmetric dimer (and also completely inactivates the enzyme).

An even sharper response to a ligand can be obtained with larger assemblies, such as the enzyme formed from 12 polypeptide chains that we discuss next.


The Allosteric Transition in Aspartate Transcarbamoylase Is Understood in Atomic Detail

One enzyme used in the early studies of allosteric regulation was aspartate transcarbamoylase from *E. coli*. It catalyzes the important reaction that begins the synthesis of the pyrimidine ring of C, U, and T nucleotides: carbamoylphosphate + aspartate → N-carbamoylaspartate. One of the final products of this pathway, cytosine triphosphate (CTP), binds to the enzyme to turn it off whenever CTP is plentiful.

Aspartate transcarbamoylase is a large complex of six regulatory and six catalytic subunits. The catalytic subunits are present as two trimers, each arranged like an equilateral triangle; the two trimers face each other and are held together by three regulatory dimers that form a bridge between them. The entire molecule is poised to undergo a concerted, all-or-none, allosteric transition between two conformations, designated as T (tense) and R (relaxed) states (Figure 3-61).

The binding of substrates (carbamoylphosphate and aspartate) to the catalytic trimers drives aspartate transcarbamoylase into its catalytically active R state, from which the regulatory CTP molecules dissociate. By contrast, the binding of CTP to the regulatory dimers converts the enzyme to the inactive T state, from which the substrates dissociate. This tug-of-war between CTP and substrates is identical in principle to that described previously in Figure 3-58 for a simpler allosteric protein. But because the tug-of-war occurs in a symmetric molecule with multiple binding sites, the enzyme undergoes a cooperative allosteric transition that will turn it on suddenly as substrates accumulate

Figure 3-60 A cooperative allosteric transition in an enzyme composed of two identical subunits. This diagram illustrates how the conformation of one subunit can influence that of its neighbor. The binding of a single molecule of an inhibitory ligand (yellow) to one subunit of the enzyme occurs with difficulty because it changes the conformation of this subunit and thereby destroys the symmetry of the enzyme. Once this conformational change has occurred, however, the energy gained by restoring the symmetric pairing interaction between the two subunits makes it especially easy for the second subunit to bind the inhibitory ligand and undergo the same conformational change. Because the binding of the first molecule of ligand increases the affinity with which the other subunit binds the same ligand, the response of the enzyme to changes in the concentration of the ligand is much steeper than the response of an enzyme with only one subunit (see Figure 3-59).






Figure 3-61 The transition between R and T states in the enzyme aspartate transcarbamoylase. The enzyme consists of a complex of six catalytic subunits and six regulatory subunits, and the structures of its inactive (T state) and active (R state) forms have been determined by x-ray crystallography. The enzyme is turned off by feedback inhibition when CTP concentrations rise. Each regulatory subunit can bind one molecule of CTP, which is one of the final products in the pathway. By means of this negative feedback regulation, the pathway is prevented from producing more CTP than the cell needs. (Based on K.L. Krause, K.W. Volz, and W.N. Lipscomb, *Proc. Natl. Acad. Sci. USA* 82:1643-1647, 1985.)

(forming the R state) or shut it off rapidly when CTP accumulates (forming the T state).

A combination of biochemistry and x-ray crystallography has revealed many fascinating details of this allosteric transition. Each regulatory subunit has two domains, and the binding of CTP causes the two domains to move relative to each other, so that they function like a lever that rotates the two catalytic trimers and pulls them closer together into the T state (see Figure 3-61). When this occurs, hydrogen bonds form between opposing catalytic subunits that help widen the cleft that forms the active site within each catalytic subunit, thereby destroying the binding sites for the substrates (Figure 3-62). Adding large amounts of substrate has the opposite effect, favoring the R state by binding in the cleft of each catalytic subunit and opposing the above conformational change. Conformations that are intermediate between R and T are unstable, so that the enzyme mostly clicks back and forth between its R and T forms, producing a mixture of these two species in proportions that depend on the relative concentrations of CTP and substrates.

Many Changes in Proteins Are Driven by Phosphorylation

Enzymes are regulated by more than the binding of small molecules. A second method that is commonly used by eucaryotic cells to regulate a protein's function is the covalent addition of a phosphate group to one of its amino acid side chains. Such *phosphorylation* events can affect the protein in two important ways.

First, because each phosphate group carries two negative charges, the enzyme-catalyzed addition of a phosphate group to a protein can cause a major conformational change in the protein by, for example, attracting a cluster of positively charged amino acid side chains. This can, in turn, affect the binding of ligands elsewhere on the protein surface, dramatically changing the protein's activity through an allosteric effect. Removal of the phosphate group by a second enzyme returns the protein to its original conformation and restores its initial activity.




Figure 3-62 Part of the on-off switch in the catalytic subunits of aspartate transcarbamoylase. Changes in the indicated hydrogen-bonding interactions are partly responsible for switching this enzyme's active site between active (yellow) and inactive conformations. Hydrogen bonds are indicated by thin red lines. The amino acids involved in the subunit-subunit interaction are shown in red, while those that form the active site of the enzyme are shown in blue. The large drawings show the catalytic site in the interior of the enzyme; the boxed sketches show the same subunits viewed from the enzyme's external surface. (Adapted from E.R. Kantrowitz and W.N. Lipscomb, *Trends Biochem. Sci.* 15:53-59, 1990.)


Second, an attached phosphate group can form part of a structure that is directly recognized by binding sites of other proteins. As previously discussed, certain small protein domains, called modules, appear very frequently in larger proteins. A large number of these modules provide binding sites for attaching their protein to phosphorylated peptides in other protein molecules. One of these modules is the SH2 domain, featured previously in this chapter, which binds to a short peptide sequence containing a phosphorylated tyrosine side chain (see Figure 3-40B). Several other types of modules recognize phosphorylated serine or threonine side chains in a specific context. As a result, protein phosphorylation and dephosphorylation events have a major role in driving the regulated assembly and disassembly of protein complexes.

Through such effects, reversible protein phosphorylation controls the activity structure and cellular localization of many types of proteins in eucaryotic cells. In fact, this regulation is so extensive that more than one-third of the 10,000 or so proteins in a typical mammalian cell are thought to be phosphorylated at any given time—many with more than one phosphate. As might be expected, the addition and removal of phosphate groups from specific proteins often occur in response to signals that specify some change in a cell's state. For example, the complicated series of events that takes place as a eucaryotic cell divides is largely timed in this way (discussed in Chapter 17), and many of the signals mediating cell-cell interactions are relayed from the plasma membrane to the nucleus by a cascade of protein phosphorylation events (discussed in Chapter 15).

A Eucaryotic Cell Contains a Large Collection of Protein Kinases and Protein Phosphatases

Protein phosphorylation involves the enzyme-catalyzed transfer of the terminal phosphate group of an ATP molecule to the hydroxyl group on a serine, threonine, or tyrosine side chain of the protein (Figure 3-63). This reaction is catalyzed by a **protein kinase**, and the reaction is essentially unidirectional because

Figure 3–63 Protein phosphorylation. Many thousands of proteins in a typical eucaryotic cell are modified by the covalent addition of a phosphate group. (A) The general reaction, shown here, entails the transfer of a phosphate group from ATP to an amino acid side chain of the target protein by a protein kinase. Removal of the phosphate group is catalyzed by a second enzyme, a protein phosphatase. In this example, the phosphate is added to a serine side chain; in other cases, the phosphate is instead linked to the $-OH$ group of a threonine or a tyrosine in the protein. (B) The phosphorylation of a protein by a protein kinase can either increase or decrease the protein's activity, depending on the site of phosphorylation and the structure of the protein.



of the large amount of free energy released when the phosphate–phosphate bond in ATP is broken to produce ADP (discussed in Chapter 2). The reverse reaction of phosphate removal, or *dephosphorylation*, is instead catalyzed by a **protein phosphatase**. Cells contain hundreds of different protein kinases, each responsible for phosphorylating a different protein or set of proteins. There are also many different protein phosphatases; some of these are highly specific and remove phosphate groups from only one or a few proteins, whereas others act on a broad range of proteins and are targeted to specific substrates by regulatory subunits. The state of phosphorylation of a protein at any moment, and thus its activity, depends on the relative activities of the protein kinases and phosphatases that modify it.

The protein kinases that phosphorylate proteins in eucaryotic cells belong to a very large family of enzymes, which share a catalytic (kinase) sequence of 250 amino acids. The various family members contain different amino acid sequences on either side of the kinase sequence (see Figure 3–12), and often have short amino acid sequences inserted into loops within it (*red arrowheads* in Figure 3–64). Some of these additional amino acid sequences enable each kinase to recognize the specific set of proteins it phosphorylates, or to bind to structures that localize it in specific regions of the cell. Other parts of the protein allow the activity of each enzyme to be tightly regulated, so it can be turned on and off in response to different specific signals, as described below.

By comparing the number of amino acid sequence differences between the various members of a protein family, one can construct an “evolutionary tree” that is thought to reflect the pattern of gene duplication and divergence that gave rise to the family. An evolutionary tree of protein kinases is shown in Figure 3–65. Not surprisingly, kinases with related functions are often located on nearby branches of the tree: the protein kinases involved in cell signaling that phosphorylate tyrosine side chains, for example, are all clustered in the top left corner of the tree. The other kinases shown phosphorylate either a serine or a threonine side chain, and many are organized into clusters that seem to reflect their function—in transmembrane signal transduction, intracellular signal amplification, cell-cycle control, and so on.

As a result of the combined activities of protein kinases and protein phosphatases, the phosphate groups on proteins are continually turning over—being added and then rapidly removed. Such phosphorylation cycles may seem wasteful, but they are important in allowing the phosphorylated proteins to switch rapidly from one state to another: the more rapid the cycle, the faster the state of phosphorylation of a population of protein molecules can change in response to a sudden stimulus that changes the phosphorylation rate (see Figure 15–10). The




Figure 3–64 The three-dimensional structure of a protein kinase.

Superimposed on this structure are *red arrowheads* to indicate sites where insertions of 5–100 amino acids are found in some members of the protein kinase family. These insertions are located in loops on the surface of the enzyme where other ligands interact with the protein. Thus, they distinguish different kinases and confer on them distinctive interactions with other proteins. The ATP (which donates a phosphate group) and the peptide to be phosphorylated are held in the active site, which extends between the phosphate-binding loop (*yellow*) and the catalytic loop (*orange*). See also Figure 3–12. (Adapted from D.R. Knighton et al., *Science* 253:407–414, 1991.)




Figure 3-65 An evolutionary tree of selected protein kinases. Although a higher eucaryotic cell contains hundreds of such enzymes, and the human genome codes for more than 500, only some of those discussed in this book are shown.

energy required to drive the phosphorylation cycle is derived from the free energy of ATP hydrolysis, one molecule of which is consumed for each phosphorylation event.

The Regulation of Cdk and Src Protein Kinases Shows How a Protein Can Function as a Microchip

The hundreds of different protein kinases in a eucaryotic cell are organized into complex networks of signaling pathways that help to coordinate the cell's activities, drive the cell cycle, and relay signals into the cell from the cell's environment. Many of the extracellular signals involved need to be both integrated and amplified by the cell. Individual protein kinases (and other signaling proteins) serve as input-output devices, or "microchips," in the integration process. An important part of the input to these proteins comes from the control that is exerted by phosphates added and removed from them by protein kinases and protein phosphatases, respectively, in the signaling network.

For a protein that is phosphorylated at multiple sites, specific sets of phosphate groups serve to activate the protein, while other sets can inactivate it. A cyclin-dependent protein kinase (Cdk) represents a good example of such a signal processing device. Kinases in this class phosphorylate serines, and they are central components of the cell-cycle control system in eucaryotic cells, as discussed in detail in Chapter 17. In a vertebrate cell, individual Cdk enzymes turn on and off in succession, as a cell proceeds through the different phases of its division cycle. When a particular one of these kinases is on, it influences various aspects of cell behavior through effects on the proteins it phosphorylates.

A Cdk protein becomes active as a serine/threonine protein kinase only when it is bound to a second protein called a *cyclin*. But, as Figure 3-66 shows, the binding of cyclin is only one of three distinct "inputs" required to activate the Cdk. In addition to cyclin binding, a phosphate must be added to a specific threonine side chain, and a phosphate elsewhere in the protein (covalently bound to a specific tyrosine side chain) must be removed. Cdk thus monitors a specific set of cell components—a cyclin, a protein kinase, and a protein phosphatase—and it acts as an input-output device that turns on if, and only if, each of these components has attained its appropriate activity state. Some cyclins rise and fall in concentration in step with the cell cycle, increasing gradually in amount until they are suddenly destroyed at a particular point in the cycle. The sudden destruction of a cyclin (by targeted proteolysis) immediately shuts off its partner Cdk enzyme, and this triggers a specific step in the cell cycle.




Figure 3-66 How a Cdk protein acts as an integrating device. The function of these central regulators of the cell cycle is discussed in Chapter 17.




Figure 3-67 The domain structure of the Src family of protein kinases, mapped along the amino acid sequence.

A similar type of microchip behavior is exhibited by the Src family of protein kinases. The *Src* protein (pronounced “sarc”) was the first tyrosine kinase to be discovered, and it is now known to be part of a subfamily of nine very similar protein kinases, which are found only in multicellular animals. As indicated by the evolutionary tree in Figure 3-65, sequence comparisons suggest that tyrosine kinases as a group were a relatively late innovation that branched off from the serine/threonine kinases, with the Src subfamily being only one subgroup of the tyrosine kinases created in this way.

The Src protein and its homologs contain a short N-terminal region that becomes covalently linked to a strongly hydrophobic fatty acid, which holds the kinase at the cytoplasmic face of the plasma membrane. Next come two peptide-binding modules, a Src homology 3 (SH3) domain and a SH2 domain, followed by the kinase catalytic domains (Figure 3-67). These kinases normally exist in an inactive conformation, in which a phosphorylated tyrosine near the C-terminus is bound to the SH2 domain, and the SH3 domain is bound to an internal peptide in a way that distorts the active site of the enzyme and helps to render it inactive (see Figure 3-12).

Turning the kinase on involves at least two specific inputs: removal of the C-terminal phosphate and the binding of the SH3 domain by a specific activating protein (Figure 3-68). As for the Cdk protein, the activation of the Src kinase signals that a particular set of separate upstream events has been completed (Figure 3-69). Thus, both the Cdk and Src families of proteins serve as specific signal integrators, helping to generate the complex web of information-processing events that enable the cell to compute logical responses to a complex set of conditions.

Proteins That Bind and Hydrolyze GTP Are Ubiquitous Cellular Regulators

We have described how the addition or removal of phosphate groups on a protein can be used by a cell to control the protein’s activity. In the examples discussed so far, the phosphate is transferred from an ATP molecule to an amino acid side chain of the protein in a reaction catalyzed by a specific protein kinase. Eucaryotic cells also have another way to control protein activity by phosphate addition and removal. In this case, the phosphate is not attached directly to the protein; instead, it is a part of the guanine nucleotide GTP, which binds very tightly to the protein. In general, proteins regulated in this way are in their active conformations with GTP bound. The loss of a phosphate group occurs when the bound GTP is hydrolyzed to GDP in a reaction catalyzed by the protein itself, and in its GDP bound state the protein is inactive. In this way, GTP-binding proteins act as on-off switches whose activity is determined by the presence or absence of an additional phosphate on a bound GDP molecule (Figure 3-70).

GTP-binding proteins (also called **GTPases** because of the GTP hydrolysis they catalyze) constitute a large family of proteins that all contain variations on the same GTP-binding globular domain. When the tightly bound GTP is hydrolyzed to GDP, this domain undergoes a conformational change that inactivates it. The three-dimensional structure of a prototypical member of this family, the monomeric GTPase called Ras, is shown in Figure 3-71.

The *Ras* protein has an important role in cell signaling (discussed in Chapter 15). In its GTP-bound form, it is active and stimulates a cascade of protein phosphorylations in the cell. Most of the time, however, the protein is in its inactive, GDP-bound form. It becomes active when it exchanges its GDP for a GTP molecule in response to extracellular signals, such as growth factors, that bind to receptors in the plasma membrane (see Figure 15-55).





Figure 3-68 The activation of a Src-type protein kinase by two sequential events. (Adapted from I. Moareti, et al., *Nature* 385:650–653, 1997.)

Figure 3-69 How a Src-type protein kinase acts as an integrating device. The disruption of the SH3 domain interaction (green) can involve replacing its binding to the indicated red linker region with a tighter interaction with the Nef protein, as illustrated in Figure 3-68.



Regulatory Proteins Control the Activity of GTP-Binding Proteins by Determining Whether GTP or GDP Is Bound

GTP-binding proteins are controlled by regulatory proteins that determine whether GTP or GDP is bound, just as phosphorylated proteins are turned on and off by protein kinases and protein phosphatases. Thus, Ras is inactivated by a *GTPase-activating protein (GAP)*, which binds to the Ras protein and induces it to hydrolyze its bound GTP molecule to GDP—which remains tightly bound—and inorganic phosphate (P_i)—which is rapidly released. The Ras protein stays in its inactive, GDP-bound conformation until it encounters a *guanine nucleotide exchange factor (GEF)*, which binds to GDP-Ras and causes it to release its GDP. Because the empty nucleotide-binding site is immediately filled by a GTP molecule (GTP is present in large excess over GDP in cells), the GEF activates Ras by *indirectly* adding back the phosphate removed by GTP hydrolysis. Thus, in a sense, the roles of GAP and GEF are analogous to those of a protein phosphatase and a protein kinase, respectively (Figure 3-72).

Large Protein Movements Can Be Generated From Small Ones

The Ras protein belongs to a large superfamily of *monomeric GTPases*, each of which consists of a single GTP-binding domain of about 200 amino acids. Over the course of evolution, this domain has also become joined to larger proteins with additional domains, creating a large family of GTP-binding proteins. Family members include the receptor-associated trimeric G proteins involved in cell signaling (discussed in Chapter 15), proteins regulating the traffic of vesicles between intracellular compartments (discussed in Chapter 13), and proteins that bind to transfer RNA and are required as assembly factors for protein synthesis on the ribosome (discussed in Chapter 6). In each case, an important biological activity is controlled by a change in the protein's conformation that is caused by GTP hydrolysis in a Ras-like domain.

The *EF-Tu protein* provides a good example of how this family of proteins works. EF-Tu is an abundant molecule that serves as an elongation factor (hence the EF) in protein synthesis, loading each amino-acyl tRNA molecule onto the ribosome. The tRNA molecule forms a tight complex with the GTP-bound form of EF-Tu (Figure 3-73). In this complex, the amino acid attached to the tRNA is masked. Unmasking is required for the tRNA to transfer its amino acid in protein synthesis, and it occurs on the ribosome when the GTP bound to EF-Tu is




Figure 3-70 GTP-binding proteins as molecular switches. The activity of a GTP-binding protein (also called a GTPase) generally requires the presence of a tightly bound GTP molecule (switch “on”). Hydrolysis of this GTP molecule produces GDP and inorganic phosphate (P_i), and it causes the protein to convert to a different, usually inactive, conformation (switch “off”). As shown here, resetting the switch requires the tightly bound GDP to dissociate, a slow step that is greatly accelerated by specific signals; once the GDP has dissociated, a molecule of GTP is quickly rebound.




Figure 3-71 The structure of the Ras protein in its GTP-bound form. This monomeric GTPase illustrates the structure of a GTP-binding domain, which is present in a large family of GTP-binding proteins. The red regions change their conformation when the GTP molecule is hydrolyzed to GDP and inorganic phosphate by the protein; the GDP remains bound to the protein, while the inorganic phosphate is released. The special role of the "switch helix" in proteins related to Ras is explained next (see Figure 3-74).

hydrolyzed, allowing the tRNA to dissociate. Since the GTP hydrolysis is triggered by a proper fit of the tRNA to the mRNA molecule on the ribosome, the EF-Tu serves as an assembly factor that discriminates between correct and incorrect mRNA-tRNA pairings (see Figure 6-66 for a further discussion of this function of EF-Tu).

Comparison of the three-dimensional structure of EF-Tu in its GTP-bound and GDP-bound forms reveals how the unmasking of the tRNA occurs. The dissociation of the inorganic phosphate group (P_i), which follows the reaction $GTP \rightarrow GDP + P_i$, causes a shift of a few tenths of a nanometer at the GTP-binding site, just as it does in the Ras protein. This tiny movement, equivalent to a few times the diameter of a hydrogen atom, causes a conformational change to propagate along a crucial piece of α helix, called the *switch helix*, in the Ras-like domain of the protein. The switch helix seems to serve as a latch that adheres to a specific site in another domain of the molecule, holding the protein in a "shut" conformation. The conformational change triggered by GTP hydrolysis causes the switch helix to detach, allowing separate domains of the protein to swing apart,





Figure 3-72 A comparison of the two major intracellular signaling mechanisms in eucaryotic cells. In both cases, a signaling protein is activated by the addition of a phosphate group and inactivated by the removal of this phosphate. To emphasize the similarities in the two pathways, ATP and GTP are drawn as APPP and GPPP, and ADP and GDP as APP and GPP, respectively. As shown in Figure 3-63, the addition of a phosphate to a protein can also be inhibitory.

Figure 3-73 An aminoacyl tRNA molecule bound to EF-Tu. The three domains of the EF-Tu protein are colored differently, to match Figure 3-74. This is a bacterial protein; however, a very similar protein exists in eukaryotes, where it is called EF-1. (Coordinates determined by P. Nissen et al., *Science* 270:1464-1472, 1995.)



through a distance of about 4 nm. This releases the bound tRNA molecule, allowing its attached amino acid to be used (Figure 3-74).

One can see from this example how cells have exploited a simple chemical change that occurs on the surface of a small protein domain to create a movement 50 times larger. Dramatic shape changes of this type also underlie the very large movements that occur in motor proteins, as we discuss next.

Motor Proteins Produce Large Movements in Cells

We have seen how conformational changes in proteins have a central role in enzyme regulation and cell signaling. We now discuss proteins whose major function is to move other molecules. These **motor proteins** generate the forces responsible for muscle contraction and the crawling and swimming of cells. Motor proteins also power smaller-scale intracellular movements: they help to move chromosomes to opposite ends of the cell during mitosis (discussed in Chapter 18), to move organelles along molecular tracks within the cell (discussed in Chapter 16), and to move enzymes along a DNA strand during the synthesis of a new DNA molecule (discussed in Chapter 5). All these fundamental processes depend on proteins with moving parts that operate as force-generating machines.




Figure 3-74 The large conformational change in EF-Tu caused by GTP hydrolysis. (A) The three-dimensional structure of EF-Tu with GTP bound. The domain at the top is homologous to the Ras protein, and its red α helix is the switch helix, which moves after GTP hydrolysis, as shown in Figure 3-71. (B) The change in the conformation of the switch helix in domain 1 causes domains 2 and 3 to rotate as a single unit by about 90° toward the viewer, which releases the tRNA that was shown bound to this structure in Figure 3-73. (A, adapted from H. Berchtold et al., *Nature* 365:126-132, 1993; B, courtesy of Mathias Sprinzl and Rolf Hilgenfeld.)

How do these machines work? In other words, how are shape changes in proteins used to generate directed movements in cells? If, for example, a protein is required to walk along a narrow thread such as a DNA molecule, it can do this by undergoing a series of conformational changes, such as those shown in Figure 3–75. With nothing to drive these changes in an orderly sequence, however, they are perfectly reversible, and the protein can wander randomly back and forth along the thread. We can look at this situation in another way. Since the directional movement of a protein does work, the laws of thermodynamics (discussed in Chapter 2) demand that such movement utilize free energy from some other source (otherwise the protein could be used to make a perpetual motion machine). Therefore, without an input of energy, the protein molecule can only wander aimlessly.

How, then, can one make the series of conformational changes unidirectional? To force the entire cycle to proceed in one direction, it is enough to make any one of the changes in shape irreversible. For most proteins that are able to walk in one direction for long distances, this is achieved by coupling one of the conformational changes to the hydrolysis of an ATP molecule bound to the protein. The mechanism is similar to the one just discussed that drives allosteric protein shape changes by GTP hydrolysis. Because a great deal of free energy is released when ATP (or GTP) is hydrolyzed, it is very unlikely that the nucleotide-binding protein will undergo the reverse shape change needed for moving backward—since this would require that it also reverse the ATP hydrolysis by adding a phosphate molecule to ADP to form ATP.

In the model shown in Figure 3–76, ATP binding shifts a motor protein from conformation 1 to conformation 2. The bound ATP is then hydrolyzed to produce ADP and inorganic phosphate (P_i), causing a change from conformation 2 to conformation 3. Finally, the release of the bound ADP and P_i drives the protein back to conformation 1. Because the transition 2 → 3 is driven by the energy provided by ATP hydrolysis, this series of conformational changes is effectively irreversible. Thus, the entire cycle goes in only one direction, causing the protein molecule to walk continuously to the right in this example.

Many motor proteins generate directional movement in this general way, including the muscle motor protein *myosin*, which walks along actin filaments to generate muscle contraction, and the *kinesin* proteins that walk along microtubules (both discussed in Chapter 16). These movements can be rapid: some of the motor proteins involved in DNA replication (the DNA helicases) propel themselves along a DNA strand at rates as high as 1000 nucleotides per second.

Membrane-bound Transporters Harness Energy to Pump Molecules Through Membranes

We have thus far seen how allosteric proteins can act as microchips (Cdk and Src kinases), as assembly factors (EF-Tu), and as generators of mechanical force and motion (motor proteins). Allosteric proteins can also harness energy derived from ATP hydrolysis, ion gradients, or electron transport processes to pump specific ions or small molecules across a membrane. We consider one example here; others will be discussed in Chapter 11.

One of the best understood pump proteins is the calcium transport protein from muscle cells. This protein, called the **Ca²⁺ pump**, is embedded in the membrane of a specialized organelle in muscle cells called the *sarcoplasmic reticulum*. The Ca²⁺ pump (also known as the Ca²⁺ ATPase) maintains the low cytoplasmic calcium concentration of resting muscle by pumping calcium out of the cytosol into the membrane-enclosed sarcoplasmic reticulum; then, in response to a nerve impulse, Ca²⁺ is rapidly released (through other channels) back into the cytosol to trigger muscle contraction. The Ca²⁺ pump is homologous to the




Figure 3–75 An allosteric “walking” protein. Although its three different conformations allow it to wander randomly back and forth while bound to a thread or a filament, the protein cannot move uniformly in a single direction.





Figure 3–76 An allosteric motor protein. The transition between three different conformations includes a step driven by the hydrolysis of a bound ATP molecule, and this makes the entire cycle essentially irreversible. By repeated cycles, the protein therefore moves continuously to the right along the thread.



Na^+ - K^+ pump that maintains Na^+ and K^+ concentration differences across the plasma membrane, both being members of a family of P-type cation pumps—so named because these proteins are autophosphorylated during their reaction cycle. A large, cytoplasmic head region binds and hydrolyzes ATP, forming a covalent bond with the phosphate released. The protein thereby transiently shifts to a high-energy, phosphorylated state that is tightly bound to two Ca^{2+} ions that were picked up from the cytosol. This form of the protein then decays to a low-energy, phosphorylated state, which causes the Ca^{2+} ions to be released into the lumen. Dephosphorylation of the enzyme finally releases the phosphate and resets the protein for its next round of ion pumping.

The head region of all P-type cation pumps is linked to a series of ten transmembrane α helices, four of which form transmembrane Ca^{2+} -binding sites for the Ca^{2+} pump. The three-dimensional structure of this protein has been deciphered by high-resolution electron microscopy and x-ray diffraction (see Figure 11–15). This has enabled biologists to derive a molecular model for pump action based on extensive biochemical data on its normal and mutant forms.

As illustrated in Figure 3–77A, the transmembrane α helices that bind the Ca^{2+} wind around each other and create a cavity for Ca^{2+} between the helices. ATP hydrolysis generates a series of major conformational changes in the cytoplasmic head, which—through its stalk connection to the transmembrane helices—changes the structure and relative orientations of some of the helices in the membrane, thereby altering the cavity in a way that pushes the Ca^{2+} ions unidirectionally across the membrane (Figure 3–77B). As for a motor protein, unidirectional transport of an ion requires the cycle to use energy, so as to impart a preferred directionality to the protein's conformational changes.

Humans have invented many different types of mechanical pumps, and it should not be surprising that cells also contain membrane-bound pumps that function in other ways. Most notable are the rotary pumps that couple the hydrolysis of ATP to the transport of H^+ ions (protons). These pumps resemble miniature turbines, and they are used to acidify the interior of lysosomes and other eucaryotic organelles. Like other ion pumps that create ion gradients, they can function in reverse to catalyze the reaction $\text{ADP} + \text{P}_i \rightarrow \text{ATP}$ if there is a steep enough gradient across their membrane of the ion that they transport.

One such pump, the ATP synthase, harnesses a gradient of proton concentration produced by electron transport processes to produce most of the ATP used in the living world. Because this ubiquitous pump has such a central role in energy conversion, we postpone discussing its three-dimensional structure and mechanism until Chapter 14.

Figure 3–77 The transport of calcium ions by the Ca^{2+} pump. (A) The structure of the pump protein, formed from a single subunit of 994 amino acids (see Figure 11–15 for details). (B) A model for ion pumping. For simplicity, only two of the four transmembrane α helices that bind Ca^{2+} are shown.

Proteins Often Form Large Complexes That Function as Protein Machines

As one progresses from small, single-domain proteins to large proteins formed from many domains, the functions the proteins can perform become more elaborate. The most impressive tasks, however, are carried out by large protein assemblies formed from many protein molecules. Now that it is possible to reconstruct most biological processes in cell-free systems in the laboratory, it is clear that each of the central processes in a cell—such as DNA replication, protein synthesis, vesicle budding, or transmembrane signaling—is catalyzed by a highly coordinated, linked set of ten or more proteins. In most such protein machines, the hydrolysis of bound nucleoside triphosphates (ATP or GTP) drives an ordered series of conformational changes in some of the individual protein subunits, enabling the ensemble of proteins to move coordinately. In this way, each of the appropriate enzymes can be placed directly into the positions where they are needed to perform successive reactions in a series. This is what occurs, for example, in protein synthesis on a ribosome (discussed in Chapter 6)—or in DNA replication, where a large multiprotein complex moves rapidly along the DNA (discussed in Chapter 5).

Cells have evolved protein machines for the same reason that humans have invented mechanical and electronic machines. For accomplishing almost any task, manipulations that are spatially and temporally coordinated through linked processes are much more efficient than the sequential use of individual tools.

A Complex Network of Protein Interactions Underlies Cell Function

There are many challenges facing cell biologists in this “post-genome” era when complete genome sequences are known. One is the need to dissect and reconstruct each one of the thousands of protein machines that exist in an organism such as ourselves. To understand these remarkable protein complexes, each must be reconstituted from its purified protein parts—so that its detailed mode of operation can be studied under controlled conditions in a test tube, free from all other cell components. This alone is a massive task. But we now know that each of these subcomponents of a cell also interacts with other sets of macromolecules, creating a large network of protein–protein and protein–nucleic acid interactions throughout the cell. To understand the cell, therefore, one will need to analyze most of these other interactions as well.

Some idea of the complexity of intracellular protein networks can be gained from a particularly well-studied example described in Chapter 16: the many dozens of proteins that interact with the actin cytoskeleton in the yeast *Saccharomyces cerevisiae* (see Figure 16–15).

The extent of such protein–protein interactions can also be estimated more generally. In particular, large-scale efforts have been undertaken to detect these interactions using the two-hybrid screening method described in Chapter 8 (see Figure 8–51). Thus, for example, this method is being applied to a large set of the 6000 gene products produced by *S. cerevisiae*. As expected, the majority of the interactions that have been observed are between proteins in the same functional group. That is, proteins involved in cell-cycle control tend to interact extensively with each other, as do proteins involved with DNA synthesis, or DNA repair, and so on. But there are also a surprisingly large number of interactions between the protein members of different functional groups (Figure 3–78). These interactions are presumably important for coordinating cell functions, but most of them are not understood.

An examination of the range of available data suggests that an average protein in a human cell may interact with somewhere between 5 and 15 different partners. Often, a different set of partners will be bound by each of the different domains in a multidomain protein; in fact, one can speculate that the unusually extensive multidomain structures observed for human proteins (see p. 462) may have evolved to generate these interactions. Given the enormous complexity of