

1995

USP 23

NF 18

THE UNITED STATES PHARMACOPEIA

THE NATIONAL FORMULARY

By authority of the United States Pharmacopeial Convention, Inc., meeting at Washington, D.C., March 8-10, 1990. Prepared by the Committee of Revision and published by the Board of Trustees

Official from January 1, 1995



UNITED STATES PHARMACOPEIAL CONVENTION, INC.
12601 Twinbrook Parkway, Rockville, MD 20852

... sodium chloride inhalation solution may be nebulized using nebulizer gases. Nebulizers are suitable for the administration of solution solutions only if they give droplets sufficiently fine and uniform in size so that the mist reaches the bronchioles. Nebulized solutions may be breathed directly from the nebulizer or the nebulizer may be attached to a plastic face mask, tent, or intermittent positive pressure breathing (IPPB) machine.

Another group of products, also known as metered-dose inhalers (MDIs) are propellant driven drug suspensions or solutions of a liquid gas propellant with or without a cosolvent and are designed for delivering metered doses of the drug to the respiratory tract. An MDI contains multiple doses, often exceeding several hundred. The most common single-dose volumes delivered are from 25 to 100 μ L (also expressed as mg) per actuation. Examples of MDIs containing drug solutions and suspensions in this pharmacopeia are *Epinephrine Inhalation Aerosol* and *Salbutamol Hydrochloride and Phenylephrine Bitartrate Inhalation Aerosol*, respectively.

Powders may also be administered by mechanical devices that generate manually produced pressure or a deep inhalation by the patient (e.g., *Cromolyn Sodium for Inhalation*).

A special class of inhalations termed inhalants consists of drugs or combination of drugs, that by virtue of their high vapor pressure can be carried by an air current into the nasal passage where they exert their effect. The container from which the inhalant is normally administered is known as an inhaler.

INJECTIONS

See *Injections* (1).

IRRIGATIONS

Irrigations are sterile solutions intended to bathe or flush open wounds or body cavities. They are used topically, never parenterally. They are labeled to indicate that they are not intended for injection.

LOTIONS

See *Solutions or Suspensions*.

LOZENGES

Lozenges are solid preparations, which are intended to dissolve and integrate slowly in the mouth. They contain one or more ingredients, usually in a flavored, sweetened base. They can be prepared by molding (gelatin and/or fused sucrose or sorbitol) or by compression of sugar based tablets. Molded lozenges are sometimes referred to as pastilles while compressed lozenges are often referred to as troches. They are usually intended for relief of local irritation or infections of the mouth or throat and may contain active ingredients intended for systemic absorption after swallowing.

OINTMENTS

Ointments are semisolid preparations intended for external application to the skin or mucous membranes.

Ointment bases recognized for use as vehicles fall into four general classes: the hydrocarbon bases, the absorption bases, the water-removable bases, and the water-soluble bases. Each therapeutic ointment possesses as its base a representative of one of these four general classes.

Hydrocarbon Bases

These bases, which are known also as "oleaginous ointment bases" are represented by *White Petrolatum* and *White Ointment*. Only small amounts of an aqueous component can be incorporated into them. They serve to keep medicaments in prolonged contact with the skin and act as occlusive dressings. Hydrocarbon bases are used chiefly for their emollient effects, and are difficult to wash off. They do not "dry out" or change appreciably on aging.

This class of bases may be divided into two groups: the first group consisting of bases that permit the incorporation of aqueous solutions with the formation of a water-in-oil emulsion (*Hydrophilic Petrolatum* and *Lanolin*), and the second group consisting of water-in-oil emulsions that permit the incorporation of additional quantities of aqueous solutions (*Lanolin*). Absorption bases are useful also as emollients.

Water-removable Bases

Such bases are oil-in-water emulsions, e.g., *Hydrophilic Ointment*, and are more correctly called "creams." (See *Creams*.) They are also described as "water-washable," since they may be readily washed from the skin or clothing with water, an attribute that makes them more acceptable for cosmetic reasons. Some medicaments may be more effective in these bases than in hydrocarbon bases. Other advantages of the water-removable bases are that they may be diluted with water and that they favor the absorption of serous discharges in dermatological conditions.

Water-soluble Bases

This group of so-called "greaseless ointment bases" is comprised of water-soluble constituents. *Polyethylene Glycol Ointment* is the only Pharmacopeial preparation in this group. Bases of this type offer many of the advantages of the water-removable bases and, in addition, contain no water-insoluble substances such as petrolatum, anhydrous lanolin, or waxes. They are more correctly called "Gels." (See *Gels*.)

Choice of Base—The choice of an ointment base depends upon many factors, such as the action desired, the nature of the medication to be incorporated and its bioavailability and stability, and the requisite shelf-life of the finished product. In some cases, it is necessary to use a base that is less than ideal in order to achieve the stability required. Drugs that hydrolyze rapidly, for example, are more stable in hydrocarbon bases than in bases containing water, even though they may be more effective in the latter.

OPHTHALMIC PREPARATIONS

Drugs are administered to the eyes in a wide variety of dosage forms, some of which require special consideration. They are discussed in the following paragraphs.

Ointments

Ophthalmic ointments are ointments for application to the eye. Special precautions must be taken in the preparation of ophthalmic ointments. They are manufactured from sterilized ingredients under rigidly aseptic conditions and meet the requirements under *Sterility Tests* (71). If the specific ingredients used in the formulation do not lend themselves to routine sterilization techniques, ingredients that meet the sterility requirements described under *Sterility Tests* (71), along with aseptic manufacture, may be employed. Ophthalmic ointments must contain a suitable substance or mixture of substances to prevent growth of, or to destroy, microorganisms accidentally introduced when the container is opened during use, unless otherwise directed in the individual monograph, or unless the formula itself is bacteriostatic (see *Added Substances* under *Ophthalmic Ointments* (771)). The medicinal agent is added to the ointment base either as a solution or as a micronized powder. The finished ointment must be free from large particles and must meet the requirements for *Leakage* and for *Metal Particles* under *Ophthalmic Ointments* (771). The immediate containers for ophthalmic ointments shall be sterile at the time of filling and closing. It is mandatory that the immediate containers for ophthalmic ointments be sealed and tamper-proof so that sterility is assured at time of first use.

The ointment base that is selected must be nonirritating to the eye, permit diffusion of the drug throughout the secretions bathing the eye, and retain the activity of the medicament for a reasonable period under proper storage conditions.

Petrolatum is mainly used as a base for ophthalmic drugs. Some absorption bases, water-removable bases, and water-soluble

bases may be desirable for water-soluble drugs. Such bases allow for better dispersion of water-soluble medicaments, but they must be nonirritating to the eye.

Solutions

Ophthalmic solutions are sterile solutions, essentially free from foreign particles, suitably compounded and packaged for instillation into the eye. Preparation of an ophthalmic solution requires careful consideration of such factors as the inherent toxicity of the drug itself, isotonicity value, the need for buffering agents, the need for a preservative (and, if needed, its selection), sterilization, and proper packaging. Similar considerations are also made for nasal and otic products.

ISOTONICITY VALUE

Lacrimal fluid is isotonic with blood, having an isotonicity value corresponding to that of a 0.9% sodium chloride solution. Ideally, an ophthalmic solution should have this isotonicity value; but the eye can tolerate isotonicity values as low as that of a 0.6% sodium chloride solution and as high as that of a 2.0% sodium chloride solution without marked discomfort.

Some ophthalmic solutions are necessarily hypertonic in order to enhance absorption and provide a concentration of the active ingredient(s) strong enough to exert a prompt and effective action. Where the amount of such solutions used is small, dilution with lacrimal fluid takes place rapidly so that discomfort from the hypertonicity is only temporary. However, any adjustment toward isotonicity by dilution with tears is negligible where large volumes of hypertonic solutions are used as collyria to wash the eyes; it is therefore important that solutions used for this purpose be approximately isotonic.

BUFFERING

Many drugs, notably alkaloidal salts, are most effective at pH levels that favor the undissociated free bases. At such pH levels, however, the drug may be unstable so that compromise levels must be found and held by means of buffers. One purpose of buffering some ophthalmic solutions is to prevent an increase in pH caused by the slow release of hydroxyl ions by glass. Such a rise in pH can affect both the solubility and the stability of the drug. The decision whether or not buffering agents should be added in preparing an ophthalmic solution must be based on several considerations. Normal tears have a pH of about 7.4 and possess some buffer capacity. The application of a solution to the eye stimulates the flow of tears and the rapid neutralization of any excess hydrogen or hydroxyl ions within the buffer capacity of the tears. Many ophthalmic drugs, such as alkaloidal salts, are weakly acidic and have only weak buffer capacity. Where only 1 or 2 drops of a solution containing them are added to the eye, the buffering action of the tears is usually adequate to raise the pH and prevent marked discomfort. In some cases pH may vary between 3.5 and 8.5. Some drugs, notably pilocarpine hydrochloride and epinephrine bitartrate, are more acid and overtax the buffer capacity of the lacrimal fluid. Ideally, an ophthalmic solution should have the same pH, as well as the same isotonicity value, as lacrimal fluid. This is not usually possible since, at pH 7.4, many drugs are not appreciably soluble in water. Most alkaloidal salts precipitate as the free alkaloid at this pH. Additionally, many drugs are chemically unstable at pH levels approaching 7.4. This instability is more marked at the high temperatures employed in heat sterilization. For this reason, the buffer system should be selected that is nearest to the physiological pH of 7.4 and does not cause precipitation of the drug or its rapid deterioration.

An ophthalmic preparation with a buffer system approaching the physiological pH can be obtained by mixing a sterile solution of the drug with a sterile buffer solution using aseptic technique. Even so, the possibility of a shorter shelf-life at the higher pH must be taken into consideration, and attention must be directed toward the attainment and maintenance of sterility throughout the manipulations.

Many drugs, when buffered to a therapeutically acceptable pH, would not be stable in solution for long periods of time. These products are lyophilized and are intended for reconstitution immediately before use (e.g., *Acetylcholine Chloride for Ophthalmic Solution*).

STERILIZATION

The sterility of solutions applied to an injured eye is of greatest importance. Sterile preparations in appropriate containers for individual use on one patient should be available in hospital, office, or other installation where accidentally or surgically traumatized eyes are treated. The method of sterilization is determined primarily by the character of the product (see *Sterilization and Sterility Assurance of Pharmaceutical Articles* (1211)).

Whenever possible, sterile membrane filtration under appropriate conditions is the preferred method. If it can be shown that product stability is not adversely affected, sterilization by heat in the final container is also a preferred method.

Buffering certain drugs near the physiological pH makes them quite unstable at high temperature.

Avoiding the use of heat by employing a bacterial filter is a valuable technique, provided caution is exercised in selection, assembly, and use of the equipment. Single-dose, presterilized disposable units are available and should be used wherever possible.

PRESERVATION

Ophthalmic solutions may be packaged in multiple-use containers when intended for the individual use of one patient where the ocular surfaces are intact. It is mandatory that multiple-use containers for ophthalmic solutions be tamper-proof so that sterility is assured at time of use. Multiple-use solutions must contain a suitable substance or substances to prevent the growth of, or to destroy, microorganisms accidentally introduced when the container is opened.

Where intended for use in surgical procedures, ophthalmic solutions, although they must be sterile, should not contain antibacterial agents, since they may be irritating to the eye.

THICKENING AGENT

A pharmaceutical grade of methylcellulose (0.4% solution, viscosity is 25 centipoises, or 0.25% if 4000 centipoises) and suitable thickening agents such as hydroxypropyl methylcellulose or polyvinyl alcohol occasionally are added to ophthalmic solutions to increase the viscosity and prolong contact of the solution with the tissue. The thickened ophthalmic solution is free from visible particles.

Suspensions

Ophthalmic suspensions are sterile liquid preparations containing solid particles dispersed in a liquid vehicle for application to the eye (see *Suspensions*). In multiple-use containers, such suspensions contain the drug in a nonirritating vehicle to prevent irritation and/or scratching of the cornea. Multiple-use suspensions should never be dispensed if there is evidence of aggregation.

Strips

Fluorescein sodium solution should be dispensed in a single-use container or in the form of a sterile, prepackaged strip. The strip releases a sufficient amount of fluorescein for diagnostic purposes when touched to the eye to detect a foreign body or a corneal abrasion. Contact with the eye may be avoided by leaching the strip with sterile water before the eye with the aid of sterile water or saline solution.

PASTES

Pastes are semisolid dosage forms that contain drug substances intended for topical application. They are made from a single phase aqueous phase (e.g., *Fluorescein Sodium Paste*). The other phase (e.g., *Zinc Oxide Paste*), consists of thick, fatty bases that ordinarily flow at body temperature, and they form protective coatings over the areas to which they are applied.

The fatty pastes appear less greasy and less irritating than ointments by reason of a high proportion of water-soluble components having an affinity for water. These pastes are used for the treatment of