

Figure 1. High-pulse repetition frequency Doppler recordings in a patient after valve replacement for aortic stenosis. Left, At baseline, a normal symmetric Doppler spectrum reaching 1 m/s was observed. Middle, During nipride, the afterload reduction was associated with a dagger-shaped Doppler spectrum peaking in late systole at 3 m/s. Right, Inotropic stimulation by dobutamine resulted in a similar intracavitary flow velocity pattern with a maximal velocity of 3.8 m/s.

#### Statistical Analysis <sup>†</sup>

Results are expressed as mean plus/minus SD. Unpaired and paired t tests were used to compare appropriate data. Fisher's exact test was performed for comparison of categorical variables. One-way ANOVA, followed by Bonferroni's test, was performed for multiple comparisons. Sensitivity and specificity were calculated as usual. Multivariate analysis was performed separately on Doppler echocardiographic and catheterization data to detect their values in predicting postoperative AFV. Optimal diagnostic accuracy level for a given factor was defined as the value of the given factor at which sensitivity and specificity are equal. A value of P less or equal to .05 was considered significant.

#### Results <sup>†</sup>

All patients underwent both preoperative and postoperative Doppler echocardiographic studies. The latter was performed 7 plus/minus 3 days (range, 2 to 10 days) after valve replacement. After surgery, nipride was given in 93 patients and dobutamine in 96 patients. Three patients did not receive nipride because of persistent hypotension (< 100 mm Hg). Four patients received neither nipride nor dobutamine because AFV was present at rest (> 3.7 m/s).

#### Hemodynamics <sup>†</sup>

In the whole population, nipride induced a decrease in systolic pressure and an increase in heart rate (Table 1). Dobutamine infusion increased heart rate, whereas systolic blood pressure remained unchanged. Patients with resting AFV had significantly higher heart rates and lower systolic pressures at baseline and during dobutamine infusion compared with pati

resting AFV. A comparison of patients according to the pres

NORRED EXHIBIT 2012 - Page 1  
Medtronic, Inc., Medtronic Vascular, Inc.,  
& Medtronic Corevalve, LLC  
v. Troy R. Norred, M.D.  
Case IPR2014-00110




Figure 1. High-pulse repetition frequency Doppler recordings in a patient after valve replacement for aortic stenosis. Left, At baseline, a normal symmetric Doppler spectrum reaching 1 m/s was observed. Middle, During nipride, the afterload reduction was associated with a dagger-shaped Doppler spectrum peaking in late systole at 3 m/s. Right, Inotropic stimulation by dobutamine resulted in a similar intracavitary flow velocity pattern with a maximal velocity of 3.8 m/s.

#### Statistical Analysis <sup>†</sup>

Results are expressed as mean plus/minus SD. Unpaired and paired *t* tests were used to compare appropriate data. Fisher's exact test was performed for comparison of categorical variables. One-way ANOVA, followed by Bonferroni's test, was performed for multiple comparisons. Sensitivity and specificity were calculated as usual. Multivariate analysis was performed separately on Doppler echocardiographic and catheterization data to detect their values in predicting postoperative AFV. Optimal diagnostic accuracy level for a given factor was defined as the value of the given factor at which sensitivity and specificity are equal. A value of *P* less or equal to .05 was considered significant.

#### Results <sup>†</sup>

All patients underwent both preoperative and postoperative Doppler echocardiographic studies. The latter was performed 7 plus/minus 3 days (range, 2 to 10 days) after valve replacement. After surgery, nipride was given in 93 patients and dobutamine in 96 patients. Three patients did not receive nipride because of persistent hypotension (< 100 mm Hg). Four patients received neither nipride nor dobutamine because AFV was present at rest (> 3.7 m/s).

#### Hemodynamics <sup>†</sup>

In the whole population, nipride induced a decrease in systolic pressure and an increase in heart rate (Table 1). Dobutamine infusion increased heart rate, whereas systolic blood pressure remained unchanged. Patients with resting AFV had significantly higher heart rates and lower systolic pressures at baseline and during dobutamine infusion compared with patients without resting AFV. A comparison of patients according to the presence or absence of preoperative AFV showed them to be similar with respect to heart rate, blood

① Mike Stwak  
882-2640

② Debbie Ikelburgen

③ Delivery System

Jennifer  
Robins

18 yo w/ mva  
Quad / vent dependent

