

# REACTION MECHANISMS IN ENVIRONMENTAL ORGANIC CHEMISTRY

Richard A. Larson  
Eric J. Weber



LEWIS PUBLISHERS  
Boca Raton Ann Arbor London Tokyo

**DOCKET**  
**A L A R M**

Find authenticated court documents without watermarks at [docketalarm.com](http://docketalarm.com).

Library of Congress Cataloging-in-Publication Data

Larson, Richard A.

Reaction mechanisms in environmental organic chemistry / Richard A. Larson and Eric J. Weber  
p. cm.

Includes bibliographical references and index.

1. Organic compounds—Environmental aspects. 2. Environmental chemistry. 3. Chemical reactions. I. Weber, Eric J. II. Title.

TD196.073L37 1994

628.5—dc20 93-1622

ISBN 0-87371-258-7

This book contains information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. A wide variety of references are listed. Reasonable efforts have been made to publish reliable data and information, but the authors and the publisher cannot assume responsibility for the validity of all materials or for the consequences of their use.

Neither this book nor any part may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, microfilming, and recording, or by any information storage or retrieval system, without prior permission in writing from the publisher.

CRC Press, Inc.'s consent does not extend to copying for general distribution, for promotion, for creating new works, or for resale. Specific permission must be obtained in writing from CRC Press for such copying.

Direct all inquiries to CRC Press, Inc., 2000 Corporate Blvd., N.W., Boca Raton, Florida 33431.

© 1994 by CRC Press, Inc.

Lewis Publishers is an imprint of CRC Press

No claim to original U.S. Government works

International Standard Book Number 0-87371-258-7

Library of Congress Card Number 93-1622

Printed in the United States of America

1 2 3 4 5 6 7 8 9 0

Printed on acid-free paper



NOVARTIS EXHIBIT 2034  
Noven v. Novartis and LTS Lohmann  
IPR2014-00550  
Page 2 of 35

## CHAPTER 2


---

# HYDROLYSIS

---

### A. INTRODUCTION

In general terms, hydrolysis is defined as a chemical transformation in which an organic molecule, RX, reacts with water, resulting in the formation of a new covalent bond with OH and cleavage of the covalent bond with X (the leaving group) in the original molecule. The net reaction is the displacement of X by OH<sup>-</sup> (Harris, 1981; Mill and Mabey, 1988)




Hydrolysis is an example of a larger class of reactions referred to as nucleophilic displacement reactions in which a nucleophile (an electron-rich species containing an unshared pair of electrons) attacks an electrophilic atom (an electron-deficient reaction center). Hydrolytic processes encompass several types of reaction mechanisms that can be defined by the type of reaction center (i.e., the atom bearing the leaving group, X) where hydrolysis occurs. The reaction mechanisms encountered most often are direct and indirect nucleophilic substitution and nucleophilic addition-elimination.

The chemical structures of hydrolyzable functional groups that are prevalent in environmental chemicals, and the reaction products resulting from their hydrolysis, are listed in Table 2.1. For these classes of chemicals, hydrolysis may be the dominant pathway for their transformation in aquatic ecosystems. Hydrolytic processes are not limited to the bodies of water such as rivers, streams, lakes, and oceans usually associated with the term aquatic ecosystems. Hydrolysis of organic chemi-


Table 2.1. Examples of Hydrolyzable Functional Groups

## 1. Halogenated Aliphatics


Nucleophilic substitution




Elimination




## 2. Epoxides




## 3. Organophosphorus Esters




## 4. Carboxylic Acid Esters




## 5. Anhydrides




## 6. Amides



## 7. Carbamates



## 8. Ureas



als also can occur in fogwater, biological systems, groundwater systems, and the aqueous microenvironment associated with soils and sediments.

## B. HYDROLYSIS KINETICS

## 1. Specific Acid and Base Catalysis

In addition to nucleophilic attack by  $\text{H}_2\text{O}$  (neutral hydrolysis), we find that hydrolytic reactions are sensitive to specific acid and specific base catalysis (i.e., catalysis by hydronium ion,  $\text{H}^+$ , and hydroxide ion,  $\text{OH}^-$ , respectively). Accordingly, hydrolysis kinetics must take into account the potential for  $\text{H}_2\text{O}$  to dissociate. Even at pH 7.0, where the concentration of  $\text{H}^+$  and  $\text{OH}^-$  is only  $10^{-7}$  M, specific acid and specific base catalysis can significantly accelerate hydrolysis kinetics. Specific acid and base catalysis occurs because the hydronium ion and hydroxide ions provide an alternative mechanism for hydrolysis that is energetically more favorable. In specific acid catalysis, hydronium ion is thought to provide a reaction pathway of lower energy by withdrawing electron density from the atom bearing the leaving group, X, thus making it more susceptible to nucleophilic attack by  $\text{H}_2\text{O}$ . Specific base catalysis occurs because  $\text{OH}^-$  is a much more reactive nucleophile than  $\text{H}_2\text{O}$  (typically by a factor of about  $10^4$ ; Streitwieser, 1962). Accordingly, a hydrolysis reaction involving nucleophilic attack by  $\text{OH}^-$  will occur at a faster rate than the pathway involving nucleophilic attack by  $\text{H}_2\text{O}$  alone.

Taking into consideration specific acid and base catalysis (the concentration of a catalyst is reflected in the rate law but is not reflected in the equilibrium constant), the hydrolysis rate term for hydrolyzable chemicals can be described by:

$$d[\text{RX}]/dt = k_{\text{hyd}}[\text{RX}] = k_{\text{a}}[\text{H}^+][\text{RX}] + k_{\text{n}}[\text{RX}] + k_{\text{b}}[\text{OH}^-][\text{RX}] \quad (2.2)$$

where  $[\text{RX}]$  is the concentration of the hydrolyzable compound,  $k_{\text{hyd}}$  is the observed or measured hydrolysis rate constant, and  $k_{\text{a}}$ ,  $k_{\text{n}}$ , and  $k_{\text{b}}$  are the rate constants for the acid-catalyzed, neutral, and base-catalyzed processes. Assuming that the individual rate processes for the acid, neutral, and base hydrolyses obey first-order kinetics with respect to the hydrolyzable chemical, RX, it is possible to write the following equation for  $k_{\text{hyd}}$ :

$$k_{\text{hyd}} = k_{\text{a}}[\text{H}^+] + k_{\text{n}} + k_{\text{b}}[\text{OH}^-] \quad (2.3)$$

From the equilibrium term for the ionization of water,  $K_{\text{w}}$ :

$$K_{\text{w}} = [\text{OH}^-][\text{H}^+] = 1 \times 10^{-14} \quad (2.4)$$

it is possible to substitute for  $[\text{OH}^-]$  in Equation 2.4, giving:

$$k_{\text{hyd}} = k_a[\text{H}^+] + k_n + k_b(K_w/[\text{H}^+]) \quad (2.5)$$

Because  $k_{\text{hyd}}$  is a pseudo first-order rate constant at a fixed pH (i.e., the hydrolysis is independent of RX concentration), the half-life for hydrolysis can be calculated from Equation 2.6.

$$t_{1/2} = \ln 2/k_{\text{hyd}} \quad (2.6)$$

## 2. pH Dependence

It is apparent from Equation 2.5 that the overall rate constant for hydrolysis,  $k_{\text{hyd}}$ , depends on pH and the magnitude of the rate constants for the individual processes. Plots of  $\log k_{\text{hyd}}$  versus pH are very useful for determining the contribution of the acid, neutral, and base terms for hydrolysis of a compound of interest at a specific pH. Figure 2.1 illustrates the  $\log k_{\text{hyd}}$  versus pH plot for several chemicals of interest. These data demonstrate that the relationship between hydrolysis kinetics and pH is dependent on the nature of the hydrolyzable functional group. For example, over typical environmental pH values (4 to 8), the neutral hydrolysis rate term for ethylene oxide and methyl chloride will dominate. Only below pH 4 will the acid-catalyzed rate term for ethylene oxide contribute to the overall hydrolysis rate term. Likewise, base-catalyzed hydrolysis of ethylene oxide and methyl chloride will not






Figure 2.1. pH-rate profiles for the hydrolysis of ethylene oxide, methyl chloride and ethyl acetate (Data taken from Mabey and Mill, 1978).

compete with neutral hydrolysis until pH 11, which is well above the pH range of natural aquatic ecosystems. By contrast, the neutral hydrolysis of ethyl acetate occurs over a much narrower pH range (6 to 7). Either below or above this pH range, the rate terms for acid- or base-catalyzed hydrolysis will dominate.

## C. HYDROLYSIS REACTION MECHANISMS

Reaction mechanisms for hydrolysis can be classified according to the type of reaction center involved. The primary distinction is made between reaction at saturated and unsaturated centers. With respect to carbon-centered functional groups, which will be the primary focus of this chapter, hydrolysis involves reactions at  $\text{sp}^3$  (saturated) or  $\text{sp}^2$  (unsaturated) hybridized carbons. Nucleophilic reactions at  $\text{sp}^3$  carbons are termed nucleophilic substitution (2.7). The reaction at  $\text{sp}^2$  carbons is termed nucleophilic addition-elimination or acyl substitution (2.8).



Because both the nucleophile,  $\text{Y}^-$ , and the leaving group,  $\text{X}^-$ , are Lewis bases, these are examples of Lewis acid-base reactions in which one Lewis base replaces another in the Lewis acid-base adduct (Jensen, 1978).

### 1. Nucleophilic Substitution

The limiting cases of nucleophilic substitution have been described as the ionization mechanism ( $\text{S}_{\text{N}}1$ , substitution-nucleophilic-unimolecular) and the direct displacement mechanism ( $\text{S}_{\text{N}}2$ , substitution-nucleophilic-bimolecular; Gleave et al., 1935). The  $\text{S}_{\text{N}}1$  and  $\text{S}_{\text{N}}2$  mechanisms describe the extremes in nucleophilic substitution reactions. Pure  $\text{S}_{\text{N}}1$  and  $\text{S}_{\text{N}}2$  reaction mechanisms, however, are rarely observed. More often a mix of these reaction mechanisms are occurring simultaneously.

#### $\text{S}_{\text{N}}1$ Mechanism

The  $\text{S}_{\text{N}}1$  mechanism begins by a rate-determining heterolytic dissociation of the substrate to an  $\text{sp}^2$  hybridized carbocation (commonly referred to as a carbonium ion) and the leaving group (2.9):

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.