



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 33/14, 31/66, 31/20	A2	(11) International Publication Number: WO 97/39759 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/US97/06712 (22) International Filing Date: 23 April 1997 (23.04.97) (30) Priority Data: 60/016,140 24 April 1996 (24.04.96) US (71) Applicant (for all designated States except US): BRIGHAM AND WOMEN'S HOSPITAL [US/US]; 75 Francis Street, Boston, MA 02115 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): STOLL, Andrew, L. [US/US]; 35 Old Winter Street, Lincoln, MA 01773 (US). SEVERUS, Wolfram, E. [DE/DE]; Badensche Strasse 7, D-10825 Berlin (DE). (74) Agent: SANZO, Michael, A.; Vinson & Elkins, 1455 Pennsylvania Avenue, N.W., Washington, DC 20004 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: OMEGA-3 FATTY ACIDS AND OMEGA-3 PHOSPHATIDYLCHOLINE IN THE TREATMENT OF BIPOLAR DISORDER		
(57) Abstract <p>The present invention is directed to a method of treating patients with bipolar disorder by administering omega-3 fatty acids. These may be administered in a substantially purified form, as part of a pharmaceutical composition, or as part of a larger molecule, e.g. a triacylglycerol, which releases free fatty acid after ingestion by a patient. The present invention is also directed to triacylglycerols which are esterified at the gamma carbon of glycerol to phosphocholine and at either the alpha or beta carbon of glycerol to an omega-3 fatty acid. These "omega-3 phosphatidylcholines" are also used in the treatment of patients with bipolar disorder.</p>		

Petition for Inter Partes Review
Of U.S. Patent 8,278,351

Exhibit

ENZYMOTEC - 1010

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Omega-3 Fatty Acids and Omega-3 Phosphatidylcholine in the Treatment of Bipolar Disorder

Field of the Invention

5 The present invention relates to medical treatments for psychiatric disorders. More specifically, it is concerned with novel methods and compositions for treating patients with bipolar disorder.

Background of the Invention

10 Patients with bipolar disorder suffer recurrent, alternating cycles of mania and depression. In a controlled clinical study performed more than a decade ago, it was reported that lecithin (phosphatidylcholine) has anti-manic properties when administered to such patients (Cohen *et al.*, *Am. J. Psychiatry* 139:1162-1164 (1982); see also Cohen *et al.*, *Am. J. Psychiatry* 137:242-243 (1980); Schreier, *Am. J. Psychiatry* 139:108-110 (1982)). More recent reports, have suggested that the beneficial effects observed for lecithin are due primarily to the metabolic release of free choline (Stoll *et al.*, *Biol. Psychiatry* 37:170-174 (1995)).

15 Although effective in reducing mania, lecithin is not widely used in treating bipolar patients. One of the main reasons for this is that 15-30 grams of lecithin per day must typically be given to a patient in order to obtain a beneficial effect and the intake of such a large quantity of lipid would, over time, tend to promote cardiovascular disease. An ideal solution to this problem would be to administer a therapeutic agent that has the same beneficial effect as lecithin
20 in controlling mania but which does not have the same adverse effect with respect to cardiovascular disease.

The present invention is directed to phosphatidylcholines in which the α or β carbon of glycerol is esterified to an omega-3 fatty acid. These fatty acids are unique among dietary fats in that they inhibit thrombosis and platelet aggregation and can lower blood pressure (see
25 Dimmitt, *Clin. Exp. Pharmacol. Physiol.* 22:204-208 (1995)). Thus, the "omega-3 phosphatidylcholines" disclosed herein produce the same effects as lecithin in bipolar patients due to the release of free choline but reduce, rather than increase, the risk that a patient will suffer a stroke or coronary thrombosis.

In addition, the present invention is directed to a method of treating bipolar disorder
30 using omega-3 fatty acids themselves, *i.e.* apart from phosphatidylcholine. These may be administered in a purified state, as part of a composition containing other therapeutic agents or

as part of another compound, e.g. a triacylglycerol, which is metabolized to release free fatty acid *in vivo*.

Summary of the Invention

5 An evaluation of mood stabilizing agents indicates that all such agents presently used to treat bipolar patients have an inhibitory effect on neuronal signal transduction systems. The present invention is based, in part, upon this discovery and the upon the recognition that omega-3 fatty acids are useful in treating pathological conditions involving excessive cell signal transduction (see e.g., Sperling, *Rheum. Dis. Clinics* 17 (1991); Sperling, *et al.*, *J. Clin. Invest.* 91:651-660 (1993)).

10 Thus, a method has been developed for treating a human patient for bipolar disorder by administering omega-3 fatty acids at a dosage sufficient to reduce or eliminate the symptoms associated with the disorder, i.e. at a dosage sufficient to reduce the frequency of mood fluctuations or lessen the severity of the mania or depression experienced by such patients. The omega-3 fatty acids should be administered at a dosage of between about 1 and about 30 grams per day. The two most preferred omega-3 fatty acids are eicosapentanoic acid and docosa-
15 hexanoic acid and these should typically be administered at daily dosages of 2-10 grams and 1-5 grams respectively. The fatty acids may be administered as the sole therapeutic agent or in conjunction with other agents known to be useful in the treatment of bipolar patients. In particular, the fatty acids may be administered either with a source of lithium or choline. In
20 addition, omega-3 fatty acids may be taken by patients as a component of another molecule, e.g. a triacylglycerol, and be metabolically released after ingestion.

The present invention is also directed to an omega-3 phosphatidylcholine useful in the treatment of bipolar disorder, consisting of glycerol esterified at both its α and β carbons to fatty acids. At least one, and preferably both, of these fatty acids is an omega-3 fatty acid and the γ
25 position of the glycerol must be esterified to phosphocholine. It is preferred that at least one of the esterified fatty acids be either eicosapentanoic acid or docosahexanoic acid. Omega-3 phosphatidylcholines with eicosapentanoic acid esterified to the α carbon and docosahexanoic acid esterified to the β carbon and vice versa are the most preferred. In all cases, the γ position of the triacylglycerol is esterified to phosphocholine.

In another aspect, the present invention is directed to a pharmaceutical composition comprising one or more of the omega-3 phosphatidylcholines discussed above. The composition should contain sufficient triacylglycerol so that one or more unit doses provides enough agent to reduce or eliminate the symptoms associated with bipolar disorder. In some instances, lithium may be also incorporated into the composition in order to improve therapeutic effects.

The present invention is also directed to a method for treating bipolar disorder in a human patient by administering an omega-3 phosphatidylcholine. It is expected that this phosphatidylcholine will typically be administered at a dosage sufficient to provide between 1 and 30 (preferably between 2 and 8) grams of free omega-3 fatty acid to the patient. Again, administration may be carried out concurrently with the administration of other therapeutic agents such as lithium.

Detailed Description of the Invention

In the following description, reference will be made to various methodologies well-known to those skilled in the art of medicine and pharmacology. Such methodologies are described in standard reference works setting forth the general principals of these disciplines. Included among the relevant references are: Goodwin, F.K. and Jamison, K.R., Manic Depressive Illness, Oxford University Press (1990); and Bloom, F. and Kupfer, D., Psychopharmacology. The Fourth Generation of Progress, Raven Press (1994).

A. Definitions

Bipolar disorder: Bipolar disorder refers to a form of psychosis characterized by abnormally severe mood swings. The patient alternates between episodes of mania and episodes of depression.

Omega-3 fatty acids: Fatty acids are long chain aliphatic molecules beginning with a methyl group and ending with a carboxyl group. Omega-3 fatty acids contain a double bond in the third position from the methyl group. Two common, long chain omega-3 fatty acids are eicosapentanoic acid (20 carbons in length) and docosahexanoic acid (22 carbons in length). These are both found in fish oils

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.