

» Go to Healthcare Professionals site

This information is for Patients & Caregivers

Register for HERConnection

View Full Prescribing Information



What is KADCYLA?

How can KADCYLA help?

What can I expect while on KADCYLA?

Are there financial assistance programs?

What other types of support are there?

What is KADCYLA?

Safety First!

Before you start exploring, please read side effect information. » [View now](#)

IN THIS SECTION

What is KADCYLA?

» [Understanding HER2+ MBC](#)

» [How KADCYLA is different](#)

How KADCYLA is thought to work

KADCYLA PATIENT SUPPORT HOTLINE



Call the KADCYLA Support Line
1 (855) KADCYLA
or 1 (855) 523-2952

» [Learn more](#)

Frequently Asked Questions

How KADCYLA is thought to work

Step 1: KADCYLA attaches to a HER2 receptor

KADCYLA is made to find HER2-positive cells and attach to them. It tells the cells to stop growing and tells the body's **immune system** to destroy them.

[Replay](#) [Step 1](#) [Step 2](#) [Step 3](#) [Step 4](#)

Next: See how KADCYLA can help

KADCYLA Patient Support Line
1 (855) KADCYLA (523-2952)

4HER Patient Support App

For women with HER2+ Breast Cancer as spread to other parts of the body (metastatic breast cancer)

Free program for patients receiving KADCYLA

after prior treatment with trastuzumab (Herceptin) and a taxane. Prior treatment could have been for the initial treatment of breast cancer or for the treatment of cancer that had spread to other parts of the body.

! Important Safety Information

What is the most important safety information I should know about KADCYLA?

KADCYLA is not the same medicine as trastuzumab (Herceptin).

Liver problems

KADCYLA may cause severe liver problems that can be life-threatening. Symptoms of liver problems may include vomiting, nausea, eating disorder (anorexia), yellowing of the skin (jaundice), stomach pain, dark urine, or itching

Heart problems

KADCYLA may cause heart problems, including those without symptoms (such as reduced heart function) and those with symptoms (such as congestive heart failure). Symptoms may include swelling of the ankles or legs, shortness of breath, cough, rapid weight gain of greater than 5 lbs in less than 24 hours, dizziness or loss of consciousness, or irregular heartbeat

Pregnancy

If you are exposed to KADCYLA during pregnancy, contact your healthcare provider right away; you are also encouraged to enroll in the **MotHER Pregnancy Registry** by calling 1-800-690-6720

If you are a mother who is breastfeeding, you should talk with your doctor about either stopping breastfeeding or stopping KADCYLA

Contact your doctor right away if you experience symptoms associated with these side effects.

What are the additional possible serious side effects of KADCYLA?

Lung problems

KADCYLA may cause lung problems, including inflammation of the lung tissue, which can be life-threatening. Signs of lung problems may include trouble breathing, cough, tiredness, and fluid in the lungs

Infusion-related reactions

Symptoms of an infusion-related reaction may include one or more of the following: the skin getting hot or red (flushing), chills, fever, trouble breathing, low blood pressure, wheezing, tightening of the muscles in the chest around the airways, or a fast heartbeat. Your doctor will monitor you for infusion-related reactions

Serious Bleeding

KADCYLA can cause life-threatening bleeding. Taking KADCYLA with other medications used to thin your blood (antiplatelet) or prevent blood clots (anticoagulation) can increase your risk of bleeding. Your doctor should provide additional monitoring if you are taking one of these other drugs while on KADCYLA. Life-threatening bleeding may also happen with KADCYLA even when blood thinners are not also being taken

Low platelet count

Low platelet count may happen during treatment with KADCYLA. Platelets help your blood to clot. Signs of low platelets may include easy bruising, bleeding, and prolonged bleeding from cuts. In mild cases there may not be any symptoms

Nerve damage

Symptoms may include numbness and tingling, burning or sharp pain, sensitivity to touch, lack of coordination, muscle weakness, or loss of muscle function

Skin reactions around the infusion site

KADCYLA may leak from the vein or needle and cause reactions such as redness, tenderness, skin irritation, or pain or swelling at the infusion site. If this happens, it is more likely to happen within 24 hours of the infusion

How will my doctor know if KADCYLA is right for me?

You must have a HER2 test to determine if your cancer is HER2-positive before taking KADCYLA, because benefit has been shown only in patients whose tumors are HER2-positive.

What are the most common side effects of KADCYLA?

The most common side effects seen in people taking KADCYLA are:

- Tiredness
- Nausea
- Pain that affects the bones, muscles, ligaments, and tendons
- Bleeding
- Low platelet count
- Headache
- Liver problems
- Constipation
- Nosebleeds

You are encouraged to report side effects to Genentech and the FDA. You may contact Genentech by calling 1-888-835-2555. You may contact the FDA by visiting www.fda.gov/medwatch or calling 1-800-FDA-1088.

Please see KADCYLA full Prescribing Information, including Most Important Safety Information, for additional Important Safety Information.

» Go to Healthcare Professionals site

This information is for Patients & Caregivers

Register for HERConnection

View Full Prescribing Information



What is KADCYLA?

How can KADCYLA help?

What can I expect while on KADCYLA?

Are there financial assistance programs?

What other types of support are there?

What is KADCYLA?

Safety First!

Before you start exploring, please read side effect information. » [View now](#)

IN THIS SECTION

What is KADCYLA?

» [Understanding HER2+ MBC](#)

» [How KADCYLA is different](#)

How KADCYLA is thought to work

How KADCYLA is thought to work

Step 2: KADCYLA goes inside the cell

KADCYLA also goes inside the cell to keep fighting from the inside.

KADCYLA PATIENT SUPPORT HOTLINE



Call the KADCYLA Support Line
1 (855) KADCYLA
or 1 (855) 523-2952

» [Learn more](#)

Frequently Asked Questions

[Replay](#) [Step 1](#) [Step 2](#) [Step 3](#) [Step 4](#)

Next: See how KADCYLA can help

KADCYLA Patient Support Line
1 (855) KADCYLA (523-2952)

4HER Patient Support App

For women with HER2+ Breast Cancer as spread to other parts of the body (metastatic breast cancer)

Free program for patients receiving KADCYLA

after prior treatment with trastuzumab (Herceptin) and a taxane. Prior treatment could have been for the initial treatment of breast cancer or for the treatment of cancer that had spread to other parts of the body.

! Important Safety Information

What is the most important safety information I should know about KADCYLA?

KADCYLA is not the same medicine as trastuzumab (Herceptin).

Liver problems

KADCYLA may cause severe liver problems that can be life-threatening. Symptoms of liver problems may include vomiting, nausea, eating disorder (anorexia), yellowing of the skin (jaundice), stomach pain, dark urine, or itching

Heart problems

KADCYLA may cause heart problems, including those without symptoms (such as reduced heart function) and those with symptoms (such as congestive heart failure). Symptoms may include swelling of the ankles or legs, shortness of breath, cough, rapid weight gain of greater than 5 lbs in less than 24 hours, dizziness or loss of consciousness, or irregular heartbeat

Pregnancy

If you are exposed to KADCYLA during pregnancy, contact your healthcare provider right away; you are also encouraged to enroll in the **MotHER Pregnancy Registry** by calling 1-800-690-6720

If you are a mother who is breastfeeding, you should talk with your doctor about either stopping breastfeeding or stopping KADCYLA

Contact your doctor right away if you experience symptoms associated with these side effects.

What are the additional possible serious side effects of KADCYLA?

Lung problems

KADCYLA may cause lung problems, including inflammation of the lung tissue, which can be life-threatening. Signs of lung problems may include trouble breathing, cough, tiredness, and fluid in the lungs

Infusion-related reactions

Symptoms of an infusion-related reaction may include one or more of the following: the skin getting hot or red (flushing), chills, fever, trouble breathing, low blood pressure, wheezing, tightening of the muscles in the chest around the airways, or a fast heartbeat. Your doctor will monitor you for infusion-related reactions

Serious Bleeding

KADCYLA can cause life-threatening bleeding. Taking KADCYLA with other medications used to thin your blood (antiplatelet) or prevent blood clots (anticoagulation) can increase your risk of bleeding. Your doctor should provide additional monitoring if you are taking one of these other drugs while on KADCYLA. Life-threatening bleeding may also happen with KADCYLA even when blood thinners are not also being taken

Low platelet count

Low platelet count may happen during treatment with KADCYLA. Platelets help your blood to clot. Signs of low platelets may include easy bruising, bleeding, and prolonged bleeding from cuts. In mild cases there may not be any symptoms

Nerve damage

Symptoms may include numbness and tingling, burning or sharp pain, sensitivity to touch, lack of coordination, muscle weakness, or loss of muscle function

Skin reactions around the infusion site

KADCYLA may leak from the vein or needle and cause reactions such as redness, tenderness, skin irritation, or pain or swelling at the infusion site. If this happens, it is more likely to happen within 24 hours of the infusion

How will my doctor know if KADCYLA is right for me?

You must have a HER2 test to determine if your cancer is HER2-positive before taking KADCYLA, because benefit has been shown only in patients whose tumors are HER2-positive.

What are the most common side effects of KADCYLA?

The most common side effects seen in people taking KADCYLA are:

- Tiredness
- Nausea
- Pain that affects the bones, muscles, ligaments, and tendons
- Bleeding
- Low platelet count
- Headache
- Liver problems
- Constipation
- Nosebleeds

You are encouraged to report side effects to Genentech and the FDA. You may contact Genentech by calling 1-888-835-2555. You may contact the FDA by visiting www.fda.gov/medwatch or calling 1-800-FDA-1088.

Please see KADCYLA full Prescribing Information, including Most Important Safety Information, for additional Important Safety Information.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.