

(54) **TREATMENT WITH ANTI-ERBB2
ANTIBODIES**

Related U.S. Application Data

(75) Inventor: **Robert L. Cohen**, San Mateo, CA (US)

(63) Continuation of application No. 09/568,322, filed on
May 9, 2000.

(60) Provisional application No. 60/134,085, filed on May
14, 1999.

Correspondence Address:

GENENTECH, INC.
1 DNA WAY
SOUTH SAN FRANCISCO, CA 94080 (US)

Publication Classification

(51) **Int. Cl.⁷** **A61K 39/395**; A61K 31/337

(52) **U.S. Cl.** **424/143.1**; 514/449

(73) Assignee: **Genentech, Inc.**


(21) Appl. No.: **10/429,519**

(57) **ABSTRACT**

(22) Filed: **May 5, 2003**

The present invention concerns the treatment of cancer with
anti-ErbB2 antibodies.

3H4 aa 541-599
 4D5 aa 529-625
 7C2 aa 22-53
 7F3 aa 22-53



3H4 epitope (SEQ ID NO:3)

VEECRVLQGLPREYV⁵⁴¹NARHCLPCHPECQPQNGSVTCFGPEADQCVACAHYKDPPFCVAR⁵⁹⁹

4D5 epitope (SEQ ID NO:4)

LPCHPECQPQNGSVTCFGPEADQCVACAHYKDPPFCVARCPSGVKPDLSYMPIWKFPDEEGACQP⁶²⁵
⁵⁶¹

FIG._1

1 MELAAALCRWGLLLALLPPGAASTQVCTGTMKLRLPA
 38 SPETHLDMLRHLYQGCQVVQGNLELTYLPTNASLSFL
 75 QDIQEVQGYVLIAHNQVRQVPLQRLRIVRGTQLFEDN
 112 YALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEI
 149 LKGGVLIQRNPQLCYQDTILWKDIFHKNNQLALTLID
 186 TNRSRA

FIG._2

TREATMENT WITH ANTI-ERBB2 ANTIBODIES

RELATED APPLICATIONS

[0001] This application is a continuation application of non-provisional application Ser. No. 09/568,322 filed May 9, 2000 under 37 CFR 1.53(b)(1), claiming priority under 35 USC 119(e) to provisional application No. 60/134,085 filed May 14, 1999, the contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention concerns the treatment of cancer with anti-ErbB2 antibodies.

BACKGROUND OF THE INVENTION

[0003] Proto-oncogenes that encode growth factors and growth factor receptors have been identified to play important roles in the pathogenesis of various human malignancies, including breast cancer. It has been found that the human erbB2 gene (also known as HER2, or c-erbB-2), which encodes a 185-kd transmembrane glycoprotein receptor (p185^{HER2}) related to the epidermal growth factor receptor (EGFR), is overexpressed in about 25% to 30% of human breast cancer (Slamon et al., *Science* 235:177-182 [1987]; Slamon et al., *Science* 244:707-712 [1989]).

[0004] Several lines of evidence support a direct role for ErbB2 in the pathogenesis and clinical aggressiveness of ErbB2-overexpressing tumors. The introduction of ErbB2 into non-neoplastic cells has been shown to cause their malignant transformation (Hudziak et al., *Proc. Natl. Acad. Sci. USA* 84:7159-7163 [1987]; DiFiore et al., *Science* 237:178-182 [1987]). Transgenic mice that express HER2 were found to develop mammary tumors (Guy et al., *Proc. Natl. Acad. Sci. USA* 89:10578-10582 [1992]).

[0005] Antibodies directed against human ErbB2 protein and the protein encoded by the rat equivalent of the erbB2 gene (neu) have been described. Drebin et al., *Cell* 41:695-706 (1985) refer to an IgG2a monoclonal antibody which is directed against the rat neu gene product. This antibody called 7.16.4 causes down-modulation of cell surface p185 expression on B104-1-1 cells (NIH-3T3 cells transfected with the neu protooncogene) and inhibits colony formation of these cells. In Drebin et al., *PNAS (USA)* 83:9129-9133 (1986), the 7.16.4 antibody was shown to inhibit the tumorigenic growth of neu-transformed NIH-3T3 cells as well as rat neuroblastoma cells (from which the neu oncogene was initially isolated) implanted into nude mice. Drebin et al. in *Oncogene* 2:387-394 (1988) discuss the production of a panel of antibodies against the rat neu gene product. All of the antibodies were found to exert a cytostatic effect on the growth of neu-transformed cells suspended in soft agar. Antibodies of the IgM, IgG2a and IgG2b isotypes were able to mediate significant in vitro lysis of neu-transformed cells in the presence of complement, whereas none of the antibodies were able to mediate high levels of antibody dependent cellular cytotoxicity (ADCC) of the neu-transformed cells. Drebin et al. *Oncogene* 2:273-277 (1988) report that mixtures of antibodies reactive with two distinct regions on the p185 molecule result in synergistic anti-tumor effects on neu-transformed NIH-3T3 cells implanted into nude mice. Biological effects of anti-neu antibodies are reviewed in

Myers et al., *Meth. Enzym.* 198:277-290 (1991). See also WO94/22478 published Oct. 13, 1994.

[0006] Hudziak et al., *Mol. Cell. Biol.* 9(3):1165-1172 (1989) describe the generation of a panel of anti-ErbB2 antibodies which were characterized using the human breast tumor cell line SKBR3. Relative cell proliferation of the SKBR3 cells following exposure to the antibodies was determined by crystal violet staining of the monolayers after 72 hours. Using this assay, maximum inhibition was obtained with the antibody called 4D5 which inhibited cellular proliferation by 56%. Other antibodies in the panel, including 7C2 and 7F3, reduced cellular proliferation to a lesser extent in this assay. Hudziak et al. conclude that the effect of the 4D5 antibody on SKBR3 cells was cytostatic rather than cytotoxic, since SKBR3 cells resumed growth at a nearly normal rate following removal of the antibody from the medium. The antibody 4D5 was further found to sensitize p185^{erbB2}-overexpressing breast tumor cell lines to the cytotoxic effects of TNF- α . See also WO89/06692 published Jul. 27, 1989. The anti-ErbB2 antibodies discussed in Hudziak et al. are further characterized in Fendly et al. *Cancer Research* 50:1550-1558 (1990); Kotts et al. *In Vitro* 26(3):59A (1990); Sarup et al. *Growth Regulation* 1:72-82 (1991); Shepard et al. *J. Clin. Immunol.* 11(3):117-127 (1991); Kumar et al. *Mol. Cell. Biol.* 11(2):979-986 (1991); Lewis et al. *Cancer Immunol. Immunother.* 37:255-263 (1993); Pietras et al. *Oncogene* 9:1829-1838 (1994); Vitetta et al. *Cancer Research* 54:5301-5309 (1994); Sliwkowski et al. *J. Biol. Chem.* 269(20):14661-14665 (1994); Scott et al. *J. Biol. Chem.* 266:14300-5 (1991); and D'souza et al. *Proc. Natl. Acad. Sci.* 91:7202-7206 (1994).

[0007] Tagliabue et al. *Int. J. Cancer* 47:933-937 (1991) describe two antibodies which were selected for their reactivity on the lung adenocarcinoma cell line (Calu-3) which overexpresses ErbB2. One of the antibodies, called MGR3, was found to internalize, induce phosphorylation of ErbB2, and inhibit tumor cell growth in vitro.

[0008] McKenzie et al. *Oncogene* 4:543-548 (1989) generated a panel of anti-ErbB2 antibodies with varying epitope specificities, including the antibody designated TA1. This TA1 antibody was found to induce accelerated endocytosis of ErbB2 (see Maier et al. *Cancer Res.* 51:5361-5369 (1991)). Bacus et al. *Molecular Carcinogenesis* 3:350-362 (1990) reported that the TA1 antibody induced maturation of the breast cancer cell lines AU-565 (which overexpresses the erbB2 gene) and MCF-7 (which does not). Inhibition of growth and acquisition of a mature phenotype in these cells was found to be associated with reduced levels of ErbB2 receptor at the cell surface and transient increased levels in the cytoplasm.

[0009] Stancovski et al. *PNAS (USA)* 88:8691-8695 (1991) generated a panel of anti-ErbB2 antibodies, injected them i.p. into nude mice and evaluated their effect on tumor growth of murine fibroblasts transformed by overexpression of the erbB2 gene. Various levels of tumor inhibition were detected for four of the antibodies, but one of the antibodies (N28) consistently stimulated tumor growth. Monoclonal antibody N28 induced significant phosphorylation of the ErbB2 receptor, whereas the other four antibodies generally displayed low or no phosphorylation-inducing activity. The effect of the anti-ErbB2 antibodies on proliferation of SKBR3 cells was also assessed. In this SKBR3 cell prolif-

eration assay, two of the antibodies (N12 and N29) caused a reduction in cell proliferation relative to control. The ability of the various antibodies to induce cell lysis in vitro via complement-dependent cytotoxicity (CDC) and antibody dependent cellular cytotoxicity (ADCC) was assessed, with the authors of this paper concluding that the inhibitory function of the antibodies was not attributed significantly to CDC or ADCC.

[0010] Bacus et al. *Cancer Research* 52:2580-2589 (1992) further characterized the antibodies described in Bacus et al. (1990) and Stancovski et al. of the preceding paragraphs. Extending the i.p. studies of Stancovski et al., the effect of the antibodies after i.v. injection into nude mice harboring mouse fibroblasts overexpressing human ErbB2 was assessed. As observed in their earlier work, N28 accelerated tumor growth whereas N12 and N29 significantly inhibited growth of the ErbB2-expressing cells. Partial tumor inhibition was also observed with the N24 antibody. Bacus et al. also tested the ability of the antibodies to promote a mature phenotype in the human breast cancer cell lines AU-565 and MDA-MB453 (which overexpress ErbB2) as well as MCF-7 (containing low levels of the receptor). Bacus et al. saw a correlation between tumor inhibition in vivo and cellular differentiation; the tumor-stimulatory antibody N28 had no effect on differentiation, and the tumor inhibitory action of the N12, N29 and N24 antibodies correlated with the extent of differentiation they induced.

[0011] Xu et al. *Int. J. Cancer* 53:401-408 (1993) evaluated a panel of anti-ErbB2 antibodies for their epitope binding specificities, as well as their ability to inhibit anchorage-independent and anchorage-dependent growth of SKBR3 cells (by individual antibodies and in combinations), modulate cell-surface ErbB2, and inhibit ligand stimulated anchorage-independent growth. See also WO94/00136 published Jan. 6, 1994 and Kasprzyk et al. *Cancer Research* 52:2771-2776 (1992) concerning anti-ErbB2 antibody combinations. Other anti-ErbB2 antibodies are discussed in Hancock et al. *Cancer Res.* 51:4575-4580 (1991); Shawver et al. *Cancer Res.* 54:1367-1373 (1994); Arteaga et al. *Cancer Res.* 54:3758-3765 (1994); and Harwerth et al. *J. Biol. Chem.* 267:15160-15167 (1992).

[0012] A recombinant humanized anti-ErbB2 monoclonal antibody (a humanized version of the murine anti-ErbB2 antibody 4D5, referred to as rhuMab HER2 or HERCEPTIN®, has been clinically active in patients with ErbB2-overexpressing metastatic breast cancers that had received extensive prior anticancer therapy. (Baselga et al., *J. Clin. Oncol.* 14:737-744 [1996]).

[0013] ErbB2 overexpression is commonly regarded as a predictor of a poor prognosis, especially in patients with primary disease that involves axillary lymph nodes (Slamon et al., [1987] and [1989], supra; Ravdin and Chamness, *Gene* 159:19-27 [1995]; and Hynes and Stern, *Biochim Biophys Acta* 1198:165-184 [1994]), and has been linked to sensitivity and/or resistance to hormone therapy and chemotherapeutic regimens, including CMF (cyclophosphamide, methotrexate, and fluorouracil) and anthracyclines (Baselga et al., *Oncology* 11(3 Suppl 2):43-48 [1997]). However, despite the association of ErbB2 overexpression with poor prognosis, the odds of HER2-positive patients responding clinically to treatment with taxanes were greater than three times those of HER2-negative patients (Ibid).

rhuMab HER2 was shown to enhance the activity of paclitaxel (TAXOL®) and doxorubicin against breast cancer xenografts in nude mice injected with BT-474 human breast adenocarcinoma cells, which express high levels of HER2 (Baselga et al., *Breast Cancer, Proceedings of ASCO*, Vol. 13, Abstract 53 [1994]).

SUMMARY OF THE INVENTION

[0014] In a first aspect, the present invention provides a method of treating a human patient susceptible to or diagnosed with a tumor in which ErbB2 protein is expressed comprising the following steps, performed sequentially:

[0015] (a) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody and, optionally, further comprising treating the patient with a therapeutically effective amount of a chemotherapeutic agent (e.g. a taxoid, such as paclitaxel or doxorubicin);

[0016] (b) surgically removing the tumor; and then

[0017] (c) treating the patient with a therapeutically effective amount of an anti-ErbB2 antibody and/or of a chemotherapeutic agent (e.g. a taxoid, such as paclitaxel or doxorubicin).

[0018] Preferably, the tumor overexpresses ErbB2 protein and is selected from the group consisting of a breast tumor, squamous cell tumor, small-cell lung tumor, non-small cell lung tumor, gastrointestinal tumor, pancreatic tumor, glioblastoma, cervical tumor, ovarian tumor, liver tumor, bladder tumor, hepatoma, colon tumor, colorectal tumor, endometrial tumor, salivary gland tumor, kidney tumor, prostate tumor, vulval tumor, thyroid tumor, hepatic tumor, head tumor and neck tumor.

[0019] The invention further provides an article of manufacture comprising a container, a composition within the container comprising an anti-ErbB2 antibody and a package insert instructing the user of the composition to treat a patient essentially according to the above method.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIG. 1 shows epitope-mapping of the extracellular domain of ErbB2 as determined by truncation mutant analysis and site-directed mutagenesis (Nakamura et al. *J. of Virology* 67(10):6179-6191 [October 1993]; Renz et al. *J. Cell Biol.* 125(6):1395-1406 [June 1994]). The anti-proliferative MAbs 4D5 and 3H4 bind adjacent to the transmembrane domain. The various ErbB2-ECD truncations or point mutations were prepared from cDNA using polymerase chain reaction technology. The ErbB2 mutants were expressed as gD fusion proteins in a mammalian expression plasmid. This expression plasmid uses the cytomegalovirus promoter/enhancer with SV40 termination and polyadenylation signals located downstream of the inserted cDNA. Plasmid DNA was transfected into 293S cells. One day following transfection, the cells were metabolically labeled overnight in methionine and cysteine-free, low glucose DMEM containing 1% dialyzed fetal bovine serum and 25 μ Ci each of 35 S methionine and 35 S cysteine. Supernatants were harvested either the ErbB2 MAbs or control antibodies were added to the supernatant and incubated 2-4 hours at 4° C. The complexes were precipitated, applied to a 10-20% Tricine SDS gradient gel electrophoresed at 100 V. The gel

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.