

Chem.

QUARTERLY REVIEWS

UNIVERSITY OF MINNESOTA
DULUTH BRANCH LIBRARY

1954, Vol. VIII, No. 2

	Page
SOME REACTIONS OF INORGANIC IODINE COM- POUNDS	
By K. J. Morgan	123
OLEFIN OXIDATION	
By L. Bateman	147
INTERMOLECULAR FORCES AND SOME PRO- PERTIES OF MATTER	
By J. S. Rowlinson	168
THE STRUCTURE OF THE ERGOT ALKALOIDS	
By A. L. Glenn	192

LONDON:
THE CHEMICAL SOCIETY

Copyright reserved by the Chemical Society

QUARTERLY REVIEWS contains articles by recognised authorities on selected topics from general, physical, inorganic, and organic chemistry. The Journal and Annual Reports interest primarily the research worker: Quarterly Reviews is designed for a wider range of readers. It is intended that each review article shall be of interest to chemists generally and not only to workers in the particular field being reviewed.

The submission of reviews for publication is welcomed, but intending authors are advised to write in the first place to the Editor, The Chemical Society, Burlington House, Piccadilly, London, W.1. Such preliminary communications should preferably be accompanied by an outline of the ground to be covered rather than by the completed manuscript.

Price to non-Fellows 30/- per annum

This material may be protected by Copyright law (Title 17 U.S. Code)

OLEFIN OXIDATION

By L. BATEMAN

(THE BRITISH RUBBER PRODUCERS' RESEARCH ASSOCIATION,
WELWYN GARDEN CITY, HERTS.)

THE interaction of olefins with molecular oxygen is not only a subject of widespread industrial importance, but is one of the most thoroughly understood chemical processes. This Review attempts to give a broad picture of the main mechanistic features. An earlier article¹ has reviewed the historical background and has given details of the method of approach and of the earlier kinetic data which were largely responsible for opening up this field.

In Section (1) we present the generally accepted chain mechanism and the simpler rate expressions which are often obeyed. Certain quantitative comparisons of olefinic reactivity derivable on this basis are then discussed. Our main concern, however, is to show how comparatively small changes in certain mechanistic details can give rise to substantial differences in the observed kinetics—so much so that a profound change in mechanism might be imagined. Section (2) deals with rate measurements under non-stationary state conditions designed to determine the propagation- and termination-rate constants separately, and emphasises the inherent limitations to accuracy which oxidation systems present in this respect.

Sections (3), (4), and (5) are concerned with the initiation of the oxidation chain, and the part played in this by the hydroperoxide which is the primary reaction product. This behaviour of the hydroperoxide is responsible for the autocatalytic character of the oxidations, and the complexity and environmental sensitivity of its decomposition serves to complicate the kinetics of the oxidations as a whole. Attention is drawn to circumstances where the fraction of hydroperoxide undergoing decomposition is large compared with that being formed, so that the character and kinetics of the process are greatly altered despite the same fundamental reactions being involved. Section (6) describes efforts to analyse the initiation process quantitatively in order that the number of oxidation chains being started under given conditions can be specified.

Under mild conditions of oxidation, the chain is long and the fraction of the hydroperoxide which decomposes to initiate fresh chains is very small. The overall yield of hydroperoxide should thus be nearly quantitative. In Section (7) serious discrepancies are interpreted in terms of the dual reactivity of peroxy-radicals towards olefins, the consequence being that the measured rate constants are composite quantities relating to both hydrogen extraction and double-bond addition.


The allylic radicals formed on removal of an α -methylene hydrogen atom from an olefin are mesomeric and hence the derived product may

¹ Bolland, *Quart. Reviews*, 1949, **3**, 1.

consist of allylic isomerides. The behaviour of mono-olefins and 1:4-diolefins in this respect is discussed in Section (8).

1. General Kinetic Behaviour

The following reaction scheme, where RH represents the olefin with an α -methylene hydrogen atom H, r_1 is the rate of chain initiation, and the k 's are the velocity coefficients of the reactions indicated, accounts for the reaction characteristics with remarkable comprehensiveness.



The more obvious of these are : (i) high yields of the hydroperoxide, RO₂H (cf. p. 162); (ii) catalysis by light and by free-radical producing substances, indicating the free-radical nature of the reaction ;² (iii) quantum yields greater than one and a proportionality between rate and the square root of the light intensity in photo-oxidations, indicating a chain reaction with mutual destruction of two chain carriers in the termination step ;⁴ (iv) a parallelism between oxidisability and the relative ease of rupture of the C-H bond in RH, indicating the importance of a hydrogen-exchange reaction such as (3) (cf. p. 149); (v) the formation of conjugated-diene hydroperoxides from 1:4-dienes, in agreement with the generation of mesomeric R· radicals as in (2) (cf. p. 164); and (vi) the marked retardation in rate produced by phenolic compounds (among others), which interfere with the propagation process by providing an alternative and easier reaction for the RO₂· radicals that does not liberate a radical equivalent to R·.⁴

It being assumed that $k_5^2 = k_4 k_6$, the above mechanism yields the rate equation (for long chains)⁵

$$r = r_1^{1/2} k_3 k_6^{-1/2} [\text{RH}] \frac{k_2 k_6^{-1/2} [\text{O}_2]}{k_3 k_4^{-1/2} [\text{RH}] + k_2 k_6^{-1/2} [\text{O}_2]} \quad (1)$$

where r is the overall rate of oxidation and the square brackets signify concentration terms.

Oxidisability at "High" Oxygen Pressures.—When reaction (2) is so much faster than (3) that $[\text{R}] \ll [\text{RO}_2]$, termination can be assumed to occur entirely by reaction (6), and equation (1) simplifies to

$$r_\infty = r_1^{1/2} k_3 k_6^{-1/2} [\text{RH}] \quad (2)$$

Equation (2) accurately expresses the observed kinetics for most olefins at

² Bateman and Bolland, Proc. XIth International Congress of Pure and Applied Chem., 1947.

³ Bateman and Gee, Proc. Roy. Soc., 1948, A, 195, 376.

⁴ Bolland and ten Have, Trans. Faraday Soc., 1947, 43, 201; Discuss. Faraday Soc., 1947, 2, 252.

⁵ Bolland, Proc. Roy. Soc., 1946, A, 186, 218.

oxygen pressures greater than 100 mm. (the "high" pressure region)—an interesting exception being discussed later. At constant r_1 , estimates of k_3/k_6^{-1} are thus obtained from $r_{\infty}/[\text{RH}]$ and these measure the relative reactivities of different olefins in reaction (3), since k_6 is not very sensitive to changes in R (see p. 153). Bolland⁶ has in this way developed a correlation between olefinic structure and oxidisability. Referring to propene, $\text{CH}_3\text{CH}=\text{CH}_2$, at 45°, he concludes that:

(i) Replacement of one or two hydrogen atoms at (a) and/or (c) by alkyl groups increases k_3 by $3 \cdot 3^n$, where n is the total number of substituents; similar replacement at (b) is without effect.

(ii) Replacement of a hydrogen atom at (a) by a phenyl group increases k_3 23-fold.

(iii) Replacement of a hydrogen atom at (a) by an alk-1-enyl group increases k_3 107-fold.

(iv) The value of k_3 appropriate to an α -methylene group contained in a cyclic structure is 1.7 times that of the group contained in an analogous acyclic structure.

These rules relate to broad variations, as implied by the assumed equivalency of different alkyl groups. The assumptions that k_6 is invariable and that benzoyl peroxide (used as a standard initiator) initiates throughout with equal efficiency also introduce second-order uncertainties. More serious discrepancies occur in special cases. Thus 2:4-dimethylpent-2-ene is at least 10 times less reactive than would be predicted,⁷ presumably because of steric hindrance at C_3 —behaviour simulated in a saturated

TABLE I

Olefin, RH ^a	E_3 ^b	$\Delta H_3 = E_3/0.4$	$\Delta(\Delta H_3)$
$\text{CH}_2=\text{CH}^*\text{CH}_3$	13.5	34	0
$\text{CH}_2=\text{CH}^*\text{CH}_2\text{Alk}$	11.5	29	5
$\text{CHAlk}=\text{CH}^*\text{CH}_2\text{Alk}$	10.5	26	8
$\text{CAlk}_2=\text{CH}^*\text{CH}_2\text{Alk}$	9	23	11
$\text{CH}_2=\text{CH}^*\text{CH}_2\text{Ph}$	10	25	9
$\text{CHAlk}=\text{CH}^*\text{CH}_2\text{CH}=\text{CHAlk}$	6	15	19
$\text{CH}=\text{CH}^*\text{CH}_2$	9.5	24	10
Alk—Alk			

All values are in kcal./mole

Reactive α -methylene group indicated by an asterisk. ^b Calculated from $E_3 = E_0 - \frac{1}{2}E_1 + \frac{1}{2}E_6$, where E_0 designates the overall activation energy of oxidations catalysed by benzoyl peroxide. E_6 is taken as zero and E_1 as 30 kcal./mole (Bolland's published values are based on $E_1 = 31$ kcal./mole).

⁶ Bolland, *Trans. Faraday Soc.*, 1950, **46**, 358. ⁷ Morris, unpublished result.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.