

agricultural
ent level of
ent.

ed

efficients of
Friction Be-
and Various
Unpublished
Agricultural
ate College,

coefficients
l materials.
ing 35 (6):

. Coefficient
r corn and
ublished re-
-52R1, Inter-
mpany, Chi-

s. C. W. Sys-
Bright Leaf
North Caro-
riment Sta-
Engineering
No. 14

ter, W. E.
green Vir-
aves. Part
weight and
cco Science

The Oxidation of Nicotine. I.

Kinetics of the Liquid Phase Reaction Near Room Temperature

Robert H. Linnell
Department of Chemistry, University of Vermont
Burlington, Vermont, U.S.A.

50798 2191

Introduction

Although the biochemical degradation and oxidation of nicotine has been extensively investigated(1), the kinetics of the reaction of oxygen with nicotine has not previously been

reported. Obabko(2) reports that heating tobacco containing 25-28% H₂O (in an inert N₂ atmosphere) to 70-75°C. does not deprive the tobacco of its ability to bind O₂ from air under fermentation conditions

(35°C. and 75% relative humidity) although oxidase is destroyed. The heat treated tobacco absorbed O₂ intensely initially but decreased with time whereas non-heat treated tobacco reaches a maximum O₂ uptake rate after some time and then decreases. Recently Wada(3) has oxidized nicotine at 30°C. with pure O₂ and found about 20% of the nicotine oxidized in 4 weeks. He found ammonia, methyl amine, nicotinic acid, myosmine, cotinine, nicotyrine and unidentified products.

Experimental

Nicotine was vacuum distilled in a Todd Column packed with glass helixes and the fraction at 97-98° (10 mm.) taken (reflux ratio 10:1). Freshly distilled, water-clear nicotine turns yellow in a few hours when exposed to air, but when sealed off in a glass ampule under vacuum it remains colorless for months. Preliminary work was done using an AMINCO Warburg apparatus at 20-40° and with various shaking rates. It was found that with N₂ in the Warburg there was not volume change, with O₂ there was a rapid uptake of gas which was roughly 5 times the uptake rate using air; rates increased with shaking speeds and increased with temperature. Nicotine acidified with HCl, H₂SO₄ and HOAc did not have any O₂ uptake activity indicating that the free base is involved in the oxidation. To obtain kinetic data an apparatus was constructed which consisted of a flat bottom flask of about 50 ml. volume fitted with a ground glass stopper for filling and a side arm which was connected to a 100 ml. gas burette




Figure 1.

(Tobacco Science 89)

<http://leafy.library.ucsf.edu/tid/nna14d00/pdf>

containing sat. NaCl-H₂O and a 3-way stop-cock for flushing the apparatus with gas. Oxygen gas at 1 atmosphere pressure was used in all experiments. The reaction flask was mounted on a shaker which had variable amplitude and speed and was immersed in a water bath regulated to $\pm 0.1^\circ\text{C}$. Experiments were performed at various shaking amplitudes and speed with nicotine samples of 1.00-5.00 ml. A sample size of 2.00 ml. and shaking at 440 cycles/minute with an amplitude of 7-8 mm. (measured at the sample) yielded rates of oxygen uptake that were independent of shaking for the temperature range of 15-40° used in this study. Nicotine stored in contact with air decreased in rate of oxygen uptake compared to freshly vacuum distilled nicotine. Vacuum storage prevents aging but air contact during sample removal gradually aged the stored nicotine; this problem was solved by using Matheson Prepurified N₂ (8 ppm O₂) to flush out the nicotine storage flask after each sample removal and in this way identical rates could be obtained from the same nicotine over a period of one week.

Oxygen was bubbled into nicotine held at temperatures from 50-70°C. The nicotine rapidly turned brown and an infrared band at 5.9 microns, absent in pure nicotine, increased in intensity as the experiment continued. The infrared band suggests cotinine is being formed(4). Oxidation for several days and then using Frankenburg's(4) technique for isolation of cotinine indicated complex reaction products. Addition of a free radical initiator, α, α' -Azodi-isobutyronitrile (AIBN), in a similar experiment, resulted in more rapid darkening of the nicotine and similar, if not identical, reaction products. Addition of an antioxidant, 4-methyl-2, 6-di-tert-butylphenol (BHT), greatly decreased rate of coloration of nicotine on oxidation and inactivated nicotine to O₂ for several hours.

Several experiments were done on a burley tobacco containing about 5% nicotine. Samples of the tobacco were heated in an atmosphere containing N₂, air or O₂ and the nicotine was determined by the method of Cundiff(5). No change in nicotine content was found with N₂ at temperatures from 85-125°C. With O₂, the nicotine was reduced to one-half its original value in 300 hours at 85°, 24 hours at 110° and 3 hours at 125°. The oxidized tobacco had a pleasant aromatic odor.

Table 1. Linear Rates of Oxygen Absorbed by Liquid Nicotine

Run	Temp. °C.	Ml. O ₂ (STP) x 10 ³ sec ⁻¹ x ml. nicotine ⁻¹	Induction period sec x 10 ⁻³	Remarks
IX-20	15	1.08	5.60	Nicotine stored 24 hrs. under N ₂ .
XIV-24	20	1.35	2.46	Nicotine stored 10 days under N ₂ .
V-18	25	1.61	1.60	Nicotine fresh from vacuum dist. No storage.
VII-19	25	1.61	2.12	Nicotine stored 24 hrs. under N ₂ .
XIII-23	25	1.59	2.22	Nicotine stored 10 days under N ₂ .
XV-25	25	1.47	1.98	Nicotine stored 14 days under N ₂ .
XVI-25	25	1.54	1.70	Nicotine stored 14 days under N ₂ .
XVII-26	25	none	29	Added 50 mg. BHT
XII-22	30	1.73	1.58	Nicotine stored 4 days under N ₂ .
VIII-20	35	2.09	0.86	Nicotine stored 24 hrs. under N ₂ .
X-21	35	2.10	0.88	Nicotine stored 3 days under N ₂ .
XI-21	40	2.34	0.64	Nicotine stored 4 days under N ₂ .
II-16	25	1.21	2.10	Nicotine yellow from storage one month in poor vacuum.
III-16	25	0.80	—	Reused nicotine from II-16. immediately.
IV-16	25	0.18	—	Reused nicotine from III-16 24 hours later.

Results and Discussion

In Figure 1 typical plots of volume O₂ used vs. time are shown. Table 1 tabulates all the results where rates are those from the linear part of the reaction. An Arrhenius plot of the maximum rates of table 1 yields an activation energy of 6.81 kcal (least squares). This is a low activation energy compared to the uncatalyzed oxidation of hydrocarbons such as tetralin where the overall activation energy is 14.9 \pm 0.6 kcal from 65-130°C.(6) If the linear rates are extrapolated back to zero oxygen uptake, and this time called the induction period $t_{ind.}$, a plot of $\log(1/t_{ind.})$ vs $1/T$ (assuming first order kinetics) would yield an activation energy for the removal of inhibitor (initiation). A value of 14.5 kcal is found. The induction period could be due to formation and decomposition of a nicotine hydroperoxide but it is premature to as-

sociate an activation energy with this proposed mechanism. A small apparent evolution of gas in the initial periods of lower temperature rate studies is probably due to decreased gas solubility in the nicotine at lower temperatures.

In the experiments reported here, 2.00 ml. nicotine, at room temp., was used, corresponding to 1.88 x 10⁻⁴ moles. For the consumption of one mole O₂ per mole nicotine we would therefore require 421 ml. O₂ at STP. Most experiments were run to a total O₂ consumption of 20-30 ml. (STP) and inhibition (deviation from linear rate) was noted at about 15 ml. (STP) O₂ consumption. The low O₂/nicotine mole ratio at which inhibition is found (about 0.03) together with inhibition by BHT and catalysis by AIBN provides strong evidence of the free-radical nature of the oxidation. This indicates nicotine oxidation follows the general mechanism

NCTN 86/05

Nicotine

marks
stored
under N₂.
stored
under N₂.
fresh
um dist.
ge.
stored
nder N₂.
stored
nder N₂.
stored
under N₂.
stored
nder N₂.
mg. BHT
stored
der N₂.
stored
der N₂.
stored
der N₂.
stored
der N₂.
fellow
ge one
poor

otinine
y.
otinine
; 24

with this
all appar-
he initial
ture rate
decreased
a at lower

ted here,
mp., was
8 x 10⁻²
n of one
we would
at STP.
to a total
(STP)
m linear
15 ml.
low O₂/
inhibi-
together
atalysis
vidence
the oxi-
ine oxi-
hanism

of olefin oxidation(7) and a nicotine hydroperoxide may be expected as an intermediate. The proposed hydroperoxide could be unstable at room temperature, yielding H₂O₂ (which could form nicotine N-oxide), cotinine, H₂O, and radicals which could continue the chain reaction and form complex polymers. Further work is underway to isolate the proposed hydroperoxide and provide more details on the mechanism of the reaction.

Acknowledgment

The author thanks the Tobacco Industry Research Committee which provided financial assistance making this work possible.

Literature Cited

1. McKennis, H. Jr., Turnbull, L. B., Wingfield, H. N. Jr., and Dewey, L. J., *J. Am. Chem. Soc.* 80, 1634 (1958); *ibid* 6597.
2. Obabko, V. A., *Vsesoyuz. Nauch.-Issledovatel. Inst. Tabach. i Makhoroch. Prom. No.* 142, 105-72

(1940); *ibid.* pp. 176-207. *Chem. Abs.* 36, 1439.

3. Wada, E., Kasaki, T. and Saito, K., *Arch. Biochem. and Biophys.* 79, 124 (1959).
4. Frankenburg, W. G. and A. A. Vaitekunas, *J. Am. Chem. Soc.* 79, 149 (1957).
5. Cundiff, R. H. and Markunas, P. C., *Anal. Chem.* 27, 1950 (1955).
6. George, P., Rideal, E. K. and Robertson, A., *Proc. Roy. Soc.* 185, 288 (1946).
7. Bateman, L., *Quart. Rev.* 8, 147 (1954).



50798 2193