

Cardiovascular effects of inhaled nitric oxide in patients with left ventricular dysfunction.

E Loh, J S Stamler, J M Hare, J Loscalzo and W S Colucci

Circulation. 1994;90:2780-2785

doi: 10.1161/01.CIR.90.6.2780

Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 1994 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://circ.ahajournals.org/content/90/6/2780>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:

<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation* is online at:

<http://circ.ahajournals.org/subscriptions/>

Downloaded from <http://circ.ahajournals.org/> by guest on February 27, 2013

Cardiovascular Effects of Inhaled Nitric Oxide in Patients With Left Ventricular Dysfunction

Evan Loh, MD; Jonathon S. Stamler, MD; Joshua M. Hare, MD;
Joseph Loscalzo, MD, PhD; Wilson S. Colucci, MD

Background Pulmonary vascular resistance (PVR) is frequently elevated in patients with advanced heart failure. Nitric oxide (NO), which contributes to the activity of endothelium-derived relaxing factor, causes relaxation of pulmonary arteries and veins in vitro. Inhalation of NO gas causes pulmonary vasodilation in patients with primary and secondary forms of pulmonary hypertension.

Methods and Results To test the hypothesis that inhalation of NO gas lowers PVR in patients with heart failure, we studied the hemodynamic effects of a 10-minute inhalation of NO (80 ppm) in 19 patients with New York Heart Association class III (n=5) and class IV (n=14) heart failure due to left ventricular (LV) dysfunction. Although inhalation of NO had no effect on pulmonary artery pressures, the PVR decreased by $31 \pm 7\%$ ($P < .001$) due to a $23 \pm 7\%$ increase ($P < .001$) in

pulmonary artery wedge pressure and despite a $4 \pm 2\%$ ($P < .05$) decrease in cardiac index. The magnitude of the decrease in PVR with inhaled NO was inversely related ($r = -.713$; $P < .001$) to the baseline PVR. Inhaled NO had no effect on heart rate, systemic arterial pressure, systemic vascular resistance, or LV peak $+dP/dt$ or $-dP/dt$.

Conclusions In patients with heart failure due to LV dysfunction, inhalation of NO causes a decrease in the PVR associated with an increase in LV filling pressure. These findings predict that inhaled NO, if used alone at this dose (80 ppm), may have adverse effects in patients with LV failure. (*Circulation*. 1994;90:2780-2785.)

Key words • nitric oxide • lung • heart failure • endothelium-derived factors

The endothelium plays an essential role in the dynamic regulation of vascular tone by synthesizing and releasing a variety of substances, one of which, endothelium-derived relaxing factor (EDRF), has the physicochemical properties of nitric oxide (NO) or a closely related substance.^{1,2} Endogenous NO produced by endothelial cells diffuses into neighboring vascular smooth muscle cells, where it binds to the heme component of guanylyl cyclase, thereby activating the enzyme, resulting in increased cyclic GMP production and relaxation.^{3,4} Arterial and venous endothelial cells in the pulmonary vasculature produce NO constitutively and in response to a variety of stimuli.⁵⁻⁸ NO appears to be involved both in the regulation of basal pulmonary vascular resistance (PVR)^{9,10} and in counterregulating the effects of vasoconstrictor substances.¹¹⁻¹⁵

PVR is frequently increased in patients with advanced heart failure. The underlying mechanism for increased PVR in heart failure is not known, but it almost certainly involves activation of vasoconstrictor pathways by the sympathetic nervous system, the renin-angiotensin system, and/or endothelin.^{16,17} Although there is evidence that endothelium-dependent vasodilation is impaired in the systemic vasculature of both animal models¹⁸ and patients with heart failure,¹⁹⁻²² it is

not known whether this mechanism contributes to increased PVR.

Inhalation of NO gas causes pulmonary vasodilation in patients with primary pulmonary hypertension²³ and pulmonary hypertension secondary to congenital heart disease²⁴ and to adult respiratory distress syndrome.²⁵ These observations suggest that inhaled NO might ameliorate pulmonary vasoconstriction, and they led to our hypothesis that inhalation of NO would lower PVR in patients with heart failure. To test this hypothesis, we studied the hemodynamic effects of a 10-minute inhalation of NO (80 ppm) in 19 patients with moderate to severe heart failure secondary to LV dysfunction from idiopathic or ischemic dilated cardiomyopathy.

Methods

Study Population

Nineteen patients with New York Heart Association functional class III (n=5) or IV (n=14) heart failure were studied. All patients were receiving digitalis, diuretics, and angiotensin-converting enzyme inhibitors. There were 15 men and 4 women, with a mean age of 52 ± 3 years. The cause of heart failure was ischemic cardiomyopathy in 10 patients and idiopathic dilated cardiomyopathy in 9. The peak VO_2 averaged 9.9 ± 1.6 mL \cdot kg⁻¹ \cdot min⁻¹. The study protocol was approved by the Committee for the Protection of Human Subjects from Research Risks at the Brigham and Women's Hospital, and written informed consent was obtained in all cases.

Hemodynamic Measurements

Vasodilators, converting enzyme inhibitors, digitalis, and diuretics were withheld on the morning of the catheterization. A 7F Swan-Ganz catheter (Arrow International, Inc) was placed in the pulmonary artery. Femoral artery pressure was monitored via an 8F side-arm sheath (Cordis Laboratories). In 10 patients, a 7F micromanometer-tipped pigtail catheter

Received June 20, 1994; revision accepted August 7, 1994.

From the Cardiovascular (E.L., J.M.H., J.L., W.S.C.) and Respiratory (J.S.S.) Divisions, Departments of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Mass.

Correspondence to Wilson S. Colucci, MD, Cardiovascular Division, Brigham and Women's Hospital, 75 Francis St, Boston, MA 02115.

© 1994 American Heart Association, Inc.

TABLE 1. Hemodynamic Effects of Inhaled NO in Patients With Congestive Heart Failure (n=19)

	Room Air	NO	P
HR, bpm	90±3	93±3	NS
MAP, mm Hg	79±3	81±3	NS
SVR, dyne · s · cm ⁻⁵	1102±104	1041±97	NS
PA, mm Hg	35±4	37±4	NS
PAWP, mm Hg	25±3	31±4	<.001
LVEDP, mm Hg; n=10	28±4	34±5	.02
PVR, dyne · s · cm ⁻⁵	226±30	119±13	<.001
PA-PAWP, mm Hg	11±1	6±0.5	<.001
SVI, mL/m ²	26±2	24±2	.03
CI, L · min ⁻¹ · m ⁻²	2.3±0.2	2.1±0.2	.03

HR indicates heart rate; bpm, beats per minute; MAP, mean arterial pressure; SVR, systemic vascular resistance; PA, mean pulmonary artery pressure; PAWP, pulmonary artery wedge pressure; LVEDP, left ventricular end-diastolic pressure; PVR, pulmonary vascular resistance; SVI, stroke volume index; and CI, cardiac index.

(Millar Industries) was placed in the left ventricle (LV), allowing for simultaneous dP/dt and right heart pressure measurements. The ECG, femoral artery pressure, pulmonary artery pressure, and LV pressure were recorded on a strip chart recorder (Electronics for Medicine, PPG Biomedical Systems Division). Cardiac output was determined by the Fick method, based on the measured oxygen uptake (model MRM 2B, Waters Instruments, Inc) and oxygen content in the pulmonary and femoral arteries.²⁶ Oxygen content was calculated from the blood hemoglobin and oxygen saturation by standard methods.²⁶ Blood oxygen saturation was determined in duplicate samples on a Ciba-Corning model 270 Co-oximeter. LV peak +dP/dt (+dP/dt) and peak -dP/dt (-dP/dt) were computed on-line by an Electronics for Medicine amplifier (model 220A). Values for heart rate, arterial pressure, pulmonary arterial pressure, pulmonary artery wedge pressure, LV systolic pressure, LV end-diastolic pressure




Fig 1. Bar graph showing effect of inhalation of NO gas (80 ppm, 10 minutes) on pulmonary artery (PA) pressures in 19 patients with heart failure secondary to left ventricular dysfunction. Measurements were made after the patients inhaled room air (shaded bars) or NO (solid bars) from a face mask for 10 minutes. *P<.001 vs room air.




Fig 2. Graph showing pulmonary artery wedge pressure before and after a 10-minute inhalation of room air or NO. *P<.001 vs room air.

(LVEDP), and LV +dP/dt and -dP/dt were calculated by averaging at least 50 consecutive beats under each experimental condition.

Inhalation of Nitric Oxide

NO gas (800 ppm) and N₂ (Airco) were mixed by use of a standard low-flow blender (Low Flow MicroBlender, Bird Products Corp) before introduction into the inspiratory limb of a closed breathing circuit attached to a face mask. The inhaled concentrations of NO and oxygen were regulated separately. The inhaled O₂ concentration was measured directly with an on-line oximeter (Ohmeda Oximeter). The inhaled concentrations of NO, nitrogen dioxide (NO₂), and the higher oxides of nitrogen (NO_x) were measured continuously by a chemiluminescence technique (Chemiluminescent NO_x-NO₂ Analyzer, Thermo Environmental Instruments, Inc). The exhaled gases were scavenged by a vacuum system.

To establish baseline conditions, patients inhaled room air (FIO₂, 21%; N₂, 79%) via the closed face mask system for 10 minutes before the baseline hemodynamic measurements. Patients then inhaled NO at 80 ppm (FIO₂, 21%; N₂, 79%) via




Fig 3. Graph showing effect of NO inhalation on pulmonary vascular resistance (PVR). *P<.001 vs room air.

face mask for 10 minutes, and hemodynamic measurements were repeated.

Statistical Methods

All data are presented as the mean \pm SEM. Differences between two observations for one variable within the same group were determined by two-tailed paired *t* test. Differences between groups were determined by two-tailed unpaired *t* test. Differences were considered significant if the null hypothesis could be rejected at the .05 probability level.

Results

Hemodynamic Effect of Inhaled NO

Baseline measurements during inhalation of room air revealed moderate LV failure with elevation of the LVEDP and mean pulmonary artery wedge pressure, and reduced stroke volume and cardiac indexes (Table 1). There was moderate reactive pulmonary hypertension, with an average PVR of 226 ± 30 dyne \cdot sec \cdot cm $^{-5}$.

Inhalation of NO caused no change in heart rate, mean systemic arterial pressure, systemic vascular resistance, or pulmonary artery pressure (systolic, diastolic, or mean) but caused a $23\pm 7\%$ increase in the mean pulmonary artery wedge pressure (Table 1, Figs 1 and 2) associated with $4\pm 2\%$ and $7\pm 2\%$ decreases in cardiac index and stroke volume index, respectively (Table 1). The mean transpulmonary pressure gradient decreased by $35\pm 7\%$ (Table 1), and the PVR decreased by $31\pm 7\%$ (Table 1 and Fig 3).

The decrease in PVR was due to the increase in pulmonary artery wedge pressure, as shown by the correlation ($r=-.848$, $P=.0001$) between the changes in PVR and pulmonary artery wedge pressure (Fig 4A) and lack of correlation with changes in pulmonary artery pressure (Fig 4B; $r=.13$) or cardiac index (Fig 4C; $r=.04$). The increase in mean pulmonary artery wedge pressure was due to an increase in LV filling pressure, as shown by the correlation ($r=.939$, $P<.0001$) between the changes in LV end-diastolic pressure and pulmonary artery wedge pressure with inhaled NO (Fig 5).

TABLE 2. Hemodynamic Characteristics of Patients With a Change in Pulmonary Artery Wedge Pressure Above or Below the Median With Inhalation of NO

	% PAWP <0.26 (n=9)	% PAWP >0.26 (n=10)	P
HR, bpm	87 \pm 4	94 \pm 3	NS
MAP, mm Hg	75 \pm 3	84 \pm 3	.02
SVR, dyne \cdot s \cdot cm $^{-5}$	987 \pm 153	1218 \pm 148	NS
PA, mm Hg	29 \pm 5	42 \pm 5	.02
PAWP, mm Hg	21 \pm 4	28 \pm 4	.02
SVI, mL/m 2	30 \pm 2	21 \pm 2	.004
CI, L \cdot min $^{-1}$ \cdot m $^{-2}$	2.6 \pm 0.2	1.9 \pm 0.2	.01
PVR, dyne \cdot s \cdot cm $^{-5}$	138 \pm 23	295 \pm 40	.002
LVEDD, cm	6.2 \pm 0.4	7.1 \pm 0.3	.04
$\dot{V}O_2$	9.6 \pm 0.1	11.7 \pm 0.8	NS

LVEDD indicates left ventricular end-diastolic dimension; $\dot{V}O_2$, peak oxygen consumption. Other abbreviations as in Table 1. n=19 for all parameters except EDD (n=16) and $\dot{V}O_2$ (n=17).

Hemodynamic Determinants of an Increase in Pulmonary Artery Wedge Pressure With Inhaled NO

The most prominent hemodynamic effect of NO inhalation was the increase in pulmonary artery wedge pressure (median increase, 26%). In the 10 patients with an increase in pulmonary artery wedge pressure of $\geq 26\%$ (mean increase, $33\pm 7\%$), the baseline pulmonary artery pressure, pulmonary vascular resistance, and LV end-diastolic dimension (by M-mode echocardiography; n=16) were higher and the cardiac index and stroke volume index were lower than in the 9 patients with an increase of $< 26\%$ (Table 2). Thus, more severe LV dysfunction (as evidenced by higher left heart filling pressures, lower stroke volume, and larger LV cavity size) was present in the patients who had the largest increases in pulmonary artery wedge pressure with inhaled NO.




Fig 4. Scatterplots of regression analyses depicting the relation between the change in pulmonary vascular resistance (PVR) with NO (vs room air) and the change in pulmonary artery wedge pressure (PAWP) (left), mean PA pressure (middle), or cardiac index (right) in 19 patients.

The baseline PVR was more than twofold higher in the group that had the largest increases in pulmonary artery wedge pressure with inhaled NO (Table 2), suggesting that resting PVR might be a determinant or predictor of the response to inhaled NO. Consistent with this view, there was a strong correlation ($r = -.713$, $P < .001$) between the baseline PVR and the decrease in PVR with inhaled NO (Fig 6).

As an alternative approach to this issue, we identified a subgroup of 5 patients who had "compensated" LV failure, as defined by a pulmonary artery wedge pressure ≤ 18 mm Hg (mean, 12 ± 2 mm Hg) and a cardiac index ≥ 2.5 $L \cdot \text{min}^{-1} \cdot \text{m}^{-2}$ (mean, 2.8 ± 0.3 $L \cdot \text{min}^{-1} \cdot \text{m}^{-2}$). In these patients, inhalation of NO has no effect on pulmonary artery wedge pressure ($+7 \pm 3\%$) or PVR ($+5 \pm 13\%$). In the remaining 14 patients with "decompensated" LV failure (mean pulmonary artery wedge pressure, 30 ± 2 mm Hg; mean cardiac index, 1.9 ± 0.1 $L \cdot \text{min}^{-1} \cdot \text{m}^{-2}$), inhalation of NO increased the pulmonary artery wedge pressure by $27 \pm 3\%$ ($P < .001$) and decreased the PVR by $43 \pm 7\%$ ($P < .001$).

Effects of Inhaled NO on LV Function

Since it has been suggested that NO can depress the contractile function of isolated cardiac myocytes,²⁷ we considered the possibility that inhaled NO exerted a negative inotropic effect on the LV. A negative inotropic effect of inhaled NO was suggested by a decrease in stroke volume index despite an increase in pulmonary artery wedge pressure (Fig 7A). However, in the 10 patients in whom it was measured, inhaled NO had no effect on LV peak $+dP/dt$, despite increasing LVEDP by 8 ± 1 mm Hg (Fig 7B). LV peak $-dP/dt$, which reflects isovolumic relaxation in the absence of changes in loading conditions or heart rate,^{28,29} was also not affected by inhaled NO (baseline, 807 ± 140 mm Hg/s; NO, 800 ± 139 mm Hg/s; $P = \text{NS}$; $n = 10$).

Discussion

The major finding of this study is that in patients with reactive pulmonary arterial hypertension secondary to LV failure, inhalation of NO causes reciprocal changes in the PVR (decrease) and LV filling pressure (increase). In patients with primary pulmonary hypertension, inhalation of NO causes a decrease in pulmonary artery pressure.²³ In contrast, in patients with LV failure, we found that inhalation of NO is associated not with a decrease in pulmonary artery pressure, but rather, with an increase in LV filling pressure that accounts for the decrease in PVR. Preliminary reports from two other groups^{30,31} also indicate a similar effect of inhaled NO on LV filling pressure in patients with LV failure.

The observed decrease in transpulmonary artery pressure gradient, particularly in the setting of no change or a small decrease in cardiac output, indicates that inhaled NO caused pulmonary vasodilation. NO diffuses readily through tissues, and therefore inhalation of NO may increase the concentration of NO in the vicinity of vascular smooth muscle cells in pulmonary resistance vessels, thereby exerting a direct vasodilator effect.

We believe that the NO-induced increase in LV filling pressure is due to a small increase in LV volume that




Fig 5. Scatterplot showing relation between the percent changes in pulmonary artery wedge pressure (PAWP) and left ventricular end-diastolic pressure (LVEDP) with inhaled NO in 10 patients.

occurred secondary to an increase in pulmonary venous return to the LV. For a given pulmonary artery pressure, a decrease in PVR will result in an increase in the net driving force for LV filling. Although an increase in LV volume would result in increases in ejection fraction and stroke volume in a normal LV, in our patients LV function was severely depressed and may have been on the flat portion of the Starling relation. In addition, an NO-induced increase in LV volume may have increased the magnitude of functional mitral regurgitation that is present in the majority of such hearts.^{32,33} Thus, an NO-induced redistribution of blood from the right ventricle to the LV may occur with no increase, or even a small decrease, in stroke volume. Since the failing LV often operates on the steep portion of the diastolic pressure/volume relation, a substantial increase in LV filling pressure might reflect only a small NO-induced increase in LV volume.




Fig 6. Scatterplot showing relation between the baseline pulmonary vascular resistance (PVR) and the percent change in PVR after inhalation of NO in 19 patients.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.