

Copolymer 1 induces T cells of the T helper type 2 that crossreact with myelin basic protein and suppress experimental autoimmune encephalomyelitis

(immunoregulation/multiple sclerosis/cytokines)

RINA AHARONI, DVORA TEITELBAUM, MICHAEL SELA*, AND RUTH ARNON

Department of Immunology, The Weizmann Institute of Science, Rehovot 76100, Israel

Contributed by Michael Sela, August 6, 1997

ABSTRACT The synthetic amino acid copolymer copolymer 1 (Cop 1) suppresses experimental autoimmune encephalomyelitis (EAE) and is beneficial in multiple sclerosis. To further understand Cop 1 suppressive activity, we studied the cytokine secretion profile of various Cop 1-induced T cell lines and clones. Unlike T cell lines induced by myelin basic protein (MBP), which secreted either T cell helper type 1 (Th1) or both Th1 and Th2 cytokines, the T cell lines/clones induced by Cop 1 showed a progressively polarized development toward the Th2 pathway, until they completely lost the ability to secrete Th1 cytokines. Our findings indicate that the polarization of the Cop 1-induced lines did not result from the immunization vehicle or the *in vitro* growing conditions, but rather from the tendency of Cop 1 to preferentially induce a Th2 response. The response of all of the Cop 1 specific lines/clones, which were originated in the (SJL/J×BALB/c)F₁ hybrids, was restricted to the BALB/c parental haplotype. Even though the Cop 1-induced T cells had not been exposed to the autoantigen MBP, they crossreacted with MBP by secretion of interleukin (IL)-4, IL-6, and IL-10. Administration of these T cells *in vivo* resulted in suppression of EAE induced by whole mouse spinal cord homogenate, in which several autoantigens may be involved. Secretion of anti-inflammatory cytokines by Cop 1-induced suppressor cells, in response to either Cop 1 or MBP, may explain the therapeutic effect of Cop 1 in EAE and in multiple sclerosis.

The existence of T lymphocyte subsets that produce distinct sets of cytokines and, as a result, perform distinct effector functions, has been well established in both mice and humans (1–3). Naive CD4⁺ lymphocytes (Th0) triggered by antigen differentiate either into T helper type 1 (Th1) or into T helper type 2 (Th2) cells that can crossregulate one another; their respective cytokines act antagonistically. Th1 cells produce interleukin (IL)-2, interferon (IFN)- γ , and tumor necrosis factor, activate cell-mediated immunity, and induce delayed-type hypersensitivity. Th2 cells produce IL-4, IL-5, IL-6, and IL-10 and down-regulate cell-mediated immunity. During the last few years many studies have implicated the preferential development and activation of Th1/Th2 subpopulations in a variety of autoimmune diseases (3–5). Experimental autoimmune encephalomyelitis (EAE) is an inflammatory autoimmune disease of the central nervous system (CNS) that serves as an animal model for multiple sclerosis (MS) (6). EAE can be induced in several animal species by immunization with myelin basic protein (MBP) (6), proteolipid protein (7), or myelin oligodendrocyte glycoprotein (8). During the progression of EAE, Th1, and not Th2, cytokines are present in the

inflammatory lesions in the CNS (4, 9). The autoreactive T cells that induce the disease generally display a Th1 phenotype, and the adoptive transfer of these Th1 cells is sufficient to induce EAE (4–6, 10, 11). On the other hand, recovery from EAE is associated with an elevation of Th2 cells and cytokines in the CNS (4, 12). Regulatory T cells that suppress the development of EAE produce cytokines that correspond to the Th2 profile and mediate their activity by secretion of these suppressive cytokines (4, 13, 14). These findings, along with the observation that Th2 cytokines can inhibit the action of Th1 cytokines (3, 4, 11), suggest that the induction and activation of Th2 cells may potentially prevent EAE and other autoimmune diseases that are mediated by Th1 cells.

Copolymer 1 (Cop 1) is a synthetic random copolymer of amino acids composed of L-Ala, L-Glu, L-Lys, and L-Tyr (15, 16). Cop 1 exerts a marked suppressive and protective effect on EAE in various animal species, including primates, and on chronic relapsing EAE. Cop 1 also was shown to slow the progression of disability and to reduce the relapse rate in MS patients (17), and it recently was approved as a drug for MS under the trade name of Copaxone. The mechanism of Cop 1 activity in EAE and MS involves high-affinity promiscuous binding to various class II major histocompatibility complex (MHC) molecules (18). This efficient MHC binding results in both competition with myelin antigens for T cell activation and induction of specific regulatory T cells. We previously have demonstrated that the unresponsiveness to EAE induced by Cop 1 is regulated by T suppressor (Ts) cells, because it can be adoptively transferred to normal recipients and abrogated by pretreatment with cyclophosphamide (19). Cop 1-specific Ts cell lines and Ts hybridomas were established from spleens of mice that had been rendered unresponsive to EAE by Cop 1 (20). These Cop 1-induced suppressor cells, or their supernatants, inhibited the *in vitro* response of an encephalitogenic line to MBP when cocultured. Furthermore, they prevented the development of EAE induced by whole mouse spinal cord homogenate (MSCH) *in vivo*. To further understand the mechanism of Cop 1 activity, we studied the cytokine secretion profile of various Cop 1-induced T cell lines and clones in response to Cop 1, as well as to the autoantigen MBP. In the following we demonstrate that Cop 1 induces CD4⁺ T cell lines/clones of the Th2 subtype, which secrete anti-inflammatory Th2 cytokines in response to either Cop 1 or MBP, and suppress EAE induced by whole spinal cord homogenate.

Abbreviations: EAE, experimental autoimmune encephalomyelitis; MS, multiple sclerosis; MSCH, mouse spinal cord homogenate; MBP, myelin basic protein; Cop 1, copolymer 1; Ts, T suppressor; Th1 and Th2, T helper types 1 and 2; IL, interleukin; IFN- γ , interferon- γ ; LN, lymph nodes; ICFA, incomplete Freund's adjuvant; CFA, complete Freund's adjuvant; CNS, central nervous system.

*To whom reprint requests should be addressed. e-mail: lisela@weizmann.weizmann.ac.il.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1997 by The National Academy of Sciences 0027-8424/97/9410821-6\$2.00/0 PNAS is available online at <http://www.pnas.org>.

MATERIALS AND METHODS

Mice. SJL/J, BALB/c and (SJL/J×BALB/c)F₁ mice were purchased from The Jackson Laboratory. Female mice, 7–12 weeks old, were used for all experiments.

Antigens and Antibodies. Cop 1 is a synthetic random basic polymer, prepared by polymerization of the *N*-carboxyanhydrides of L-alanine, γ -benzyl-L-glutamate, ϵ , ϵ -N-trifluoroacetyl L-lysine, and L-tyrosine (15) followed by removal of blocking groups. Two Cop 1 batches (batches 02095 and 55495) obtained from Teva Pharmaceutical Industries (Petach Tikva, Israel) were used through the study, with average molecular mass of 6,000 Da and 5,800 Da, respectively. MBP was isolated from spinal cords of mice or rats, as previously described (21). MSCH was prepared as previously described (19). Lysozyme from egg white was obtained from Sigma. Cell lines producing the mAbs, mouse anti-mouse I-A^d (MK-D6), were obtained from American Type Culture Collection. Monoclonal anti-mouse I-E^k antibodies crossreactive with I-E^d were purchased from Serotec.

T Cell Lines and Clones. T cell lines were established either from spleens of mice that had been rendered unresponsive to EAE by subcutaneous injection of Cop 1 or MBP (5–10 mg/mouse), emulsified in incomplete Freund's adjuvant (ICFA, Difco) 15 to 35 days earlier, or from lymph nodes (LN) of mice that had been immunized with Cop 1 or MBP (200 μ g/mouse) emulsified in complete Freund's adjuvant (CFA, Difco) supplemented with 4 mg/ml of *Mycobacterium tuberculosis* H37Ra (Difco), 10 days earlier. Cells were cultured and selected *in vitro* using the immunizing antigen (1–0.5 mg/plate), as described (20). Every 14–21 days, cells were stimulated by 3-day exposure to Cop 1 or MBP presented on syngeneic-irradiated (3,000 rad) spleen cells (50×10^6 /plate), followed by propagation in 10% supernatant of Con A-activated normal mouse spleen cells as T cell growth factor. Control lines specific to lysozyme were similarly obtained by subcutaneous immunization with lysozyme (5 mg/mouse) emulsified in ICFA and stimulated by exposures to lysozyme. Cloning of T cell lines was performed by limiting dilution at 0.3 cells/well.

Proliferation Assay. T-Cells (1.5×10^4) were cultured with irradiated spleen cells (5×10^5) and with the indicated antigens. At the end of 48-hr incubation, cultures were pulsed with 1μ Ci [³H]thymidine and harvested 6–12 hr later. Results are expressed as mean cpm thymidine incorporation for triplicate cultures. SDs were under 20% of the mean cpm.

Cytokine Assays. Spleen cells (5×10^6 /ml) or T cell from lines and clones (1×10^6 /ml), were incubated with the indicated antigens, presented on irradiated spleen cells (5×10^6 /ml), in a final volume of 1 ml. Supernatants were collected 24 hr later and assayed for cytokine levels using either indicator cells or mAbs in ELISA.

Cytokine assay by indicator cells. The presence of IL-2 or IL-4 in culture supernatants was evaluated by their ability to support the proliferation of the IL-2-dependent CTLD line and the IL-4-dependent CT4-S line, respectively. The tested supernatants were incubated with the indicator cells (1×10^4 /well) at a 1:1 dilution for 48 hr and then labeled with 1μ Ci-thymidine. Results are expressed as mean cpm thymidine incorporation for triplicate cultures, and the SDs were less than 20%.

Cytokine assay by ELISA. IL-2, IL-4, IL-6, IL-10, IFN- γ , and tumor necrosis factor- α were measured using a quantitative sandwich ELISA using pairs of mAbs obtained from Pharmingen, according to the manufacturer's instructions. The threshold detection for all cytokines was 10–50 pg/ml. Results are expressed in ng as mean concentration of duplicate culture supernatants (SDs under 20%), measured in duplicate wells by ELISA (SDs under 10%).

Induction of EAE. (SJL/J×BALB/c)F₁ 2–3-month-old female mice were injected in all four footpads with 3.5 mg/mouse spinal cord homogenate emulsified in a 1:1 ratio in CFA supplemented with 4 mg/ml H37Ra. Pertussis toxin (0.25 ml, 250 ng, Sigma) was injected intravenously, immediately after, and 48 hr later. Mice were examined daily for signs of EAE and assessed for clinical severity using a 1–5 score as described (20).

RESULTS

Cop 1-specific T cell lines were established from spleens of (SJL/J×BALB/c)F₁ mice that had been rendered unresponsive to EAE by injection of Cop 1 in ICFA 15–35 days earlier or from lymph nodes of mice that had been immunized with Cop 1 in enriched CFA 10 days earlier. The cells were cultured and selected *in vitro* by repeated exposures to Cop 1. Twelve different Cop 1-specific T cell lines and clones were generated and characterized. The lines were tested for their CD4/CD8 phenotype, and in all cases 96% of the cells were found to bear the CD4 phenotype.

IL-2 and IL-4 Secretion by Cop 1-Specific Lines. The IL-2 and IL-4 secretion of eight different Cop 1 Ts lines and clones first was measured using specific indicator cell lines. CTLD and CT4-S lines were used for measuring IL-2 and IL-4 secretion, respectively. T cell lines/clones were tested with various antigen concentrations, and the results obtained with the optimal antigen dose is presented. All of the Cop 1-specific lines and clones secreted IL-4, and not IL-2, when incubated with Cop 1 (Fig. 1). This restriction toward IL-4 secretion was observed in the Cop 1 lines and clones originating from spleens of mice that had been rendered unresponsive to EAE with Cop 1 in ICFA (S-1–4, S-22–1, S-22–5, S-2, and S-3), as well as in the Cop 1 lines originating from LN of mice that had been immunized with Cop 1 in CFA (LN-1, LN-3, and LN-7). The whole spleen cell population from the Cop 1/ICFA immunized mice (Cop1/ICFA spleen cells) secreted both IL-2 and IL-4 in response to Cop 1 (see Figs. 1 and 3A) However, after one or two cycles of stimulation with Cop 1, when Cop 1-specific lines have been established, only IL-4 could be detected. The clones originating from LN of mice immunized with Cop 1/CFA exhibited a mixed IL-2/IL-4 secretion for a longer period (IL-2 was measured for 10 stimulations), but




FIG. 1. IL-2 and IL-4 secretion by T cell lines/clones. Eight different Cop 1-specific T cell lines and clones, spleen cells of mice injected with Cop 1 in ICFA, and a rat MBP-specific line (EF-S-RBP) were cultured with Cop 1 (10 μ g/culture) or rat MBP (20 μ g/culture). The presence of IL-2 and IL-4 in the supernatants was determined by their ability to support IL-2-dependent CTLD and IL-4-dependent CT4S cell lines. Results are expressed as mean cpm of thymidine incorporation for triplicate cultures. SDs were under 20% of the mean cpm. Results are from one representative experiment of more than 10 performed.

eventually they lost their ability to secrete IL-2 as well. In contrast to the Cop 1 lines/clones, the EF-S-RBP effector line originating in mice that had been immunized with the encephalitogenic antigen rat MBP in CFA and selected using MBP, secreted IL-2 from the initial stimulation through the whole study period, when cocultured with the immunizing antigen. IL-4 could not be detected by the MBP-specific line in response to MBP (Fig. 1).

Comparison Between Cop 1- and MBP-Induced Suppressor Lines. Two different lines originating in mice that had been rendered unresponsive to EAE by injection of either Cop 1 or mouse MBP emulsified in ICFA were selected *in vitro* using the homologous antigen. Each line exhibited a high proliferation response to its specific antigen. The cytokine profile of these two lines, after three *in vitro* stimulations, was tested by using mAbs in ELISA (Fig. 2). For each line the response to the optimal antigen concentration is presented. Whereas both lines responded to the nonspecific mitogen Con A with production of high amounts of all the cytokines tested, they differed in the secretion profile obtained when each line was exposed to its specific antigen. The MBP line reacted to its specific antigen MBP by secretion of Th1 cytokines (IL-2, IFN- γ and tumor necrosis factor- α), as well as Th2 cytokines (IL-4, IL-6, and IL-10). In contrast, the Cop 1 line secreted only Th2 cytokines, but not Th1 cytokines, in response to Cop 1, although it still had the potential of producing Th1 cytokines as indicated by the response to Con A. Thus, whereas the MBP line exhibited a mixed Th1/Th2 cytokine secretion, the Cop 1 line was confined to the Th2 pathway.

Restriction of Cop 1 Specific F₁-Ts Lines. The F₁-Ts-Cop 1 lines/clones were originated from (SJL/J \times BALB/c)F₁ mice. The restriction of S-2 line to the two parental haplotypes SJL/J (H-2^s) and BALB/c (H-2^d) is shown in Table 1. Whereas the response to the mitogen Con A was not restricted to one of the parental haplotypes, a restricted pattern was observed in response to Cop 1. Maximal proliferation as well as secretion of IL-4, IL-6, and IL-10 were obtained when Cop 1 was presented on BALB/c or F₁ antigen-presenting cells (APC). On the other hand, the proliferation and Th2 secretion in response to Cop 1 on SJL/J APC were much lower, and IL-4 was not secreted when Cop 1 was presented by SJL/J. Neither IL-2 nor IFN- γ were measured in response to Cop 1 with all strains of APC. Similar results were obtained with the other Cop 1-specific lines/clones. Thus, the activity of F₁-Ts-Cop1 lines/clones toward Cop 1, manifested both by proliferation




FIG. 2. Comparison between cytokine secretion by Cop 1- and MBP-induced Ts lines. Quantitative ELISA of supernatants of two T cell lines established from spleens of mice, which had been rendered unresponsive to EAE by injection of Cop 1 or MBP in ICFA, stimulated by no antigen, Cop 1 (50 μ g/ml), MBP (100 μ g/ml), and ConA (5 μ g/ml). Results are expressed as mean cytokine concentration of duplicate wells. SDs for both assays were under 20% of the mean.

Table 1. Restriction of Cop 1-specific S-2 line to the parental haplotypes

	Antigen	F ₁	SJL/J	BALB/c
Proliferation, $\times 10^{-3}$	—	0.21	0.26	0.16
	Cop 1	19.50	4.80	21.61
	Con A	n.d.	n.d.	n.d.
IL-2, ng/ml	—	<0.05	<0.05	<0.05
	Cop 1	<0.05	<0.05	<0.05
	Con A	16.7	14.7	18.7
INF- γ , ng/ml	—	<0.05	<0.05	<0.05
	Cop 1	<0.05	<0.05	<0.05
	Con A	9.8	9.2	9.1
IL-4, ng/ml	—	<0.3	<0.3	<0.3
	Cop 1	2.0	<0.3	2.1
	Con A	10.8	10.3	11.7
IL-6, ng/ml	—	<0.1	<0.1	<0.1
	Cop 1	14.1	6.5	15.1
	Con A	36.6	36.8	35.5
IL-10, ng/ml	—	<0.2	<0.2	<0.2
	Cop 1	9.8	2.1	8.6
	Con A	12.5	10.9	11.4

The Cop 1-specific line S-2 was tested for its restriction to the parental haplotype, by comparing its proliferation and cytokine secretion in response to Cop 1 (50 μ g/ml) or Con A (5 μ g/ml), on irradiated spleen cells (5×10^6 cells/ml) from (SJL/J \times BALB/c)F₁, SJL/J, or BALB/c origin. SDs of all the indicated values were under 20%. Results are from one representative experiment of two performed. n.d., not determined.

and by Th2 cytokine secretion, was restricted to the BALB/c H-2^d haplotype.

A further analysis of the restriction of the Cop 1-specific lines/clones within the H-2^d region was performed using anti-I-A^d and anti-I-E^d mAbs (Table 2). Whereas no inhibition was obtained by antibodies to the I-E subregion of the BALB/c haplotype, the response to Cop 1 manifested by both proliferation and IL-4 secretion was strongly inhibited in the presence of I-A^d antibodies. Similarly the other Cop 1-specific lines/clones were inhibited only by antibodies to the I-A^d. Thus a restriction of the response toward Cop 1 to the I-A subregion of the H-2^d was demonstrated.

Crossreactivity of Cop 1-Specific F₁-Ts Lines/Clones with MBP. The pattern of cytokine secretion by several F₁-Ts-Cop 1 lines/clones in response to Cop 1 and to the autoantigen MBP was studied. A detailed proliferation and cytokine profile of S-2 line during its development is demonstrated in Fig. 3. Initially, when whole spleen cell population from mice that had been rendered unresponsive to EAE by Cop 1 was tested (Fig. 3A) a low response to Cop 1 was measured in comparison to the response obtained with the mitogen Con A. The spleen cells reacted to Cop 1 by proliferation and by secretion of both Th1 (IL-2 and IFN- γ) and Th2 (IL-4, IL-6, and IL-10) cytokines. On the other hand, neither proliferation nor cytokine

Table 2. Restriction of Cop 1-specific S-2 line within the H-2^d region

	Antigen	Inhibitor	
		—	Anti-I-A ^d / Anti-I-E ^d
Proliferation, cpm	—	243	748 / 390
	Cop 1	7,405	796 / 8,573
IL-4, cpm	—	28	43 / 64
	Cop 1	9,685	61 / 8,563

The Cop 1-specific line S-2 was tested for its restriction within the H-2^d region, by inhibition of the proliferation and IL-4 secretion in response to Cop 1 (10 μ g/well) presented on BALB/c irradiated spleen cells (5×10^6 /well), with mAbs specific either to the I-A^d or I-E^d region. SDs of all the indicated values were under 20%. Results are from one representative experiment of two performed.




FIG. 3. Proliferation and cytokine secretion profile of S-2 line during its development. (A) Initial response of spleen cells. (B) After 6 weeks in culture. (C) After 6 months in culture. Cells were cultured with no antigen, Cop 1 (50 $\mu\text{g}/\text{ml}$), MBP (100 $\mu\text{g}/\text{ml}$), and ConA (5 $\mu\text{g}/\text{ml}$). Proliferation was measured by thymidine incorporation for triplicate cultures. Cytokine concentration was measured by quantitative ELISA in duplicate wells for each one of duplicate culture supernatants. SDs were under 20% of the mean. Results represent one of three independent experiments.

secretion could be measured when the whole spleen cell population was incubated with MBP.

Six weeks later, after the cells had been exposed to Cop 1 for three cycles of stimulation, the response to Cop 1 was more confined (Fig. 3B). The S-2 line responded to Cop 1 by proliferation, but not by secretion of IL-2 and IFN- γ . These Th1 cytokines were secreted only in response to the mitogen Con A. On the other hand, higher amounts of IL-4, IL-6, and IL-10 were secreted by these cells in response to Cop 1, approaching the amounts induced by Con A. Moreover, S-2 cells secreted IL-4 not only in response to Cop 1 but also in response to MBP (7-fold more than the secretion with no antigen).

After 6 months of *in vitro* growth in which the S-2 line had been repeatedly exposed to Cop 1, IL-2 and IFN- γ were not detected in response to Cop 1, nor after stimulation with Con A (Fig. 3C). Although the cells did not secrete Th1 cytokines, they secreted high amounts of Th2 cytokines in response to Cop 1 and to Con A. The crossreactive response to MBP, which had been observed after 2 months only on the level of IL-4 secretion, now was demonstrated on the level of all Th2 cytokines measured (IL-4, IL-6, and IL-10). Furthermore, this

line secreted even higher amounts of Th2 cytokines in response to MBP than to Cop 1.

It should be noted that although all of the Cop 1-specific lines/clones secreted significant amounts of Th2 cytokines in response to Cop 1, they differed in the extent of crossreactivity with MBP. Other lines/clones obtained demonstrated a smaller degree of crossreactivity with MBP, usually on the level of IL-4 secretion. The crossreactivity with MBP was exhibited only by Cop 1-induced lines/clones. Thus control lysozyme specific lines, which secreted Th2 cytokines in response to lysozyme, did not respond at all to MBP or to Cop 1 (data not shown).

Suppressive Activity of Cop 1-Specific F₁-Ts Lines *in Vivo*. The ability of the Cop 1-specific lines/clones to prevent EAE induced by whole spinal cord homogenate *in vivo* was investigated. Activated F₁-Ts-Cop1 cells were injected to (SJL/J \times BALB/c)F₁ mice followed by EAE induction. The S-2 line originating from spleens of mice that had been rendered unresponsive to EAE with Cop 1 in ICFA was tested both after 2 months (Fig. 4A) and after 6 months (Fig. 4B) of *in vitro* growth. This line was significantly effective in suppressing EAE, because the mean clinical score of mice injected with the line was always lower than that of the control group. However the S-2 cells grown *in vitro* for 6 months were more effective in disease prevention than those grown for 2 months. Thus, after 2 months 20 \times 10⁶ cells inhibited only 40% and 58% of disease incidence and severity respectively, $P < 0.04$, whereas 10 \times 10⁶ cells of the line grown for 6 months inhibited 80% incidence and 86% severity of the disease, $P < 0.01$.

The Cop 1-specific clone S-22-1, originating similarly to S-2, completely prevented the disease, and none of the animals showed any signs of EAE ($P < 0.002$). This is in contrast to the lysozyme-specific line, which did not show any inhibitory effect on the disease ($P < 0.7$), even though the same number of cells were injected (Fig. 4C).




FIG. 4. Inhibition of MSCH-induced EAE by T cell lines and clones. (A) Cop 1-specific line S-2 after 2 months in culture (20 \times 10⁶/mouse). (B) S-2 line after 6 months in culture (10 \times 10⁶/mouse). (C) Cop 1-specific clone S-22-1 and lysozyme-specific line Lys-1 after 6 months in culture (15 \times 10⁶/mouse). (D) Cop 1-specific clone LN-3 after 6 months in culture (20 \times 10⁶/mouse). Cells were injected intravenously 3 days after stimulation with Cop 1 to (SJL/J \times BALB/c)F₁ mice 5–10 in a group, followed by EAE induction by MSCH. Control mice were induced with EAE alone. Results are expressed as mean daily clinical score of 5–10 mice in a group. Statistical analysis by Student's *t* test: (A) S-2 after 2 months in culture, $P < 0.04$. (B) S-2 after 6 months in culture, $P < 0.01$. (C) L-22-1, $P < 0.002$, Lys-1, $P < 0.7$. (D) LN-3 clone, $P < 0.09$.

The suppressive activity of another Cop 1-specific clone, LN-3, originating from LN of mice that had been immunized with Cop 1 in CFA and grown *in vitro* for 6 months, also was tested. As demonstrated (Fig. 4D), this clone also induced EAE suppression, but it was less effective than S-2 and S-2-1, because even after 6 months, 20×10^6 cells induced only 60% and 50% inhibition of the disease incidence and severity, respectively ($P < 0.09$).

DISCUSSION

The cumulative evidence on the protective role of Th2 cells in a variety of cellular immune responses, including autoimmune diseases, led to the design of immuno-intervention approaches aimed at inducing an antigen-specific shift from the Th1 pathogenic response to the protective Th2 response (4, 5, 13, 14, 22). In this study we demonstrated that Cop 1, which had been shown to suppress EAE as well as MS (16, 17), induces a predominant Th2 response. All 12 T cell lines and clones generated from Cop 1-immunized mice and selected for Cop 1 exhibited a Th2 cytokine profile and secreted high amounts of IL-4, IL-6, and IL-10 in response to Cop 1 (Figs. 1 and 3A, Tables 1 and 2). Secretion of tumor necrosis factor- α never was detected by Cop 1-induced cells (data not shown). IL-2 and IFN- γ were secreted when the whole spleen cell population was tested (Figs. 1 and 3). However, after the cells were repeatedly exposed to Cop 1 and T cell lines/clones were established, these Th1 cytokines could not be detected, whereas CD4⁺ Th2 cells prevailed.

The bias toward the Th2 phenotype was found in the Cop 1 lines/clones originating from spleens of mice that had been rendered unresponsive to EAE by injection of Cop 1 in ICFA 15–35 days earlier. This regimen has been previously shown to induce antigen-specific suppressor cells (19, 20), and the Cop 1 lines/clones originating in this way showed strong disposition toward the Th2 pathway, because after 1–3 cycles of stimulation with Cop 1 they already were confined to the Th2 pathway (Figs. 2 and 3B). Furthermore, even clones originating from lymph nodes of mice that had been immunized with Cop 1 in enriched CFA 10 days earlier, a procedure usually used to obtain effector T cell lines (6), exhibited this tendency to Th2 diversion. In this case a mixed Th1/Th2 secretion was detected for a longer period but eventually, those clones, too, lost the ability to secrete IL-2, and a complete shift to IL-4 secretion was observed. T cell lines obtained from LN of mice that produced Th2 cytokines also were demonstrated in the case of mice that had been immunized with altered peptide ligand of proteolipid protein and CFA (23).

It should be noted that our T cell lines were not grown in the presence of either IL-4 or neutralizing antibodies to Th1 interleukins, which frequently are used to selectively promote Th2 polarized cells (3). Rather, T cell growth factor medium that contained the whole spectrum of cytokines secreted by normal spleen cells in response to Con A was used as a growth factor source. In this way the role of the specific antigen in determining the cytokine profile could be revealed. Indeed, the effector line EF-S-RBP, secreted IL-2, and not IL-4, when cocultured with its specific antigen MBP (Fig. 1). Furthermore, even when MBP was injected according to the suppression regime (in ICFA, 1 month earlier), the MBP-specific line obtained exhibited a mixed Th1/Th2 cytokine profile (Fig. 2). In contrast, the Cop 1-induced lines were confined to the Th2 pathway already after three stimulations. These findings indicate that the polarization of Cop 1-induced lines was not obtained due to the immunization vehicle or the *in vitro* culture conditions, but rather due to the uniqueness of Cop 1, which preferentially induces Th2 protective response. This trait could stem from the polymeric nature of Cop 1, which contains multiple major histocompatibility complex class II binding epitopes, and does not require processing for presentation

(18). This results in high density of the antigen on the antigen-presenting cells, thus favoring Th2 development (3).

Of interest is the genetic restriction of the Cop 1-specific lines that were originated from (SJL/J \times BALB/c)F₁ mice to the BALB/c (H-2^d) parental haplotype (Table 1), and within the H-2^d region, to the I-A and not the I-E subregion (Table 2). This restriction was manifested by proliferation and by Th2 cytokine secretion. We demonstrated before that the unresponsiveness to EAE of the BALB/c strain results from the high level of regulatory cells because it could be abrogated by pretreatment with cyclophosphamide (19). The restriction of the Cop 1 Th2 lines to the BALB/c haplotype is also consistent with the dominant BALB/c genetic predisposition toward Th2 differentiation (24), which was reported in the literature for *Leishmania major* (4, 25) and autoimmune diabetes (4, 26).

The crossreactivity of Cop 1 with the natural autoantigen MBP, on the level of B as well as T cell response, previously was demonstrated (16, 27, 28). This crossreactivity was shown to correlate with Cop 1 suppressive activity *in vivo* (29). It was therefore interesting to find that the crossreactivity between Cop 1 and MBP also is expressed on the level of cytokine secretion (Fig. 3). This crossreactivity, manifested only on the level of the Th2 cytokine secretion, became more evident with more stimulations with Cop 1, probably due to selection of crossreactive clone from the cell line population. Thus, initially it was confined to the signature Th2 cytokine, IL-4 (Fig. 3B), but with additional stimulations with Cop 1, the crossreactivity with MBP became much more pronounced and expanded to the three Th2 cytokines tested (Fig. 3C). The S-2 line secreted even higher amounts of Th2 cytokines in response to MBP to which it had never been exposed before, than to Cop 1, which had been used for immunization and stimulation. Interestingly, whereas the response of encephalitogenic lines to MBP is Th1- and H-2^b-restricted (4, 6), the crossreactive response to MBP of our Cop 1-specific lines is Th2- and H-2^d-restricted. The way by which Cop 1 induces this crossreactive Th2-confined response has not been revealed in this study. Whether the Cop 1-specific cells recognize a common "suppressive" determinant shared by Cop 1 and MBP, but different from the encephalitogenic regions of MBP that induce Th1 inflammatory processes, or bear a mutated T cell receptor that transduces different signals as a consequence of antigen binding, is not clear at present.

The most meaningful criterion for the biological relevance of the Cop 1-specific Th2 lines is their ability to suppress the disease *in vivo*. Indeed a significant suppression was demonstrated by these cells on the development of EAE, as reflected both in the incidence and in the clinical score of the disease (Fig. 4). The relevance of Cop 1 Th2-specific lines to the *in vivo* effect induced by Cop 1 is not obvious, because whole spleen population demonstrated a mixed Th1/Th2 response (Figs. 1 and 3A). However it should be noted that adoptive transfer of protection was demonstrated using whole spleen cell population from Cop 1-treated mice as well (19). Because activated T cells cross the blood brain barrier irrespective of their antigenic specificity (30), it is possible that Cop 1-specific cells induced or stimulated by the injection of Cop 1 in the periphery, penetrate through the blood brain barrier to the CNS. The Cop 1-induced cells, which crossreact with MBP on the level of Th2 cytokines, are further stimulated by exposure to MBP, which is abundantly present in the CNS, and can be presented by Ia-inducible glia cells (30). Thus the secretion of the anti-inflammatory cytokines is carried on or even amplified in the case of the crossreactive response to MBP (Fig. 3C). Interestingly, even those Cop 1 lines that showed only small or moderate crossreactivity with MBP induced an *in vivo* protective effect on the disease (Fig. 4A and D). This effect can be attributed to the dominant Th2-inducing effect of IL-4. It was found that small amounts of IL-4 produced by T cells and accumulated at the site of the T cells can reach a threshold

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.