EXHIBIT 1003

DOCKET ALARM Find authenticated court documents without watermarks at <u>docketalarm.com</u>.



WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

A61K 31/565 // (A61K 31/565, 31:565)	A1	 (11) International Publication Number: WO 97/41868 (43) International Publication Date: 13 November 1997 (13.11.97)
 1) International Application Number: PCT/US9 12) International Filing Date: 28 April 1997 (2 30) Priority Data: 28 April 1997 (2 30) Priority Data: 28 April 1997 (2 30) Priority Data: 8 May 1996 (08.05.96) 31) Applicant: AMERICAN HOME PRODUCTS COL TION [US/US]; Five Giralda Farms, Madison, N. 0874 (US). 32) Inventor: GAST, Michael, Jay; 230 Country Phoenixville, PA 19460 (US). 34) Agents: ALICE, Ronald, W.; American Home Produ poration, Patent Law Dept 2B, One Campus Dr sippany, NJ 07054 (US) et al. 	97/0707 28.04.9' 28.04.9' U RPOR/ J 0794 y Lan ucts Co rive, Pa	 (81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: ORAL CONTRACEPTIVE

(57) Abstract

DOCKET

LARM

Α

This invention provides a method of contraception which comprises administering to a female of childbearing age for 28 days per menstrual cycle a combination of a progestin at a daily dosage equivalent to 40-125 μ g levonorgestrel and an estrogen at a daily dosage equivalent to 10-15 μ g ethinyl estradiol for 23-25 days beginning on day 1 of the menstrual cycle, followed by administering an estrogen at a daily dosage equivalent to 5-15 μ g ethinyl estradiol for 3-5 days.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

A	L	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
A	M	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
A	Т	Austria	FR	France	LU	Luxembourg	SN	Senegal
A	U	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
A	Z	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
B	A	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
B	B	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
B	E	Belgium	GN	Guinca	MK	The former Yugoslav	TM	Turkmenistan
B	F	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
B	G	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
B,	J	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
B	R	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
B	Y	Belarus	IS	Iceland	MW	Malawi	US	United States of America
C	A	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
C	F	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
C	G	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
C	Н	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
C	1	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
C	M	Cameroon		Republic of Korea	PL	Poland		
C	N	China	KR	Republic of Korea	PT	Portugal		
C	U	Cuba	KZ	Kazakstan	RO	Romania		
C	Z	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
D	E	Germany	LI	Liechtenstein	SD	Sudan		
D	К	Denmark	LK	Sri Lanka	SE	Sweden		
E	E	Estonia	LR	Liberia	SG	Singapore		

R M Find authenticated court documents without watermarks at <u>docketalarm.com</u>.

DOCKET

Δ

Α

- 1 -

ORAL CONTRACEPTIVE

BACKGROUND OF THE INVENTION

- The vast majority of oral contraceptives consist of a combination of a progestin and estrogen that are administered concurrently for 21 days followed either by a 7 day pill free interval or by the administration of a placebo for 7 days in each 28 day cycle. The most important aspects of a successful oral contraceptive product are effective contraception, good cycle control (absence of spotting and breakthrough bleeding and occurrence of withdrawal bleeding), and minimal side effects. Combination oral
- 10 contraceptives have traditionally acted by suppression of gonadotropins. In addition, it appears that the progestin component is primarily responsible for contraceptive efficacy through inhibition of ovulation, and other peripheral effects which include changes in the cervical mucus (which increase the difficulty of sperm entry into the uterus) and the endometrium (which reduce the likelihood of implantation). The estrogenic component
- 15 intensifies the anovulatory effect of the progestin, and is also important for maintaining cycle control.

Since the introduction of oral contraceptives (OCs) over a quarter-century ago, research has been directed toward developing preparations that minimize the potential for side effects while maintaining efficacy and normal menstrual patterns. The first-

- 20 generation OCs contained more progestin and estrogen than was necessary to prevent conception. Adverse hemostatic and metabolic changes, clinical problems, and side effects were associated with these high-dose preparations. In 1978, the World Health Organization (WHO) recommended that the focus of OC research should be the development of products containing the lowest possible dose levels of estrogen and
- 25 progestin.

The first reductions in steroid content in a combination pill were focused on estrogen because it, rather than progestin, was thought to be related to the most serious side effects. Reduction in progestin content followed, as evidence mounted that lowering progestin intake might lower the risk of cardiovascular complications such as

- 30 stroke and ischemic heart disease. [Kay CR, Am J Obstet Gynecol 142:762 (1982)]. However, this evidence was not as clear as that implicating estrogen in thromboembolic disorders. [Inman WHW, Br Med J 2:203 (1970); Stolley PD, Am J Epidemiol 102:197 (1975)]. The need for a balance between estrogens and progestins to minimize adverse effects on carbohydrate metabolism and on lipid and lipoprotein levels was also
- 35 recognized. [Bradley DD, N Engl J Med 299:17 (1978); Wynn V, Lancet 1:1045 (1979)]. Researchers then found that the synergistic action between progestin and

estrogen in a balanced ratio successfully inhibited ovulation at low levels of both components.

Research into low-dose progestins was advanced significantly by the development of norgestrel (Ng) and levonorgestrel (LNg). Levonorgestrel is the biologically active moiety of racemic norgestrel. It is strongly progestational, has no inherent estrogenic activity, is antiestrogenic, and possesses good biologic activity. The contraceptive effects of levonorgestrel are manifested throughout the hypothalamic-pituitary-gonadal-target organ axis.

Ethinyl estradiol (EE) is the estrogen most frequently used in combination OCs.
In attempts to fulfill the WHO objective, the dosage of EE in marketed OC formulations has been steadily reduced from that found in earlier OCs. Thromboembolic mortality decreased when the amount of synthetic estrogen in OC formulations was reduced from 100 µg to 50 µg. Subsequently, a significant reduction in fatal myocardial infarctions was reported for women using OCs with 30 µg of EE rather than 50 µg of EE. [Meade

15 TW, Br Med J 280:1157 (1980)].

DOCKE.

In keeping with the goal of reducing the total steroidal dosage, while maintaining contraceptive efficacy, good cycle control, and minimizing side effects, numerous regimens have been developed in which the progestin/estrogen combination is administered either as a fixed dosage combination (monophasic) or as biphasic or

- 20 triphasic regimens in which the dosage of the combination is varied either once or twice throughout the menstrual cycle. In these regimens, the progestin/estrogen combination is typically administered for 21 days followed by either a 7-day pill free period or the administration of a non-contraceptive placebo (or iron supplement) for 7 days. In these regimens, 3-ketodesogestrel (3-KDSG), desogestrel (DSG), levonorgestrel (LNg),
- 25 gestodene (GTD), norgestrel (NG), and norethindrone (NE) are typically used as the progestin while ethinyl estradiol (EE); 17β-estradiol, and mestranol are typically the estrogenic components.

Several examples of attempts at reducing the total steroidal dosage are provided below.

- 30 Erlich (German Patent DE 4,104,385 C1 and U.S. Patent 5,280,023) discloses sequential contraceptive regimens consisting of the administration of an estrogen which effects a disturbance of follicle stimulation, followed by the administration of a combination of a progestin/estrogen in a dose at least adequate to inhibit ovulation. The regimen is administered for a total of 28 days per cycle. It is preferred that the
- 35 estrogen is administered for 5-14 days per cycle and the progestin/estrogen combination is administered for 23-14 days per cycle, so that the total administration is for 28 days per cycle. Specific regimens include (a) 4 mg estradiol for 7 days followed by 21 days

R M Find authenticated court documents without watermarks at <u>docketalarm.com</u>.

DOCKET



Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time** alerts and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.

