

Review Article

Rational Design of Stable Lyophilized Protein Formulations: Some Practical Advice

John F. Carpenter,^{1,2,6} Michael J. Pikal,³ Byeong S. Chang,⁴ and Theodore W. Randolph^{1,5}

Received March 7, 1997; accepted May 15, 1997

KEY WORDS: protein drugs; design of formulations; lyophilization; stabilization of proteins.

WHY USE LYOPHILIZATION TO PREPARE STABLE PROTEIN DRUG PRODUCTS?

Early in the development of a protein therapeutic it is essential to design a formulation that is stable during shipping and long-term storage. Obviously, an aqueous liquid formulation is the easiest and most economical to handle during manufacturing, and is the most convenient for the end user. However, many proteins are susceptible to chemical (e.g., deamidation or oxidation) and/or physical degradation (e.g., aggregation and precipitation) in liquid formulations (1,2). It may be possible to design an aqueous formulation to slow protein degradation adequately, under controlled storage conditions (i.e., constant temperature and minimal agitation). However, during shipping, when precise control of conditions is not always feasible, products can be subjected to numerous stresses that denature proteins. These include agitation, high and low temperatures, and freezing (2). Furthermore, although a formulation and shipping system might be designed to circumvent damage from these stresses, it still may not be possible to inhibit damage sufficiently during long-term storage. For example, there are cases where conditions that minimize chemical degradation foster physical damage and vice versa (1,2). Then, conditions that provide a compromise affording the requisite long-term stability cannot be found.

All of these difficulties theoretically can be avoided with a properly prepared lyophilized formulation. In the dried solid, degradative reactions can be avoided or slowed sufficiently, such that the protein product remains stable for months or years at ambient temperatures (3–6). Furthermore, short-term

excursions in temperature control during shipping are usually not damaging to a lyophilized protein (6). Even in cases where two, or more, degradative pathways require different conditions for maximum thermodynamic stabilization, the reduced reaction rates in a dried product can allow for long-term stability. Thus, in general, whenever preformulation studies indicate that sufficient protein stability cannot be achieved in aqueous liquid formulations, lyophilization provides the most attractive alternative.

However, lyophilization requires sophisticated processing and is time consuming and expensive, relative to simply filling vials with a liquid formulation (3–10). Also—of greatest concern for the current review—without appropriate stabilizing excipient(s) most protein preparations are at least partially denatured by the freezing and dehydration stresses encountered during lyophilization (2,3–6,11–16). The result is often irreversible aggregation of a fraction of the protein population, either immediately after processing or after storage (e.g., 15,16). Because most protein drugs are delivered parenterally, only a few percent of aggregated protein will be unacceptable. Finally, simply designing a formulation that allows the protein to survive the lyophilization process does not assure stability during long-term storage in the dried solid (6,13–16). A poorly formulated lyophilized product, in which the protein is sufficiently reactive to require storage at subzero temperature, should not be considered a success.

The purpose of this mini-review is to provide some practical guidelines for designing formulations that protect proteins during freezing and drying, and that are stable during shipping and long-term storage at ambient temperatures. Also, as will be discussed briefly, formulations must be designed with consideration of the physical constraints on processing conditions needed to obtain a proper final cake with a low residual moisture. All of these issues have been reviewed previously in detail (3–10). Furthermore, relevant new reviews by us and others will appear this year in books edited by Vincent Lee and Louis Rey. We will not discuss the design and optimization of lyophilization cycles. Nor will we digress from practical advice about excipient choices to address the debates about the mechanisms by which these compounds stabilize proteins (see 2,5,6,11–16). The pharmaceutical scientist who has had extensive experience bringing lyophilized protein products to market may not benefit

¹ University of Colorado Center for Pharmaceutical Biotechnology, School of Pharmacy, University of Colorado Health Sciences Center, Denver, Colorado 80262.

² Department of Pharmaceutical Sciences, School of Pharmacy, University of Colorado Health Sciences Center, Denver, Colorado 80262.

³ School of Pharmacy, University of Connecticut, Storrs, Connecticut 06269.

⁴ Amgen, Inc., Thousand Oaks, California.

⁵ Department of Chemical Engineering, University of Colorado, Boulder, Colorado.

⁶ To whom correspondence should be addressed. (e-mail: John.Carpenter@UCHSC.edu)



greatly from this review. Rather, our goal is to provide a starting point for the researcher for whom design of stable lyophilized protein formulations is still a new and major challenge.

WHAT CONSTRAINTS GOVERN THE DESIGN OF THE FORMULATION

There are so many factors to consider when designing a proper lyophilized formulation, that the task when viewed as a whole can appear overwhelming. This need not be the case, if the major constraints governing success are well understood.

Protein Stability

First, it must be remembered that the whole reason for lyophilizing the product is because the protein of choice is unstable. The most sensitive element in the formulation is the protein, and the primary concern in formulation design must be the choice of excipients that provide optimal stability. This is the issue on which we will focus in detail below.

Final Product Configuration

Secondly, the final product configuration must be clearly defined prior to starting formulation efforts. Issues to be considered include route of administration, which is often parenteral, other agents to be co-administered to the patient, product volume, protein concentration, and whether the product can be lyophilized in vials or whether alternative systems such as syringes must be employed. Also, if the final product is intended for multi-use, it will be necessary to include a preservative in the formulation, which may reduce protein stability.

Formulation Tonicity

In choosing excipients, designing an isotonic solution might be a concern. Mannitol or glycine are usually good choices as tonicity modifiers. As explained below, these excipients are often preferable to NaCl, which due to its relatively low eutectic melting and glass transition temperatures, can make a formulation more difficult to lyophilize properly (3-10,17-20). Also, if the product has a relatively low mass of protein per vial, often it will be necessary to have a bulking agent in the formulation to prevent the protein from being lost from the vial during drying (e.g., 4-6). Mannitol or glycine can also serve this role because they usually crystallize to a substantial degree during lyophilization and form a mechanically strong cake (4-6). However, it must be realized that crystalline excipients when used alone will usually *not* provide adequate stability to most proteins during processing or storage in the dried solid (12-14).

Cake Structure

Finally, the dried product must have an elegant cake structure, which is mechanically strong and has not undergone any collapse and/or eutectic melting and in which the residual moisture is relatively low (ca. 1 g H₂O/100 g dried solid). If the product collapses, it will not only be aesthetically unacceptable, but also it could have excessively high residual moisture, and reconstitution time will be prolonged (3-6,17,18).

Product Glass Transition Temperature

Also, to assure long-term stability of the protein in the dried solid, the glass transition temperature (T_g) of the amorphous phase in the product, which contains the protein, must exceed the planned storage temperature (4-6,14-16). Since water is a plasticizer of the amorphous phase (3-6,9,17,18), low residual moisture is needed to insure that T_g is greater than the highest temperature encountered during shipping and storage (usually greater than 40°C).

Product Collapse Temperature

In general, achieving these goals requires maintaining the product temperature below its glass transition temperature during the lyophilization cycle (3-10,17,18). During primary drying, when ice is sublimed, the product must be maintained below the collapse temperature, which usually coincides with the thermotropic transition that has been referred to the glass transition temperature of the maximally freeze-concentrated amorphous phase of the sample (T_g') or as the softening temperature of the amorphous phase (T_s) (3-10,17,18). Also, it is necessary to keep the product temperature below the eutectic melting temperature of any crystalline component. In practice, these temperatures can be determined using either differential scanning calorimetry (DSC) or freeze-drying microscopy. The ability to determine collapse temperature is essential to formulation development (19,20).

Drying a product below the collapse temperature carries a price (3-10). The lower the sample temperature, the slower and more expensive the drying cycle becomes. In general, freeze-drying below -40°C is not practical (3-10). Also, there are physical limits in the temperatures to which samples can be reduced, which are dependent on the lyophilizer and sample configuration (3-10). As the formulation is being developed, the pharmaceutical scientist should work closely with the process engineers, who will be designing the lyophilization cycles. It is especially important to know how the large-scale lyophilizers, which will be used for commercial production, compare to the research-scale unit that is used during formulation development. Often, the large units do not have the same level of control of process parameters as do the small research units, and in part due to the large size of a production unit, intertrial variation in product temperature during the process may be greater. Finally, input from a researcher knowledgeable in the physics of freeze-drying will help prevent the formulation scientist from arbitrarily rejecting useful formulations. There are ways (see reviews 3-10) in which the process parameters can be manipulated such that relatively rapid and controlled drying can be achieved with products that have relatively low collapse temperatures.

It is clear that one goal of formulation design is to provide the highest collapse temperature that is practical, within the constraints of maintaining protein stability. The collapse temperature (i.e., the T_g') of the product will be dictated primarily by the formulation composition. If the protein is present at a level exceeding about 20% (wt/wt) of all solute it can have a relatively large effect on T_g' . Although it is often difficult to measure the T_g' of pure protein solutions with DSC, it has been found that adding increasing amounts of protein to most formulations leads to a higher T_g' (S.D. Allison, B.S. Chang, T.W. Randolph, M.J. Pikal and J.F. Carpenter, unpublished

observations). By extrapolation it appears that pure protein solutions have a T_g' of about -10°C , which is much higher than that of most pure excipient solutions (e.g., T_g' of sucrose is about -32°C). Thus, from a process economy viewpoint, one desires a high ratio of protein to stabilizer in the formulation (cf. 6). However, stability normally increases as the weight ratio of stabilizer to protein increases, so typically a compromise must be made between providing a high collapse temperature and adequate protein stabilization (e.g., 2,4-6,11,16). Also, as will be discussed below, protein resistance to freezing damage often improves as the protein concentration increases (2,6,22). Thus, in general stability is best at both high protein concentration and high weight ratio of stabilizer to protein. Therefore, in turn, stability optimization may lead to very high total solids content, which creates processing difficulties. Formulations with total solids in excess of 10% (w/w) may be difficult to process (3,7-10).

Also, the manner in which the formulation is treated prior to applying a vacuum can alter the T_g' . Usually such treatment involves an annealing step, which results in removing some fraction of a given component from the amorphous phase (6). For example, if glycine is used as a crystalline bulking agent, depending on the freezing protocol, a significant fraction of the glycine molecules may remain in the amorphous phase of the sample (6). Glycine has a relatively low T_g' (e.g., ca. -42°C ; 6,20). Thus, it is important to crystallize as much as possible, which in turn should increase the T_g' of the amorphous phase and make drying more rapid and economical. To design the optimum protocol for excipient crystallization, DSC can be used to simulate the processing conditions used during freezing and annealing. This approach is described in Carpenter and Chang (6).

AT WHAT STEPS IS STABILIZATION OF THE PROTEIN REQUIRED?

Essentially every step from vial filling to final reconstitution of the dried product can damage the protein and require formulation components to inhibit degradation (1,2,6,11,21,22). During the rapid steps (e.g., filling, freezing, drying and rehydration) the major problem is usually physical damage, which is typically manifested as formation of oligomeric and/or precipitated protein molecules (1,2,6,15,21,22). Normally the transition from solution to solid slows the rate of physical changes more than it slows chemical changes, so chemical degradation in the dried solid is often the more serious storage stability problem (e.g., 6,15,16). However, protein aggregates can form during storage/reconstitution (e.g., 6,13-16). These degradative processes can be minimized if protein unfolding (here, meaning even a small fraction of the total molecular population) is inhibited during the most damaging stresses of freezing and drying (6,15,16). Thus, a primary focus of formulation design should be protecting the protein during these steps, so that the dried formulation immobilizes the native protein in a chemically inert solid matrix having both high T_g and low residual moisture (5,6,14,16).

Stabilization During Freezing

Whether a given protein is susceptible to freezing damage depends of many factors, beyond the inclusion of the appropriate

stabilizers (see below) in the formulation. In general, the three most important parameters to consider are protein concentration, buffer choice, and freezing protocol (2,6,21-24).

Increasing protein concentration leads to increased resistance to denaturation during freezing (2,6,22,24). This phenomenon can be demonstrated by simply determining the percentage protein aggregated after freeze-thawing, which varies inversely with protein concentration (e.g., 22). Normally, it would be expected that increasing protein concentration would increase aggregation, and this would be the case if the fraction of protein molecules unfolded during freezing were independent of concentration. However, it is now thought that increasing protein concentration directly reduces freezing-induced protein unfolding. It has been speculated that damage during freezing involves protein denaturation during formation of the ice-water interface (21,22). Assuming that only a finite number of protein molecules can be denatured at this interface, then increasing the initial protein concentration will lead to a smaller percentage of damaged molecules. For practical purposes, it is sufficient simply to consider protein concentration as an important variable to examine, and to include the highest possible concentration in testing during formulation development.

Buffer choice can also be critical. The main culprits here are sodium phosphate and potassium phosphate, which can undergo drastic changes in pH during freezing and annealing (6,23,24). With sodium phosphate, the dibasic form will readily crystallize, resulting in a frozen sample in which the pH in the remaining amorphous phase (containing the protein) can be reduced to 4 or lower (23,24). With potassium phosphate, the dihydrogen salt crystallizes, giving a final pH near 9 (23,24). The risk of alteration in pH and its damage to proteins can be minimized by increasing the initial cooling rate, limiting the duration of annealing steps and minimizing the buffer concentration, all of which reduce opportunity for salt crystallization (6,24). Rapid freezing, without annealing also limits the length of exposure of protein to denaturing conditions in the frozen state (6,24). Although other excipients can aid in inhibiting the pH change (24), the best approach is to avoid using sodium phosphate or potassium phosphate buffers. Buffers that have minimal pH change upon freezing include citrate, histidine and Tris (22,24; T.J. Anchordoquy and J.F. Carpenter, unpublished observations).

In studies in which complications due to buffer pH changes have been avoided, it has been found that the degree of protein damage during freezing correlates directly with cooling rate, with more damage found at higher cooling rates where surface area of ice is larger (21,22). It has been speculated that this is due to protein denaturation during the formation of the ice-water interface (21,22). More rapid cooling leads to smaller ice crystals, which have a greater surface area to volume ratio than larger crystals. Since cooling rates will usually be dictated by the physical constraints of the lyophilizer, excessively rapid cooling probably will not be a problem (3-10). However, some proteins are so sensitive to freezing, that even with slow, controlled cooling they will be denatured (e.g., 21,24).

Stabilization During Drying and Storage in the Dried Solid

Even if the entire population of protein molecules survives the freezing step, there will be denaturation during subsequent

dehydration, unless the appropriate stabilizers are added (15,16,25–28). Simply stated, removal of the protein molecule's hydration shell, which occurs during lyophilization, destabilizes the native conformation (15,16,25–28). To date, infrared spectroscopic studies with dozens of proteins have shown that, in the absence of the appropriate stabilizer(s) (e.g., sucrose) proteins will be unfolded in the dried solid (15,16,25–28). If samples are rehydrated immediately, the degree of damage (e.g., percent of aggregation) correlates directly with how "non-native" the infrared spectrum of the dried protein appeared (15,16,25–28). Thus, reducing post-rehydration damage is dependent on minimizing the unfolding during freezing and drying. Moreover, even if 100% native molecules are recovered in samples rehydrated immediately, there can be a substantial fraction of unfolded molecules in the dried solid (15,16,25–28). Intramolecular refolding during rehydration can dominate the intermolecular interactions leading to aggregation, thereby giving 100% native protein on reconstitution.

Fortunately, appropriate excipients can prevent or at least minimize unfolding, and the success of the formulation can be judged immediately by examining the protein secondary structure in the dried solid with infrared spectroscopy (15,16,25–28). More importantly, in the few studies published to date, it has been shown that stability during long-term storage in the dried solid is dependent of retention of native protein during freeze-drying (15,16). Even for samples stored at temperatures well below the formulation T_g , damage arose rapidly (e.g., within weeks) if the protein was unfolded in the dried solid. Therefore, infrared spectroscopy, which can be used immediately after lyophilization to determine if protein unfolded has arisen, should be considered another essential tool for the protein formulation scientist.

WHICH EXCIPIENTS ARE THE BEST FIRST CHOICES?

After this review of all of the dangers of lyophilization and all the factors to be considered it might seem that rapid development of a stable lyophilized formulation would be an impossible task. Fortunately with a rational approach to formulation design, most formulation problems are quickly resolved. Here we will provide the rationale for the initial choices of formulation components. In some cases, the "initial formulation" may be all that is needed for the final marketed product. The composition to be given, with various minor modifications, has already been used with success with protein drugs (e.g., 16). We wish to stress that for any lyophilized formulation, the minimum number of components necessary for protein stability and cake structure should be used. No excipient should be added unless there are data to document that it has a beneficial role in the formulation.

Specific Conditions for Stability of a Given Protein

Before choosing the appropriate "general" stabilizers, which are effective at protecting most proteins, it is absolutely essential that the formulation be optimized for the specific factors that increase the physical and chemical stability of a given protein. For example, simply avoiding extremes in pH can drastically reduce the rate of deamidation (1). Moreover, it has been found that the resistance of a protein to unfolding

during freeze-drying can be dramatically increased by optimizing the pH of solution (e.g., 15). Also, other specific ligands that increase protein stability (e.g., by increasing the free energy of unfolding) should be investigated. The stabilizing effects of heparin and other polyanions on growth factors (e.g., 29) provide a good example. Another important factor to be considered is the effect of ionic strength on protein unfolding and aggregation. It must be recognized that during freezing, the ionic strength may increase 50-fold as ice formation concentrates all solutes (5,6,9,19). The persons responsible for bulk drug purification and pharmaceutical preformulation often already have insight into these issues. Thus, it is imperative that the formulation scientist confer with these people, prior to embarking of design of a lyophilized formulation.

Even with specific solution conditions optimized for protein stability, it probably will be necessary to add other protective excipients, if the protein is to survive lyophilization and long-term storage in the dried solid. First, let us consider some compounds that have been used for lyophilized protein formulation but which do *not* provide stability and may actually foster damage during storage. We will then provide an outline of a simple, but effective formulation, and the rationale for the choice of the components will be discussed.

Excipients that Can Fail to Stabilize Proteins

With the goal of obtaining a strong cake structure during a rapid lyophilization cycle, polymers such as dextran and hydroxyethyl starch, which have relatively high collapse temperatures, are attractive excipients. Also, the T_g of the final dried product will be high (e.g., $>90^\circ\text{C}$) with these polymers (15). Unfortunately, these polymers do not inhibit protein unfolding during lyophilization and they typically fail to provide stability during subsequent storage (15,30). The failure to inhibit lyophilization-induced denaturation is presumably because the polymers are too bulky to hydrogen bond to the protein in the place of the water that is lost during dehydration and/or because the polymers form a separate amorphous phase from the protein (5,6). Although when used alone such polymers are not good choices as stabilizers, as described below, they could be prove useful in combination with certain disaccharide protein stabilizers.

Among the numerous compounds tested it appears that the most effective stabilizers during the lyophilization cycle are disaccharides (2,5,6,11,15,16,25–28). However, one group of compounds that should be avoided are the reducing sugars. These compounds may effectively inhibit protein unfolding during the lyophilization cycle, but during storage in the dried solid they have the propensity to degrade proteins via the Maillard reaction between carbonyls of the sugar and free amino groups on the protein (31). The result can be a brown syrup containing degraded protein instead of a white cake containing active protein drug. Usually, the only way to slow this process significantly is to store the product at subzero temperatures, which defeats the purpose of a lyophilized product. Compounds in this undesirable category include glucose, lactose, maltose and maltodextrins.

As noted earlier, crystalline bulking agents such as mannitol and glycine do not provide protection during lyophilization (6,11,12,25). However, some effective lyophilized formulations employing mixtures of these two agents have been developed

and marketed. In these cases the appropriate ratio of mannitol:glycine led to a significant fraction of the compounds remaining amorphous (e.g., 30). Presumably this amorphous fraction was sufficient to inhibit protein unfolding during lyophilization and to provide stability during long-term storage. However, we caution against such an approach because achieving just the right processing conditions, in combination with the appropriate excipient ratio, can be time consuming and tricky.

Rational Choice of Stabilizing Excipients

So what are appropriate, rational choices for excipients? To provide a concrete example, let's make the following assumptions about a fictitious case. 1) The protein drug will be formulated at 2 mg/ml. 2) The major routes of degradation are aggregation immediately after lyophilization/rehydration and deamidation during storage in the dried solid. 3) Optimizing specific conditions (e.g., using a citrate buffer at pH 6.0) only reduces aggregation upon freeze-drying and reconstitution to about 10% and deamidation still proceeds at an unacceptably rapid rate during storage, even when the product is stored 20°C below its T_g . 4) A crystalline bulking agent (e.g., mannitol) is desired to form a mechanically strong and elegant cake.

At this point, the major component missing is a nonreducing disaccharide, which forms an amorphous phase with the protein in the dried solid and serves as the primary stabilizer. The main choices are sucrose or trehalose (5,6). These compounds are relatively effective at protecting proteins during freezing and usually excellent at inhibiting unfolding during dehydration (5,6,15,16,25–28). Freezing protection depends on the initial bulk concentration of the sugar, and sometimes concentrations exceeding 5% (wt/vol) are needed to maximize stabilization (2,6). In contrast, protection during drying depends on the final mass ratio between the sugar and the protein (5,6). Generally, a weight ratio of sugar to protein of at least 1:1 is required for good stability, with optimal stability being reached at around 5:1. In practice, it is not necessary to determine the most appropriate concentration for each type of protection. Rather, with the protein concentration held constant, a range of sugar concentrations can be tested during formulation screening to discern the optimal concentration for retention of native protein in the dried solid and the resultant reduction in aggregation upon rehydration.

In general, the optimal sugar concentration for stabilizing the protein during lyophilization will also provide storage stability, if the final dried powder has a T_g well above the storage temperature (5,6,15,16). For example, assuming that a high room temperature of about 30°C is the maximum intended storage temperature, then a product containing a native protein and with a T_g of > 50°C should be stable. Since residual moisture lowers T_g , the condition $T_g > 50^\circ\text{C}$ must apply to the maximum water content allowed by the product specifications. It is imperative that DSC be used to measure the T_g of each product to be certain that this goal is achieved.

Both sucrose and trehalose have advantages and disadvantages. 1) Trehalose has a higher T_g at any moisture content and, thus, is more easily lyophilized (32). In addition, the condition $T_g > 50^\circ\text{C}$ will hold at higher residual water contents for trehalose. However, a skilled process engineer should be able to design economical, effective cycles for either sugar. Also, in products with a relatively high protein concentration, the

protein could contribute to an increased T_g , which serves to minimize the advantages of trehalose. 2) Trehalose is also more resistant than sucrose to acid hydrolysis. Hydrolysis of these disaccharides produces reducing sugars, which must be avoided. Usually this is not a problem, unless very low pH's of around 4 or lower are employed. 3) Sucrose appears to be more effective at inhibiting unfolding during lyophilization (unpublished observations). This difference has been most obvious when there is a relatively high protein concentration and a need to employ a relatively high initial concentration of sugar. This need can develop when the protein is very unstable during freezing and/or there is a relatively high protein content in the dried solid. Evidence to date indicates that less effective stabilization by trehalose is due to the greater propensity of this sugar to phase separate from polymers (e.g., the protein in a formulation) during freezing and drying (S.D. Allison, T.W. Randolph, B.S. Chang and J.F. Carpenter, unpublished observation). Whether this is a problem with a given formulation cannot be predicted. Hence, the capacity to protect a protein must be examined for each formulation. 4) Sucrose is commonly used in parenteral products that are approved by the Food and Drug Administration (33). In contrast, trehalose has not yet been used in an approved product, probably because to date there has not been sufficient practical benefit to justify using this sugar in place of sucrose. Safety of trehalose will most likely not be a concern. Thus, if there is a clear advantage of trehalose over sucrose in a given product, use of trehalose should not hinder regulatory approval.

At this point, our example formulation might be complete, as is the case for many proteins. However, let's assume that even with sucrose completely inhibiting detectable protein unfolding, as assessed with structural analysis of the dried solid with infrared spectroscopy, after rehydration there is still about 1% aggregated protein. Since there are no aggregates in the starting material, it is assumed that at some point during freezing and drying a very small fraction of the protein population was unfolded. During rehydration some of these molecules refolded, but others formed aggregates. This actually appears to be a common problem, as is the formation of aggregates in processing steps prior to lyophilization. Fortunately, aggregate formation can usually be inhibited by including a nonionic surfactant, such as polysorbate (Tween), in the formulation. The required concentrations generally are very low (e.g., <0.5% wt/vol). The optimal concentration can be discerned by titrating surfactant into the lyophilization formulation containing all of the other components. Excessive amounts of surfactant should be avoided, because these compounds are liquids at room temperature, and, hence, if present at high enough concentration they can lower the formulation glass transition temperatures (T_g' and T_g). Usually, however, at the very low concentrations needed to optimize protein stability this is not a problem.

The surfactant can be viewed as the "finishing touch". However, often it is advantageous to add the surfactant well before formulating specifically for lyophilization, to inhibit unfolding and aggregation induced by interfaces (e.g., as caused by air bubble entrainment or by vial-liquid interfaces) during processing (34).

Most important for our purposes, is the capacity of a surfactant to inhibit the formation of aggregates during lyophilization/rehydration. Currently it is not known at which steps the surfactant's protection is operative. Surfactants have

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.