

THE  
MERCCK  
MANUAL  
VOLUME I

GENERAL MEDICINE

15th EDITION

NPS Ex. 2166  
CFAD v. NPS  
IPR2015-01093

800-631-6888  
EXHIBIT  
2170

THE  
MERCK  
MANUAL

FIFTEENTH EDITION

1st Edition - 1899  
2nd Edition - 1901  
3rd Edition - 1905  
4th Edition - 1911  
5th Edition - 1923  
6th Edition - 1934  
7th Edition - 1940  
8th Edition - 1950

9th Edition - 1956  
10th Edition - 1961  
11th Edition - 1966  
12th Edition - 1972  
13th Edition - 1977  
14th Edition - 1982  
15th Edition - 1987

VOLUME I

GENERAL MEDICINE

FIFTEENTH EDITION

THE  
MERCK  
MANUAL

OF  
DIAGNOSIS AND THERAPY

VOLUME I

GENERAL MEDICINE

Robert Berkow, M.D., *Editor-in-Chief*  
Andrew J. Fletcher, M.B., B.Chir., *Assistant Editor*

*Editorial Board*

Philip K. Bondy, M.D.	Fred Plum, M.D.
L. Jack Faling, M.D.	John Romano, M.D.
Alvan R. Feinstein, M.D.	G. Victor Rossi, Ph.D.
Eugene P. Frenkel, M.D.	John H. Talbott, M.D.
Robert A. Hoekelman, M.D.	Paul H. Tanser, M.D.
Robert G. Petersdorf, M.D.	

*Published by*

MERCK SHARP & DOHME RESEARCH LABORATORIES

*Division of*  
MERCK & Co., INC.  
*Rahway, N.J.*

1987

MERCK & CO., INC.  
*Rahway, N.J.*  
*U.S.A.*

MERCK SHARP & DOHME  
*West Point, Pa.*

MERCK SHARP & DOHME INTERNATIONAL  
*Rahway, N.J.*

MERCK SHARP & DOHME RESEARCH LABORATORIES  
*Rahway, N.J. West Point, Pa.*

MSD AGVET DIVISION  
*Woodbridge, N.J.*

HUBBARD FARMS, INC.  
*Walpole, N.H.*

MERCK CHEMICAL MANUFACTURING DIVISION  
*Rahway, N.J.*

CALGON CORPORATION  
Water Management Division  
*Pittsburgh, Pa.*  
Commercial Division  
*St. Louis, Mo.*

KELCO DIVISION  
*San Diego, Calif.*

---

Library of Congress Catalog Card Number 1-31760  
ISBN Number 0-911910-07-7  
ISSN Number 0076-6526

First Printing—August 1987

Copyright© 1987 by MERCK & Co., Inc.  
All rights reserved. Copyright under the Universal Copyright  
Convention and the International Copyright Convention.  
Copyright reserved under the Pan-American Copyright Convention.

Printed in the U. S. A.

signs are generalized swelling and inflammation of the finger, tenderness over the flexor tendon sheaths, careful maintenance of a flexed finger position, and exquisite pain on active or passive finger extension. Fever, lymphangitis, lymphadenitis, and leukocytosis are usual. **Treatment** is surgical drainage plus antibiotics. Gram stain of the pus should dictate antibiotic choice; streptococci and staphylococci are the usual pathogens.

---

## 6. BACTEREMIA AND SEPTIC SHOCK

---

**Bacteremia** connotes *invasion of the circulation by bacteria*. The term **septicemia** is reserved for *situations in which bacteremia is associated with clinical manifestations of infection*. Bacteremia commonly, and usually transiently, accompanies various surgical manipulations (eg, incision of an abscess); or it may result from colonization of indwelling intravenous devices and urethral catheters. (For infants, see also **NEONATAL SEPSIS AND NEONATAL MENINGITIS** under **NEONATAL INFECTIONS** in Vol. II, Ch. 23.) Bacteremia may be intermittent or sustained, and may cause severe consequences. In patients who abuse IV narcotics, gram-positive bacteremia is common and may lead to right-sided bacterial endocarditis even in the absence of cardiac murmurs. The bacteremia of left-sided bacterial endocarditis is usually sustained and may be prolonged. Gram-negative bacteremia is usually intermittent and generally follows primary infection in the GU tract, biliary tree, GI tract, lungs, or, less commonly, skin, bones, or joints. In many patients with chronic diseases no primary focus of infection is apparent.

### Symptoms and Signs

Few clinical manifestations are unique to bacteremia. Although variable, fever is almost always present and may be intermittent, with wide diurnal variations (septic, or "spiking"). Chills are common at the onset. Skin eruptions are also common, and may be petechial, purpuric, papular, pustular, or vesicular. Usually gram-negative bacteremia begins abruptly with chills, fever, nausea, vomiting, diarrhea, and prostration.

### Diagnosis

The presence of bacteremia is established by blood cultures, which should be performed for both aerobic and anaerobic organisms. A single negative culture does not exclude bacteremia; moreover, in some patients, especially those with prior antibiotic therapy, blood cultures never do become positive. If the patient is not very ill, no more than 6 blood cultures should be performed. In severely ill patients, 2 blood cultures taken 30 min apart before treatment is instituted should suffice. In most cases, treatment is required before results of blood cultures become available.

### Complications

**Metastatic infection** of the meninges or of serous cavities, such as the pericardium or larger joints, may occur. Endocarditis (see also **ENDOCARDITIS** in Ch. 27) may be the sequel of bacteremia if the pathogen is a streptococcus or staphylococcus, but rarely occurs as a result of gram-negative bacteremias. **Metastatic abscesses** may occur almost anywhere and, when extensive, produce symptoms and signs characteristic of infection in the organ affected. Multiple abscess formation is particularly common with staphylococcal bacteremia. Bacteremia may result in **septic shock**, which is discussed separately, below.

### Prognosis

Transient bacteremias associated with surgical procedures, indwelling IV catheters, or urinary catheters are often undetected and probably do not require therapy. However, persistent bacteremia is dangerous; the prognosis depends on the ability to eliminate the source of infection with surgery or antibiotics, and on the status of the underlying disease. When multiple organisms are recovered consistently (polymicrobial bacteremia), a poor outcome can be expected. Bacteremia unresponsive to treatment because of inadequate antibiotic therapy, poor host resistance, or delay in diagnosis is often fatal.

**Treatment** of bacteremia is discussed below with septic shock.

## SEPTIC SHOCK

When bacteremia is associated with inadequate tissue perfusion, especially with gram-negative organisms or meningococci, **septic shock** with hypotension, vascular collapse, renal failure, and death may ensue. Septic shock usually occurs when bacteremia is due to gram-negative organisms and generally in hospitalized patients with underlying diseases. Predisposing factors include diabetes mellitus; cirrhosis; leukemia; lymphoma; disseminated carcinoma; surgical procedures; antecedent

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.