

## Evidence for a First Order Mechanism in Amide Hydrolysis

By J. W. BARNETT and CHARMIAN O'CONNOR\*

(Department of Chemistry, University of Auckland, Auckland, New Zealand)

**Summary** Hydrolysis of acetanilide derivatives in concentrated acids is confirmed as an A-1 mechanism by use of  $w$  and  $\phi$  parameters.

AMIDES generally exhibit a maximum in their hydrolysis rate profiles at moderate acidity,<sup>1</sup> and in the case of some nitro-derivatives of acetanilide this is followed by a local minimum at ca. 70–75%  $w/w$  sulphuric acid.<sup>2,3</sup> We recently reported<sup>4</sup> this phenomenon in the hydrolysis of *N*-acetylsulphanilic acid in sulphuric acid. On the basis of

molecular. Vinnik *et al.*<sup>3</sup> considered specifically that the activated complex incorporated the protonated substrate and a sulphuric acid molecule.


We now report that the phenomenon of increasing rate in concentrated acid media occurs for a large number of substituted acetanilides (I) undergoing hydrolysis in sulphuric acid, and in an extreme case in perchloric acid also. For those substrates where the rate increases were sufficiently large over an adequate range of acidity, to justify mathematical analysis, we have tested the data using the

TABLE

Typical rate data for hydrolysis of substituted acetanilides in concentrated acid media and analysis of the data by use of Bunnett  $w$  and Bunnett–Olsen linear free energy relationships.

Temp. (°C)	Position of substituent <sup>a</sup>					$10^5 k_{\psi} (\text{s}^{-1})$		$w$	$\phi$
	2	3	4	5	6	% $w/w$ $\text{H}_2\text{SO}_4$ 80.0	96.6		
100.0			$\text{SO}_3\text{H}$			14.2	99.6	–0.48	–0.39
95.3			$\text{SO}_3\text{H}$			7.36	52.0	–0.52	–0.41
90.0			$\text{SO}_3\text{H}$			5.03	30.9	–0.43	–0.38
80.3			$\text{SO}_3\text{H}$			2.09	15.5	–0.49	–0.40
80.0			$\text{SO}_3\text{H}$			1.78	6.43	–0.34	–0.26
100.0			Cl			0.499	1.12		
105.0			Cl			0.930	2.47		
100.0			Br			0.574	1.44		
105.0			Br			1.55	4.09		
100.0			I			2.29	11.9	–0.51	–0.45
100.0		OH	$\text{SO}_3\text{H}$			44.5	297	–0.44	–0.37
100.0	$\text{SO}_3\text{H}$		$\text{CH}_3$			0.905	4.31		
100.0	Br		Br		Br	258	578	–0.21	–0.15
55.0			$\text{NO}_2$			0.233 <sup>b</sup>	13.9 <sup>b</sup>		
65.0		$\text{NO}_2$		$\text{NO}_2$		1.50 <sup>b</sup>	16.7 <sup>b</sup>	–0.48	–0.47
						% $w/w$ $\text{HClO}_4$ 50.0			
100.0	Br		Br		Br	128	551	–0.52	–0.16
25.0	$\text{NO}_2$		$\text{NO}_2$			30.6	<0.1		

<sup>a</sup> Substituent is H unless stated otherwise; <sup>b</sup> interpolated from ref. 2; <sup>c</sup> interpolated from ref. 3(b).



increasing values of activation energy over this region of increasing rate, the mechanism of hydrolysis was interpreted<sup>2,3,4</sup> as having changed from bimolecular to uni-

Bunnett  $w^5$  and Bunnett–Olsen linear free energy<sup>6</sup> relationships (see Table). In all cases correlation coefficients were good (generally better than 0.99). The values of  $w$  all lie in the range –2.5 to 0.0 and the values of  $\phi$  are all negative (although some are < –0.34, the original limit of values found by Bunnett and Olsen<sup>6</sup>) and thus they are characteristic of reactions in which water is not involved in the rate determining step. These results record the first application of  $w$  and  $\phi$  values to an A-1 mechanism for amide hydrolysis.

(Received, 28th February 1972; Com. 333.)

<sup>1</sup> C. J. O'Connor, *Quart. Rev.*, 1970, **24**, 553.

<sup>2</sup> J. A. Duffy and J. A. Leisten, *J. Chem. Soc.*, 1960, 853.

<sup>3</sup> (a) M. I. Vinnik, I. M. Medvetskaya, L. R. Andreeva, and A. E. Tiger, *Russ. J. Phys. Chem.*, 1967, **41**, 128; (b) M. I. Vinnik and I. M. Medvetskaya, *ibid.*, p. 947.

<sup>4</sup> J. W. Barnett and C. J. O'Connor, *Tetrahedron Letters*, 1971, 2161.

<sup>5</sup> J. F. Bunnett, *J. Amer. Chem. Soc.*, 1961, **83**, 4956; 4968; 4973; 4978.

<sup>6</sup> J. F. Bunnett and F. Olsen, *Canad. J. Chem.*, 1966, **44**, 1899; 1917.