



US008754131B2

(12) **United States Patent**
Sawa et al.

(10) **Patent No.:** US 8,754,131 B2
(45) **Date of Patent:** *Jun. 17, 2014

(54) **AQUEOUS LIQUID PREPARATION
CONTAINING
2-AMINO-3-(4-BROMOBENZOYL)PHENYLACETIC
ACID**

(71) Applicant: **Senju Pharmaceutical Co., Ltd.**, Osaka
(JP)

(72) Inventors: **Shirou Sawa**, Hyogo (JP); **Shuhei
Fujita**, Hyogo (JP)

(73) Assignee: **Senju Pharmaceutical Co., Ltd.**, Osaka
(JP)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **14/165,976**

(22) Filed: **Jan. 28, 2014**

(65) **Prior Publication Data**

US 2014/0142183 A1 May 22, 2014

Related U.S. Application Data

(62) Division of application No. 13/687,242, filed on Nov.
28, 2012, now Pat. No. 8,669,290, which is a division
of application No. 13/353,653, filed on Jan. 19, 2012,
now Pat. No. 8,497,304, which is a division of
application No. 10/525,006, filed as application No.
PCT/JP2004/000350 on Jan. 16, 2004, now Pat. No.
8,129,431.

(30) **Foreign Application Priority Data**

Jan. 21, 2003 (JP) 2003-012427

(51) **Int. Cl.**
A01N 37/18 (2006.01)
A61K 31/165 (2006.01)
A01N 37/44 (2006.01)
A61K 31/24 (2006.01)
A01N 37/10 (2006.01)
A61K 31/19 (2006.01)

(52) **U.S. Cl.**
USPC **514/619**; 514/535; 514/570; 514/618

(58) **Field of Classification Search**
USPC 514/619, 535, 570
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,045,576 A 8/1977 Welstead, Jr. et al.
4,683,242 A 7/1987 Poser
4,910,225 A 3/1990 Ogawa et al.
5,110,493 A 5/1992 Cherng-Chyi et al.
5,475,034 A 12/1995 Yanni et al.

5,597,560 A 1/1997 Bergamini et al.
5,603,929 A 2/1997 Desai et al.
5,653,972 A 8/1997 Desai et al.
5,942,508 A 8/1999 Sawa
5,998,465 A 12/1999 Hellberg et al.
6,162,393 A 12/2000 De Bruij et al.
6,274,592 B1 8/2001 Sawa
6,274,609 B1 8/2001 Yasueda et al.
6,319,513 B1 11/2001 Dobrozsi
6,369,112 B1 4/2002 Xia
6,383,471 B1 5/2002 Chen et al.
6,395,746 B1 5/2002 Cagle et al.
2001/0056098 A1 12/2001 Sawa
2007/0082857 A1 4/2007 Sawa

FOREIGN PATENT DOCUMENTS

AU 707 119 9/1995
CA 2 013 188 9/1990
EP 0 306 984 3/1989
JP 62-126124 6/1987
JP 1-104023 4/1989
JP 02083323 3/1990
JP 2-124819 5/1990
JP 5-223052 8/1993
JP 9-503791 4/1997
JP 11-228404 8/1999
JP 2002-308764 10/2002
WO 96/14829 5/1996
WO 00/59475 10/2000
WO 01/15677 3/2001
WO 02/13804 2/2002

OTHER PUBLICATIONS

New Drugs in Japan, 2001, 2001 Edition, Published by Yakuji Nippo
Ltd., May 11, 2001, pp. 27-29, and its English translation of the
material portions.

ISTA Pharmaceuticals, "New Drug Applications: Xibrom", [http://
www.drugs.com/nda/xibrom_040525.html](http://www.drugs.com/nda/xibrom_040525.html), accessed online Sep.
19, 2007.

Nolan et al., "The Topical Anti-Inflammatory and Analgesic Proper-
ties of Bromfenic in Rodents", Agents and Actions, vol. 25, No. 1-2,
pp. 77-85, Aug. 1988.

(Continued)

Primary Examiner — Layla Soroush

(74) Attorney, Agent, or Firm — Wenderoth Lind & Ponack,
L.L.P.

(57) **ABSTRACT**

An aqueous liquid preparation of the present invention con-
taining 2-amino-3-(4-bromobenzoyl)phenylacetic acid or its
pharmacologically acceptable salt or a hydrate thereof, an
alkyl aryl polyether alcohol type polymer such as tyloxapol,
or a polyethylene glycol fatty acid ester such as polyethylene
glycol monostearate is stable. Since even in the case where a
preservative is incorporated into said aqueous liquid prepa-
ration, the preservative exhibits a sufficient preservative
effect for a long time, said aqueous liquid preparation in the
form of an eye drop is useful for the treatment of blepharitis,
conjunctivitis, scleritis, and postoperative inflammation.
Also, the aqueous liquid preparation of the present invention
in the form of a nasal drop is useful for the treatment of
allergic rhinitis and inflammatory rhinitis (e.g. chronic rhini-
tis, hypertrophic rhinitis, nasal polyp, etc.).

(56)

References Cited

OTHER PUBLICATIONS

Corrected partial English translation of New Drugs in Japan, 2001, 2001 Edition, Published by Yakuji Nippo Ltd., May 11, 2001, pp. 27-29, previously submitted on Apr. 11, 2005.

Complete English translation of New Drugs in Japan, 2001, 2001 Edition, Published by Yakuji Nippo Ltd., May 11, 2001, pp. 27-29.

Notice of Opposition dated Feb. 19, 2009 issued by EPO in connection with the corresponding European patent application and Opposition.

<http://medical-dictionary.thefreedictionary.com/prophylactic> accessed Dec. 15, 2009.

H. Scott et al., "Comparing the Surface Chemical Properties and the Effect of Salts on the Cloud Point of a Conventional Nonionic Surfactant, Octoxynol 9 (Triton X-100), and of its Oligomer, Tyloxapol (Triton WR-1339)", Journal of Colloid and Interface Science, vol. 205, pp. 496-502, 1998.

1


**AQUEOUS LIQUID PREPARATION
CONTAINING
2-AMINO-3-(4-BROMOBENZOYL)PHENYLACETIC
ACID**

TECHNICAL FIELD

The present invention relates to an aqueous liquid preparation containing 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof. More particularly, the present invention relates to an aqueous liquid preparation containing 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof and an alkyl aryl polyether alcohol type polymer or a polyethylene glycol fatty acid ester.

BACKGROUND ART

Benzoylphenylacetic acid derivatives including bromfenac (generic name) of formula (I):



of which chemical name is 2-amino-3-(4-bromobenzoyl)phenylacetic acid are known as disclosed in JP-A-23052/1977 and its corresponding U.S. Pat. No. 4,045,576. 2-Amino-3-(4-bromobenzoyl)phenylacetic acid, its pharmacologically acceptable salt and a hydrate thereof are known as a non-steroidal anti-inflammatory agent, and they are effective against inflammatory diseases of anterior or posterior segment of the eye, such as blepharitis, conjunctivitis, scleritis, and postoperative inflammation in the field of ophthalmology, and its sodium salt has been practically used in the form of eye drops ("New Drugs in Japan, 2001", 2001 Edition, Published by Yakuji Nippo Ltd., May 11, 2001, p. 27-29).

The eye drop as mentioned above is designed to stabilize 2-amino-3-(4-bromobenzoyl)phenylacetic acid by means of addition of a water-soluble polymer (e.g. polyvinylpyrrolidone, polyvinyl alcohol, etc.) and a sulfite (e.g. sodium sulfite, potassium sulfite, etc.) (Japanese patent No. 2,683,676 and its corresponding U.S. Pat. No. 4,910,225).

In addition, as an eye drop other than the above-mentioned one, Japanese patent No. 2,954,356 (corresponding to U.S. Pat. Nos. 5,603,929 and 5,653,972) discloses a stable ophthalmic composition which comprises incorporating an antibacterial quaternary ammonium polymer and boric acid into an acidic ophthalmic agent. The acidic agent described therein includes, for example, 2-amino-3-(4-bromobenzoyl)phenylacetic acid.

Further, in Japanese patent No. 2,954,356, there is the following description—"Benzalkonium chloride is a widely used preservative in ophthalmic solutions. However, benzalkonium chloride and other quaternary ammonium compounds are generally considered to be incompatible with ophthalmic compositions of drugs with acidic groups, such as nonsteroidal anti-inflammatory drugs. These preservatives

2

In these prior art references, there is no disclosure that alkyl aryl polyether alcohol type polymers or polyethylene glycol fatty acid esters are able to stabilize an aqueous liquid preparation of 2-amino-3-(4-bromobenzoyl)phenylacetic acid or its pharmacologically acceptable salt, and inhibit decrease in preservative effect of benzalkonium chloride and other quaternary ammonium compounds.

DISCLOSURE OF THE INVENTION

It is an object of the present invention to provide an aqueous liquid preparation comprising 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof, which is stable within a pH range giving no irritation to eyes and in which, when a preservative such as benzalkonium chloride is incorporated therein, preservative effect of the preservative does not substantially deteriorate.

Another object of the invention is to provide a method for stabilizing an aqueous liquid preparation of 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof.

Further object of the invention is to provide an aqueous liquid preparation comprising 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof and a preservative wherein, when specifically a quaternary ammonium salt such as benzalkonium chloride is incorporated as a preservative, decrease in preservative effect of said preservative is inhibited.

As a result of various studies, the inventors of the present invention have found that, by adding, for example, an alkyl aryl polyether alcohol type polymer such as tyloxapol, or a polyethylene glycol fatty acid ester such as polyethylene glycol monostearate to an aqueous liquid preparation of 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof, the aqueous solution becomes stable within a pH range giving no irritation to eyes, and change of the 2-amino-3-(4-bromobenzoyl)phenylacetic acid over time can be inhibited, and furthermore, when the aqueous solution contains a preservative, deterioration in the preservative effect of said preservative can be inhibited for a long period of time. The inventors of the present invention have further studied extensively and completed the present invention.

Namely, the present invention relates to:

- (1) An aqueous liquid preparation comprising 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof, and an alkyl aryl polyether alcohol type polymer or a polyethylene glycol fatty acid ester,
- (2) The aqueous liquid preparation according to the above (1), wherein the alkyl aryl polyether alcohol type polymer has a polymerization degree of 3 to 10, the alkyl contains 1 to 18 carbon atoms, the aryl is a phenyl residue, and the polyether alcohol is represented by the formula $O(CH_2CH_2O)_xH$ in which X is an integer of 5 to 100,
- (3) The aqueous liquid preparation according to the above (1) or (2), wherein the alkyl aryl polyether alcohol type polymer is tyloxapol,
- (4) The aqueous liquid preparation according to the above (1), wherein the carbon number of the fatty acid in the polyethylene glycol fatty acid ester is 12 to 18,
- (5) The aqueous liquid preparation according to the above (1) or (4), wherein the polyethylene glycol fatty acid ester is polyethylene glycol monostearate,

3

aryl polyether alcohol type polymer is selected from a range of minimum concentration of 0.01 w/v to maximum concentration of 0.5 w/v

(7) The aqueous liquid preparation according to any one of the above (1), (2) or (4), wherein the concentration of the polyethylene glycol fatty acid ester is selected from a range of minimum concentration of 0.02 w/v % to maximum concentration of 0.1 w/v %,

(8) The aqueous liquid preparation according to any one of the above (1) to (7), wherein the concentration of the 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof is 0.01 to 0.5 w/v,

(9) The aqueous liquid preparation according to any one of the above (1) to (8), wherein benzalkonium chloride is contained as a preservative,

(10) The aqueous liquid preparation according to anyone of the above (1) to (9), wherein the pharmacologically acceptable salt of 2-amino-3-(4-bromobenzoyl)phenylacetic acid is a sodium salt,

(11) The aqueous liquid preparation according to any one of the above (1) to (10), wherein the pH of the aqueous liquid preparation is within a range of 7 to 9,

(12) The aqueous liquid preparation according to the above (11), wherein the pH of the aqueous liquid preparation is within a range of 7.5 to 8.5,

(13) The aqueous liquid preparation according to any one of the above (1) to (12), wherein the aqueous liquid preparation is an eye drop,

(14) The aqueous liquid preparation according to any one of the above (1) to (12), wherein the aqueous liquid preparation is a nasal drop,

(15) An eye drop comprising sodium 2-amino-3-(4-bromobenzoyl)phenylacetate hydrate and 0.01 to 0.5 w/v % of tyloxapol,

(16) An eye drop comprising sodium 2-amino-3-(4-bromobenzoyl) phenylacetate hydrate and 0.02 to 0.1 w/v of polyethylene glycol monostearate,

(17) A method for stabilizing 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof in an aqueous liquid preparation, which comprises incorporating tyloxapol or polyethylene glycol monostearate into an aqueous liquid preparation containing 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof, and

(18) A method for inhibiting decrease in preservative effect of a preservative in an aqueous liquid preparation of 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof, which comprises incorporating tyloxapol or polyethylene glycol monostearate into an aqueous liquid preparation containing 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof and a preservative.

According to the present invention, a stable aqueous liquid preparation containing 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof can be prepared by incorporating an alkyl aryl polyether alcohol type polymer such as tyloxapol, or a polyethylene glycol fatty acid ester such as polyethylene glycol monostearate into an aqueous liquid preparation con-

4

thereof. Also, an aqueous liquid preparation of the present invention, wherein a preservative is incorporated, has a sufficient preservative effect.

Therefore, the aqueous liquid preparation of the present invention is advantageously used as an eye drop for the treatment of, for example, blepharitis, conjunctivitis, scleritis, and postoperative inflammation. In addition, such aqueous liquid preparation can be used as a nasal drop for the treatment of, for example, allergic rhinitis and inflammatory rhinitis (e.g. chronic rhinitis, hypertrophic rhinitis, nasal polyp, etc.).

The pharmacologically acceptable salt of 2-amino-3-(4-bromobenzoyl)phenylacetic acid includes, for example, an alkali metal salt such as sodium salt and potassium salt, and an alkaline earth metal salt such as calcium salt and magnesium salt, among which sodium salt is especially preferable.

2-Amino-3-(4-bromobenzoyl)phenylacetic acid and its pharmacologically acceptable salt can be prepared according to the method as described in JP-A-23052/1977 (corresponding to U.S. Pat. No. 4,045,576) or by a similar method thereof. These compounds can be obtained as their hydrate depending on synthetic conditions and recrystallization conditions. The hydrate includes 1/2 hydrate, 1 hydrate, and 3/2 hydrate, among which 3/2 hydrate is preferable.


In the aqueous liquid preparation of the present invention, the content (concentration range) of 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof is usually about 0.01 to 0.5 w/v %, preferably about 0.05 to 0.2 w/v %, especially about 0.1 w/v %, and it is preferable to appropriately vary the content depending on the purpose of use and the degree of disease to be treated.

The carbon number of the alkyl in the an alkyl aryl polyether alcohol type polymer which is a non-ionic surfactant used as a stabilizer for 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof is approximately 1 to 18. Specifically, the alkyl group includes, for example, methyl, ethyl, propyl, isopropyl, cyclopropyl, butyl, isobutyl, sec-butyl, tert-butyl, cyclobutyl, pentyl, isopentyl, neopentyl, tert-pentyl, 1-ethylpropyl, 4-methylpentyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 1,2-dimethylbutyl, 2-ethylbutyl, cyclopentyl, hexyl, cyclohexyl, heptyl, isoheptyl, octyl, isooctyl, nonyl, isononyl, decyl, isodecyl, undecyl, isoundecyl, dodecyl, isododecyl, tridecyl, isotridecyl, tetradecyl, isotetradecyl, pentadecyl, isopentadecyl, hexadecyl, isohexadecyl, heptadecyl, isoheptadecyl, octadecyl, isooctadecyl, and isomers thereof, among which octyl and its isomer (e.g. isooctyl, sec-octyl, 1-methylheptyl, 1-ethylhexyl, 2-ethylhexyl, 1-propylpentyl, 1,5-dimethylhexyl, 1,1,3,3-tetramethylbutyl, etc.) are preferable, and 1,1,3,3-tetramethylbutyl which is an isomer of octyl groups is especially preferable.

The aryl in the alkyl aryl polyether alcohol type polymer can be preferably a phenyl residue. The polyether alcohol can be represented by the formula $O(CH_2CH_2O)_xH$ in which X is an integer of 5 to 100, preferably 5 to 30, more preferably 8 to 10. The average polymerization degree is preferably about 3 to 10.

Among the above-mentioned alkyl aryl polyether alcohol

5



The fatty acid of the polyethylene glycol fatty acid ester which is a non-ionic surfactant used as a stabilizer for 2-amino-3-(4-bromobenzoyl)phenylacetic acid or a pharmacologically acceptable salt thereof or a hydrate thereof can be preferably a fatty acid having the carbon number of 12 to 18. Specific examples of such polyethylene glycol fatty acid esters are polyethylene glycol monostearate (e.g. polyoxyl 8 stearate, polyoxyl 40 stearate, etc.), polyethylene glycol monolaurate, polyethylene glycol monooleate, polyethylene glycol diisostearate, polyethylene glycol dilaurate, polyethylene glycol dioleate, and the like. Among these compounds, polyethylene glycol monostearate is preferable, and polyoxyl 40 stearate is especially preferable. The polyoxyl 40 stearate is a monostearic acid ester of an ethylene oxide condensed polymer, and can be represented by the formula C₁₇H₃₅COO (CH₂CH₂O)_nH which is a non-ionic surfactant and n is about 40.

Although the content (concentration range) of the alkyl aryl polyether alcohol type polymer in the aqueous liquid preparation of the present invention depends on the kind of compounds used, the minimum concentration is about 0.01 w/v and the maximum concentration is about 0.5 w/v %. With respect to the tyloxapol content (concentration range), for example, the minimum content is about 0.01 w/v %, 0.02 w/v or 0.03 w/v %, and the maximum content is about 0.05 w/v %, 0.1 w/v %, 0.3 w/v % or 0.5% w/v, and preferably the minimum content is about 0.02 w/v % and the maximum content is about 0.05 w/v %.

Although the content (concentration range) of the polyethylene glycol fatty acid ester in the aqueous liquid preparation of the present invention depends on the kind of compounds used, it is within a range of about 0.02 w/v % of minimum concentration to about 0.1 w/v % of maximum concentration. For example, the content (concentration range) of polyethylene glycol monostearate is within a range of about 0.02 w/v of minimum content to about 0.1 w/v of maximum content, and preferably within a range of about 0.02 w/v % of the minimum content to about 0.05 w/v of the maximum content.

The incorporation ratio of tyloxapol in the aqueous liquid preparation of the invention is within a range of the minimum content of about 0.1 or 0.2 part by weight to the maximum content of about 0.5, 1, 3 or 5 parts by weight, relative to 1 part by weight of 2-amino-3-(4-bromobenzoyl)phenylacetic acid or its pharmacologically acceptable salt or a hydrate thereof.

The incorporation ratio of polyethylene glycol monostearate in the aqueous liquid preparation of the present invention is within a range of the minimum content of about 0.2 part by weight to the maximum content of about 0.5 or 1 part by weight, relative to 1 part by weight of 2-amino-3-(4-bro-

6

The preservative used in the present invention includes, for example, quaternary ammonium salts (e.g. benzalkonium chloride, benzethonium chloride, etc.), chlorhexidine gluconate, and the like, among which benzalkonium chloride is especially preferable.

Further, so long as the purpose of the present invention is achieved, conventional various additives such as isotonicity buffers, thickeners, stabilizers, chelating agents, pH controlling agents, perfumes and the like may be appropriately added to the aqueous liquid preparation of the present invention. The isotonicity include sodium chloride, potassium chloride, glycerine, mannitol, sorbitol, boric acid, glucose, propylene glycol and the like. The buffers include, for example, phosphate buffer, borate buffer, citrate buffer, tartarate buffer, acetate buffer, boric acid, borax, amino acids, and the like. The thickeners include polyvinylpyrrolidone, carboxymethylcellulose, carboxypropylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinyl alcohol, sodium polyacrylate, and the like. The stabilizers include sulfites such as sodium sulfite and the like. The chelating agents include sodium edetate, sodium citrate, condensed sodium phosphate and the like. The pH controlling agents include hydrochloric acid, sodium hydroxide, phosphoric acid, acetic acid and the like. The perfumes include 1-menthol, borneol, camphor, Eucalyptus oil, and the like.

With respect to the concentrations of the above various additives in the aqueous liquid preparation of the present invention,

the isotonic is incorporated into an osmotic pressure ratio of about 0.8 to 1.2, and the concentrations of the buffer and the thickener to be added are about 0.01 to 2 w/v and 0.1 to 10 w/v %, respectively.

The pH of the aqueous liquid preparation of the present invention is adjusted to about 6 to 9, preferably about 7 to 9, especially about 7.5 to 8.5.

So long as the purpose of the present invention is achieved, other same or different kind of active ingredients may be appropriately added.

The aqueous liquid preparation of the present invention can be prepared by per se known method or according to the method as described in the Japanese Pharmacopoeia. 14th Edition, General Rules for Preparations, Solutions or Ophthalmic solutions.

The aqueous liquid preparation of the present invention can be applied to warm-blooded animals such as human, rat, mouse, rabbit, cow, pig, dog, cat, and the like.

The aqueous liquid preparation of the present invention can be prepared easily by dissolving the above-mentioned components in, for example, distilled water or sterile purified water. For example, the aqueous liquid preparation in the form of an eye drop can be used for the treatment of inflammatory diseases in anterior or posterior segment of the eye such as blepharitis, conjunctivitis, scleritis, postoperative inflammation, and the like. The dose of the aqueous liquid preparation containing 0.1 w/v of sodium 2-amino-3-(4-bromobenzoyl)phenylacetate hydrate is, for example, administered to an adult 3 to 6 times daily in an amount of 1 to 2 drops per one time. Depending on the degree of diseases, frequency of dosing is appropriately controlled.

BEST MODE FOR CARRYING OUT THE INVENTION

The present invention is illustrated by way of the following

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.