

PROLENSA (bromfenac ophthalmic solution) 0.07%

PROLENSA (bromfenac ophthalmic solution) 0.07% is a nonsteroidal anti-inflammatory drug (NSAID) indicated for the treatment of postoperative inflammation and reduction of ocular pain in patients who have undergone cataract surgery.

(/our-products/rx-pharmaceutical/prolelsa-bromfenac-ophthalmic-solution-007-copy2/about-prolelsa)

INDICATION FOR PROLENSA (bromfenac ophthalmic solution) 0.07%

PROLENSA (bromfenac ophthalmic solution) 0.07% is a nonsteroidal anti-inflammatory drug (NSAID) indicated for the treatment of postoperative inflammation and reduction of ocular pain in patients who have undergone cataract surgery.

IMPORTANT SAFETY INFORMATION ABOUT PROLENSA

- PROLENSA contains sodium sulfite, a sulfite that may cause allergic type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people.
- All topical nonsteroidal anti-inflammatory drugs (NSAIDs), including bromfenac, may slow or delay healing. Concomitant use of topical NSAIDs and topical steroids may increase the potential for healing problems.
- There is the potential for cross-sensitivity to acetylsalicylic acid, phenylacetic acid derivatives, and other NSAIDs, including bromfenac. Use with caution in patients who have previously exhibited sensitivities to these drugs.
- There have been reports that ocularly applied NSAIDs may cause increased bleeding of ocular tissues (including hyphemas) in conjunction with ocular surgery. Use with caution in patients with known bleeding tendencies or who are receiving other medications which may prolong bleeding time.
- Use of topical NSAIDs may result in keratitis. Patients with evidence of corneal epithelial breakdown should immediately discontinue use of topical NSAIDs.

INDICATION FOR LOTEMAX GEL (loteprednol etabonate ophthalmic gel) 0.5%

LOTEMAX GEL (loteprednol etabonate ophthalmic gel) 0.5% is indicated for the treatment of post-operative inflammation and pain following ocular surgery.

IMPORTANT SAFETY INFORMATION ABOUT LOTEMAX GEL

- LOTEMAX GEL is contraindicated in most viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella, and also in mycobacterial infection of the eye and fungal diseases of ocular structures.
- Prolonged use of corticosteroids may result in glaucoma with damage to the optic nerve, defects in visual acuity and fields of vision. If this product is used for 10 days or longer, IOP should be monitored.
- Use of corticosteroids may result in posterior subcapsular cataract formation.
- Use of steroids after cataract surgery may delay healing and increase the incidence of bleb formation and occurrence of perforations in those with diseases causing corneal and scleral thinning. The initial prescription and renewal of the medication order should be made by a physician only after examination of the patient with the aid of magnification, and where appropriate, fluorescein staining.
- Prolonged use of corticosteroids may suppress the host response and thus increase the hazard of secondary ocular infection. In acute purulent conditions, steroids may mask infection or enhance existing infections.
- Use of corticosteroid medication in the treatment of patients with a history of herpes simplex requires great caution. Use of ocular steroids may prolong the course and exacerbate the severity of herpetic infections of the eye.

corneal epithelial defects, diabetes mellitus, ocular surface diseases (e.g., dry eye syndrome), rheumatoid arthritis, or repeat ocular surgeries within a short period of time may be at increased risk for corneal adverse events which may become sight threatening. Topical NSAIDs should be used with caution in these patients. Post-marketing experience with topical NSAIDs suggests that use more than 24 hours prior to surgery or use beyond 14 days post-surgery may increase patient risk for the occurrence and severity of corneal adverse events.

- *PROLENSA* should not be instilled while wearing contact lenses. The preservative in *PROLENSA*, benzalkonium chloride, may be absorbed by soft contact lenses. Lenses may be reinserted after 10 minutes following administration of *PROLENSA*.
- The most commonly reported adverse reactions in 3%-8% of patients were anterior chamber inflammation, foreign body sensation, eye pain, photophobia, and blurred vision.

Please [click here](#) ([/Portals/109/-/m/BL/United%20States/Files/Package%20Inserts/Pharma/prolensa-insert.pdf](#)) for Prescribing Information about *PROLENSA*.

- Fungal infections of the cornea are particularly prone to develop coincidentally with long-term local steroid application. Fungus invasion must be considered in any persistent corneal ulceration where a steroid has been used or is in use.
- Patients should not wear contact lenses when using *LOTEMAX GEL*.
- The most common ocular adverse drug reactions were anterior chamber inflammation (5%), eye pain (2%) and foreign body sensation (2%).

Please [click here](#) ([/Portals/109/-/m/BL/United%20States/Files/Package%20Inserts/Pharma/lotemaxg-package-insert.pdf](#)) for Prescribing Information about *LOTEMAX GEL*.

US/PRA/15/0041(1)

Professional Resources

Meetings and Exhibits ([/ecp/for-your-practice/professional-calendar/meetings-and-exhibits/](#))
 Educational Meetings/Dinners ([/ecp/for-your-practice/professional-calendar/education/](#))
 Resource Materials ([/ecp/for-your-practice/resource-materials/](#))
 Training & Tools ([/ecp/for-your-practice/training/](#))

Products

Online Ordering ([/ecp/online-ordering/online-ordering-overview/](#))
 Contact Lenses ([/ecp/our-products/contact-lenses](#))
 Contact Lens Care ([/ecp/our-products/contact-lens-care/](#))
 OTC Ophthalmics ([/ecp/our-products/otc-ophthalmics/](#))
 Eye Vitamins ([/ecp/our-products/eye-vitamins](#))
 Newsroom ([/our-company/news](#))
 Careers ([/our-company/careers](#))
 Investor Relations ([/our-company/investor-relations/](#))
 Grants & Contributions ([/our-company/grants-and-charitable-contributions/grants-and-charitable-contributions-landing](#))

Customer Support

Worldwide Locations ([/our-company/worldwide-locations](#))
 Customer policies and forms ([/reference/customer-policies-and-forms/](#))

Legal

Legal Notice ([/reference/legal-notice/](#))
 Privacy Policy ([/reference/privacy-policy/](#))
 Safety Data Sheets (SDS) ([/ecp/for-your-practice/resource-materials/material-safety-data-sheets](#))

© 2016 Bausch & Lomb Incorporated.

All information and materials on this site pertain to the U.S. only, unless otherwise indicated.