

THE MERCK INDEX

AN ENCYCLOPEDIA OF
CHEMICALS, DRUGS, AND BIOLOGICALS

THIRTEENTH EDITION

Editorial Staff

Maryadele J. O'Neil, *Senior Editor*

Ann Smith, *Senior Associate Editor*

Patricia E. Heckelman, *Associate Editor*

John R. Obenchain Jr., *Editorial Assistant*

Jo Ann R. Gallipeau, *Technical Assistant*

Mary Ann D'Arecca, *Administrative Associate*

Susan Budavari, *Editor Emeritus*

Published by
Merck Research Laboratories
Division of

MERCK & CO., INC.

Whitehouse Station, NJ

IPR2015-01099
IPR2015-01097
IPR2015-01100
IPR2015-01105

2001

Lupin EX1096
Page 1

MERCK & CO., INC.
Whitehouse Station, NJ
USA

1st Edition—1889
2nd Edition—1896
3rd Edition—1907
4th Edition—1930
5th Edition—1940
6th Edition—1952
7th Edition—1960
8th Edition—1968
9th Edition—1976
10th Edition—1983
11th Edition—1989
12th Edition—1996

Library of Congress Catalog
Card Number 89-60001
ISBN Number 0911910-13-1

Copyright © 2001 by MERCK & CO., INC.
All rights reserved. Copyright under the Universal Copyright Convention
and the International Copyright Convention.
Copyright reserved under the Pan-American Copyright Convention.

Printed in the USA

Page 2


er, alcohol, ether. LD₅₀ i.p. in female
crystals from ethanol-water, mp 197-

ingestion of *C. ruscifolia* has been
presence of tutin. Symptoms include
irium, convulsions.

[14-5] Tricyclo[4.4.0.0^{3,8}]decane.
C 88.16%, H 11.84%. Synthesis:
oc. **84**, 3412 (1962); Gautier, Des-
45, 297 (1967); Whitlock, Siefken,
9 (1968). Absolute configuration:
Letters **1968**, 5467.




[268-36-4] Butylcarbamic acid
[methyl]-2-methylpentyl ester; carba-
[methyl]-2-methylpentyl ester of butylcar-
[methyl]-2-propyl-1,3-propanediol dicar-
[methyl]ethylene butylcarbamate carba-
[methyl]trian. C₁₃H₂₆N₂O₄; mol wt 274.36.
%, O 23.33%. Prep'd from 2-meth-
[methyl]carbamate + butyl isocyanate:
9 (1960 to Carter Prod.). Compre-
[methyl]sberg et al., *Anal. Profles Drug*



roethane + hexane (1:2), mp 49-

[6879-02-3] 9,11,12,13,13a,14-
oxydibenzo[*f,h*]pyrrolo[1,2-*b*]-
oxyphenanthro[9,10:6',7']indol-
393.47. C 73.26%, H 6.92%, N
indolizidine alkaloid; isomeric
of *l*-isomer from *Tylophora cre-*
daceae: E. Gellert et al., *J. Chem.*
et al., *J. Pharm. Sci.* **59**, 1501
: E. Gellert, R. Rudzats, *J. Med.*
mer isolated from *Ficus septica*,
dl-form and structure: E. Gellert
Gellert, *Austr. J. Chem.* **23**, 2503
is: G. R. Donaldson et al., *Biochem.*
104 (1968); E. Battaner, D. Vas-
254, 316 (1971). Mode of action
grollman, *Mol. Pharmacol.* **8**, 538
Antibiotics (N.Y.) **6**, 47 (1983).




dl-Form. Needles from chloroform + methanol, mp 219-221°.

l-Form. Crystals from methanol, dec 218-220°. uv max: 263, 342, 360 nm (log ε 4.81, 3.25, 3.09). [α]_D²⁵ -45° (c = 0.74 in chloroform). pKa (50% aq ethanol): 6.7.

d-Form. Crystals from methanol, mp 220-222°. [α]_D²² +20.5°.

9899. Tylophorine. [482-20-2] (13aS)-9,11,12,13,13a,14-Hexahydro-2,3,6,7-tetramethoxydibenzo[*f,h*]pyrrolo[1,2-*b*]isoquinoline; 2,3,6,7-tetramethoxyphenanthro[9,10:6',7']indolizidine. C₂₄H₂₇NO₄; mol wt 393.47. C 73.26%, H 6.92%, N 3.56%, O 16.26%. Major alkaloid from *Tylophora asthmatica* Wight et Arn., *Asclepiadaceae*. Also found in other *Asclepiadaceae*, *Moraceae*, *Urticaceae* and *Lauraceae*. Naturally occurring form originally isolated and reported to be levorotatory; later corrected to dextrorotatory. Isoln: A. N. Ratnagiriswaran, K. Venkatachalam, *Indian J. Med. Res.* **22**, 433 (1935); R. N. Chopra et al., *Arch. Pharm.* **275**, 236 (1937); T. R. Govindachari et al., *J. Chem. Soc.* **1954**, 2801. Structure: *eidem*, *Tetrahedron* **9**, 53 (1960). Absolute configuration: *eidem*, *J. Chem. Soc. Perkin Trans. I* **1974**, 1161. Synthesis of the *dl*-form: *eidem*, *Chem. & Ind. (London)* **1960**, 664; N. A. Khatri et al., *J. Am. Chem. Soc.* **103**, 6387 (1981). Synthesis and verification of dextrorotation of naturally occurring form: T. F. Buckley 3rd, H. Rapoport, *J. Org. Chem.* **48**, 4222 (1983); J. E. Nordlander, F. G. Njoroge, *ibid.* **52**, 1627 (1987). Stereoselective synthesis of *l*-form: M. Ihara et al., *Tetrahedron Letters* **29**, 4135 (1988). Biosynthesis: Mulchandani et al., *Phytochemistry* **8**, 1931 (1969); *ibid.* **10**, 1047 (1971); D. S. Bhakuni, V. K. Mangla, *Tetrahedron* **37**, 401 (1981). Pharmacology: C. Gopalakrishnan et al., *Indian J. Med. Res.* **69**, 513 (1979); **71**, 940 (1980).




Crystals, dec 282-284°. [α]_D²³ +15° (c = 0.7 in chloroform); [α]_D²⁵ +73° (c = 0.7 in chloroform). uv max in ethanol: 257, 286, 339, 356 nm (log ε 4.7, 4.42, 3.28, 3.19). Unstable in solutions, decomposition with yellowing sets in promptly accompanied by decreasing rotatory strength.

l-Form. Colorless crystals, dec 286-287°. [α]_D²⁷ -11.6° (c = 1.07 in chloroform). uv max 255, 290, 340, 352 nm (log ε 4.74, 4.49, 3.30, 2.93). Sol in chloroform; slightly sol in abs alc, ether, cold benzene. Practically insol in water.

dl-Form. Crystals from chloroform + ethanol, mp 292°.

9900. Tylosin. [1401-69-0] Tylan. C₄₆H₇₇NO₁₇; mol wt 916.10. C 60.31%, H 8.47%, N 1.53%, O 29.69%. Macrolide antibiotic isolated from a strain of *Streptomyces fradiae* found in soil from Thailand: Hamill et al., *Antibiot. & Chemother.* **11**, 328 (1961); *eidem*, *US 3178341* (1965 to Lilly). Prodn in batch and chemostat cultures: P. P. Gray, S. Bhuwathanapun, *Biotech. Bioeng.* **22**, 1785 (1980). Partial structure: Morin, Gorman, *Tetrahedron Letters* **1964**, 2339. Structure: Morin et al., *ibid.* **1970**, 4737; Achenbach et al., *Ber.* **108**, 2481 (1975). Configurational study: S. Omura et al., *Tetrahedron Letters* **1977**, 1045. Abs config: *eidem*, *J. Antibiot.* **33**, 915 (1980); N. D. Jones et al., *ibid.* **35**, 420 (1982). Synthesis of *tylonolide*, the aglycone: S. Masamune et al., *J. Am. Chem. Soc.* **98**, 7874 (1976); K. Tatsuta et al., *Tetrahedron Letters* **22**, 3997 (1981). Relationship of ribosomal binding and antibacterial properties: J. W. Corcoran et al., *J. Antibiot.* **30**, 1012 (1977). Biosynthesis

studies: E. T. Seno et al., *Antimicrob. Ag. Chemother.* **11**, 455 (1977); S. Omura et al., *J. Antibiot.* **31**, 254 (1978).





Crystals from water, mp 128-132°. [α]_D²⁵ -46° (c = 2 in methanol). uv max: 282 nm (E_{1cm}^{1%} 245). Soly in water at 25°: 5 mg/ml. Sol in lower alcohols, esters and ketones, in chlorinated hydrocarbons, benzene, ether. Solns are stable at pH 4-9; at pH <4 another active compd, *desmycosin* is formed.

Hydrochloride. C₄₆H₇₇NO₁₇·HCl. Crystals from ethanol + ether, mp 141-145°.

THERAP CAT (VET): Antibacterial.

9901. Tyloxapol. [25301-02-4] 4-(1,1,3,3-Tetramethylbutyl)phenol polymer with formaldehyde and oxirane; oxyethylated tertiary octylphenol formaldehyde polymer; *p*-isooctyl-polyoxyethylenephenol formaldehyde polymer; tyloxopal; Al-evaire; Superinone; Triton A-20; Triton WR-1339. Nonionic detergent with surface-tension-reducing properties. Prepn: Bock, Rainey, *US 2454541* (1948 to Rohm & Haas); J. W. Cornforth et al., *Nature* **168**, 150 (1951). Use to induce exptl hyperlipidemia: A. Kellner et al., *J. Exp. Med.* **93**, 373 (1951); P. E. Schurr et al., *Lipids* **7**, 68 (1972). Use as ophthalmic excipient: D. E. Guttman et al., *J. Pharm. Sci.* **50**, 305 (1961). Phase behavior of mixtures with water: K. Westesen, *Int. J. Pharm.* **102**, 91 (1994); K. Westesen, M. H. J. Koch, *ibid.* **103**, 225 (1994). Tissue distribution and excretion: R. L. DeAngelis et al., *Xenobiotica* **25**, 521 (1995).



d²⁰ 1.0963. Cloud point: 92-97°. Slowly but freely misc with water. Sol in benzene, toluene, chloroform, carbon tetrachloride, carbon disulfide, acetic acid. Alkaline pH. Oxidized by metals.

Combination with colfosceril palmitate and hexadecanol, see Exosurf[®]

USE: Pharmaceutic aid (dispersing agent; excipient). To induce exptl hyperlipidemia in animal models.

THERAP CAT: Mucolytic.

9902. Tymazoline. [24243-97-8] 4,5-Dihydro-2-[[5-methyl-2-(1-methylethyl)phenoxy]methyl]-1*H*-imidazole; 2-[[thymyloxy]methyl]-2-imidazoline; 2-(thymyloxymethyl)glyoxalidine; 2-[[*p*-mentha-1,3,5-trien-2-yloxy]methyl]-2-imidazoline. C₁₄H₂₀N₂O; mol wt 232.32. C 72.38%, H 8.68%, N 12.06%, O 6.89%. Prepd from an alkyl thymyloxyacetimidate and ethylenediamine: Sonn, *US 2149473* (1939 to Ciba); Djerassi,

Consult the Name Index before using this section.

Page 1751

Page 3