

Home > > Rx Pharmaceutical > PROLENSA (bromfenac ophthalmic solution) 0.07% > No Generic Equivalent For ProleNSA

PROLENSA (bromfenac ophthalmic solution) 0.07%

PROLENSA HOME

PROLENSA[®]
(bromfenac ophthalmic solution) 0.07%

?

Is There a Generic Substitute for PROLENSA?



Your eye doctor has prescribed specific medications for you after cataract surgery to help control postsurgical pain and inflammation in your eye(s).

Sometimes your pharmacy will ask if you want a generic substitution for branded medications. If you are prescribed *PROLENSA* after cataract surgery, it is important that you understand that there is no generic equivalent for this drug. If you are asked by your pharmacist to substitute your prescription, you should contact your eye doctor.

Why did my physician prescribe PROLENSA (bromfenac ophthalmic solution) 0.07%?

PROLENSA is a nonsteroidal anti-inflammatory drug (NSAID) indicated to treat inflammation and reduce eye pain in patients after cataract surgery. There is no generic available for *PROLENSA*.¹ If you have further questions about *PROLENSA*, talk to your eye doctor.

FDA-approved Brand

Product Attributes	PROLENSA ¹
Treats inflammation and pain after cataract surgery	✓
No generic equivalent	✓
Established safety profile	✓
No shaking required	✓
Consistent dose in each drop	✓

IPR2015-01099
IPR2015-01097
IPR2015-01100
IPR2015-01105

Lupin EX1152
Page 1

INDICATION FOR PROLENSA (bromfenac ophthalmic solution) 0.07%

PROLENSA (bromfenac ophthalmic solution) 0.07% is a nonsteroidal anti-inflammatory drug (NSAID) indicated to treat inflammation and reduce eye pain in patients after cataract surgery.

IMPORTANT SAFETY INFORMATION ABOUT PROLENSA

PROLENSA contains sodium sulfite, a substance that may cause allergic reactions, some severe, in certain people. In addition, tell your doctor if you have had an allergic reaction to any other medications.

Slow or delayed healing may occur while using non-steroidal anti-inflammatory drugs (NSAIDs) such as *PROLENSA*.

Use of NSAIDs in the eye may result in certain serious eye conditions that can be sight threatening. Tell your doctor immediately about any side effect that you may have.

Replace bottle cap after using and do not touch dropper tip to any surface in order to avoid contamination of the bottle.

Remove contact lenses prior to using *PROLENSA*. The preservative in *PROLENSA*, benzalkonium chloride, may be absorbed by soft contact lenses. Lenses may be reinserted after 10 minutes following administration of *PROLENSA*.

If more than one eye medicine is being used, use the eye medicines at least 5 minutes apart.

The most common side effects (seen in 3%-8% of patients) were inflammation of the eye, foreign body sensation, eye pain, light sensitivity, and blurred vision.

Vision and Eye Health

Vision and Age
Eye Infections & Irritations
Diseases and Disorders
Find a Doctor
[PDF](#) Eye Exam Check List

Products

Contact Lenses
Contact Lens Care
Dry Eye Products
Allergy/Redness Relief
Rx Pharmaceutical
Eye Vitamins
Surgical Products
Vision Accessories

Customer Support

Contact Us
Customer policies and forms
Rebate Center
Worldwide Locations

Legal

Legal Notice
Privacy Policy
Safety Data Sheets (SDS)

Newsroom

Careers
Investor Relations
Grants & Contributions

© 2016 Bausch & Lomb Incorporated.

All information and materials on this site pertain to the U.S. only, unless otherwise indicated.