

NONIONIC SURFACTANTS

EDITED BY

MARTIN J. SCHICK

LEVER BROTHERS CO.
RESEARCH CENTER
EDGEWATER, NEW JERSEY

1967

MARCEL DEKKER, INC., NEW YORK

IPR2015-01099

IPR2015-01097

Page 54474

COPYRIGHT © 1966 by MARCEL DEKKER, INC.

ALL RIGHTS RESERVED

No part of this book may be reproduced in any form, by photostat, microfilm, or any other means, without the written permission of the publisher.

MARCEL DEKKER, INC.
95 Madison Avenue, New York, New York 10016

LIBRARY OF CONGRESS CATALOG CARD NUMBER 66 22492

PRINTED IN THE UNITED STATES OF AMERICA

PERMISSIONS

Academic Press Inc., New York, N. Y.

- Fig. 14.11. M. J. Schick, *J. Colloid Sci.*, **17**, 801 (1962) (Fig. 1).
Fig. 14.13. *Ibid.*, **18**, 373 (1963) (Fig. 3).
Fig. 15.6. P. Becher, *ibid.*, **16**, 49 (1961) (Fig. 1).
Fig. 15.8. *Ibid.*, **17**, 325 (1962) (Fig. 2).
Fig. 17.1. K. Shinoda, T. Nakagawa, B. Tamamuchi, and T. Isemura, *Colloidal Surfactants*, 1963, p. 140 (Fig. 2.27).
Fig. 17.2. *Ibid.*, p. 141 (Fig. 2.28).
Fig. 17.3. M. E. L. McBain and F. Hutchinson, *Stabilization and Related Phenomena*, 1955, p. 169 (Fig. 5.14).
Fig. 17.4. B. A. Mulley and A. D. Metcalf, *J. Colloid Sci.*, **17**, 525 (1962) (Fig. 2).
Fig. 17.5. *Ibid.*, **17**, 526 (1962) (Fig. 3).
Fig. 17.7. K. Shinoda, T. Nakagawa, B. Tamamuchi, and T. Isemura, *Colloidal Surfactants*, 1963, p. 142 (Fig. 2.29).
Fig. 17.8. T. Nakagawa, K. Kuriyama, and H. Inoue, *J. Colloid Sci.*, **15**, 275 (1960) (Fig. 5).
Fig. 17.9. W. N. Maclay, *ibid.*, **11**, 281 (1956) (Fig. 6).
Fig. 17.10. M. J. Schick, *ibid.*, **17**, 812 (1962) (Fig. 7).
Fig. 17.11. W. N. Maclay, *ibid.*, **11**, 277 (1956) (Fig. 4).
Fig. 17.28. B. A. Mulley and A. D. Metcalf, *ibid.*, **19**, 503 (1964) (Fig. 6).
Fig. 17.29. *Ibid.*, **19**, 504 (1964) (Fig. 2).
Fig. 17.30. S. Ross, *ibid.*, **6**, 502-504 (1951) (Figs. 1-3).
Fig. 19.12. E. Maackor, *ibid.*, **6**, 492 (1951) (Fig. 1).
Fig. 19.14. E. Maackor and J. H. van der Waals, *ibid.*, **7**, 535 (1952) (Figs. 3 and 4).
Fig. 21.2. J. F. Danielli, K. G. A. Pankhurst, and A. C. Riddell, *Recent Progress in Surface Science*, 1964, p. 61 (Fig. 2).
Table 17.3. T. Nakagawa, K. Kuriyama, and H. Inoue, *J. Colloid Sci.*, **15**, 270 (1960) (Table II).
Table 17.9. A. M. Mankowich, *ibid.*, **14**, 133 (1959) (Table II).
Table 17.12. S. Ross, *ibid.*, **6**, 500 (1951) (Table II).
Table 21.1. A. G. Brown, W. C. Thuman, and J. W. McBain, *ibid.*, **8**, 502 (1953) (Table I).

CHAPTER 22

CONFIGURATION OF THE POLYOXYETHYLENE CHAIN IN BULK

M. Rösch

CHEMISCHE FABRIK STOCKHAUSEN & CIE.
KREFELD, GERMANY

| | |
|--|-----|
| 22.1. Introduction | 753 |
| 22.2. X-ray Investigations of Organic Chain-Molecules | 754 |
| 22.3. X-ray Investigations of Ionic Surfactants | 756 |
| A. Soaps | 756 |
| B. Miscellaneous Ionic Surfactants | 758 |
| 22.4. X-Ray Investigations of Nonionic Surfactants | 758 |
| A. Polyoxyethylenes (and Polyglycols) | 758 |
| B. Polypropylene Oxide | 763 |
| C. Polyoxyethylene Fatty Acid and Fatty Acid Glycerol Esters | 764 |
| D. Polyoxyethylene Alkylaryl Hydroxy Compounds | 764 |
| E. Polyoxyethylene Alkanol and Glycerol Ethers | 766 |
| 22.5. Speculative Considerations of the Fine-Structure of the Polyoxyethylene Chain and Macromolecular Lattice of Nonionic Surfactants | 774 |
| A. Chain Contraction in Polyoxyethylenes | 774 |
| B. Molecular Lattice of Nonionic Surfactants | 777 |
| 22.6. Investigation of Fine-Structure of Polyoxyethylene Chain by Alternate Methods | 782 |
| A. Electron Microscopy and Electron Diffraction | 782 |
| B. Melt Behavior | 783 |
| C. Crystallization Phenomena | 785 |
| 22.7. Conclusions | 791 |
| References | 792 |

22.1. INTRODUCTION

Studies of polyoxyethylene compounds in their bulk form have been the subject of several recent investigations for the elucidation of the configuration and fine-structure of the polyoxyethylene chain. Although it is not yet possible to present a definite picture of the structure of the polyoxyethylene chain based on these results, important conclusions and speculations can be made. The phenomenological part is covered first in Sections 22.2 to 22.4, followed by a speculative part in Section 22.5. Finally, in Section 22.7 an attempt is made to summarize in the light of our present knowledge the concepts of the configuration of the

polyoxyethylene chain in polyoxyethylene or its derivatives, such as nonionic surfactants, making allowance for some inherent uncertainties.

Our experimental knowledge of the "configuration of the polyoxyethylene chain in bulk" is mainly based on X-ray investigations. Therefore, in this chapter, results from X-ray investigations which have contributed so significantly to the general understanding of the configuration and fine-structure of organic chain-molecules are covered extensively in Sections 22.2 to 22.4. Besides the so-called short-distance- or wide angle-diffraction techniques, principally the low-angle-diffraction patterns have contributed to the characterization of the structure and dimensions of organic chain molecules. It is well-known that the crystal or crystallite structure as well as the unit cell dimensions in high-molecular-weight substances can be determined from both wide- and low-angle-diffraction patterns of the same object. Since the X-ray method has only recently been applied more extensively to investigations of nonionic surfactants, it appears desirable to review briefly in Section 22.3 our extensive knowledge from the preceding X-ray investigations of ionic surfactants.

The investigation of the fine structure of the polyoxyethylene chain by alternate experimental methods is covered in Section 22.6. Another object of this chapter, besides the specific discussion of nonionic surfactants, is also to review general concepts applicable to the fine-structure of paraffin chain compounds.

22.2. X-RAY INVESTIGATIONS OF ORGANIC CHAIN-MOLECULES

Early work has led to the generally accepted zigzag structure of paraffin-chain compounds, in which the centers of the C atoms lie in one plane and the distance between alternate CH_2 groups in the backbone chain amounts to 2.5 Å (1-3). This was followed by a complete structural analysis of the unit cell of paraffins (4-7). The paraffin chains are parallel to the c axis of the orthorhombic (odd number of C atoms) unit cell or of the monoclinic (even number of C atoms) unit cell with a cross-sectional area of 18.5 Å², lateral distances d_1 of 3.7 and 4.1 Å, and a distance of 3.5 Å between the CH_3 groups in the terminal positions of paraffin-chain layers.

Figure 22.1 presents the fine-structure of the paraffin chain. It is worth noting in connection with the subsequent considerations of polyoxyethylene derivatives that Mueller and Shearer in 1923 discussed various configurations of the paraffin chain, namely the zigzag, the screw, and a kind of meander structure (8). The schematic configuration of paraffin chain layers is illustrated in Figure 22.2, together with the reflections of the long period of paraffin chains.

Similarly, a lamellar crystal structure comparable to that of paraffins has been found for fatty acid esters and high-molecular-weight ketones (9). Although some analogy with paraffin chains exists, a difference in chain orientation was observed with the high-molecular-weight ketones, and especially with the fatty acids, and is illustrated in Figure 22.3. Dimers of carboxyl groups (linked by hydrogen bridges) connect the layers of paraffin chains, and the distance between the layer planes corresponds to two rather than one molecular length as determined from




Fig. 22.1. Fine structure of paraffin chain.




Fig. 22.2. Schematic arrangement of paraffin chains.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.