

[54] **GASTRIC ACID SECRETION INHIBITING SUBSTITUTED 2-(2-BENZIMIDAZOLYL)-PYRIDINES, PHARMACEUTICAL PREPARATIONS CONTAINING SAME, AND METHOD FOR INHIBITING GASTRIC ACID SECRETION**

[75] Inventors: **Ulf K. Junggren**, Mölnlycke; **Sven E. Sjöstrand**, Kungsbacka, both of Sweden

[73] Assignee: **Aktiebolaget Hassle**, Mölndal, Sweden

[21] Appl. No.: **27,277**

[22] Filed: **Apr. 5, 1979**

[30] **Foreign Application Priority Data**

Apr. 14, 1978 [SE] Sweden 7804231

[51] Int. Cl.³ **A61K 31/44; C07D 401/12**

[52] U.S. Cl. **424/263; 546/271; 546/301; 548/329**

[58] Field of Search **546/271; 424/263**

[56] **References Cited**

U.S. PATENT DOCUMENTS

4,045,563 8/1977 **Berntsson et al.** 546/271
 4,045,564 8/1977 **Berntsson et al.** 546/271
 4,182,766 1/1980 **Krasso et al.** 546/271 X

FOREIGN PATENT DOCUMENTS

1804450 5/1970 **Fed. Rep. of Germany** 546/271

OTHER PUBLICATIONS

Berntsson et al., Chem. Abst., 1976, vol. 85, No. 149139q.

Hideg et al., Chem. Abst., 1971, vol. 75, No. 98570g.


Primary Examiner—Henry R. Jiles

Assistant Examiner—Richard A. Schwartz

Attorney, Agent, or Firm—Brumbaugh, Graves, Donohue & Raymond

[57] **ABSTRACT**

The present invention relates to novel compounds of the formula



wherein R^1 and R^2 are the same or different and are each hydrogen, alkyl, halogen, carbomethoxy, carboethoxy, alkoxy, or alkanoyl, R^6 is hydrogen, methyl or ethyl, R^3 , R^4 and R^5 are the same or different and are each hydrogen, methyl, methoxy, ethoxy, methoxyethoxy or ethoxyethoxy whereby R^3 , R^4 and R^5 are not all hydrogen, and whereby when two of R^3 , R^4 and R^5 are hydrogen the third of R^3 , R^4 and R^5 is not methyl. The compounds are potent gastric acid secretion inhibitors.


29 Claims, No Drawings

**GASTRIC ACID SECRETION INHIBITING
SUBSTITUTED
2-(2-BENZIMIDAZOLYL)-PYRIDINES,
PHARMACEUTICAL PREPARATIONS
CONTAINING SAME, AND METHOD FOR
INHIBITING GASTRIC ACID SECRETION**

The present invention relates to new compounds having valuable properties in affecting gastric acid secretion in mammals, including man, as well as the process for their preparation, method of affecting gastric acid secretion and pharmaceutical preparations containing said novel compounds.

The object of the present invention is to obtain compounds which affect gastric acid secretion, and which inhibit exogenously or endogenously stimulated gastric acid secretion. These compounds can be used in the treatment of peptic ulcer disease.


It is previously known that compounds of the formulas I and II



wherein R^1 and R^2 are each selected from the group consisting of hydrogen, alkyl, halogen, cyano, carboxyl, carboxyalkyl, carboalkoxy, carboalkoxyalkyl, carbamoyl, carbamoyloxy, hydroxy, alkoxy, hydroxyalkyl, trifluoromethyl and acyl in any position, R^3 is selected from the group consisting of hydrogen, alkyl, acyl, carboalkoxy, carbamoyl, alkylcarbamoyl, dialkylcarbamoyl, alkylcarbonylmethyl, alkoxy carbonylmethyl, and alkylsulphonyl, and R^4 is selected from the group consisting of straight and branched alkylene groups having 1 to 4 carbon atoms, whereby at most one methylene group is present between S and the pyridyl group, and whereby the pyridyl group may be further substituted with alkyl or halogen, possess an inhibiting effect on gastric acid secretion.

It has now, however, surprisingly been found that the compounds defined below possess a still greater inhibiting effect than those given above.

Compounds of the invention are those of the general formula III



wherein R^1 and R^2 are same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, carbethoxy, alkoxy, and alkanoyl, R^6 is selected from the group consisting of hydrogen, methyl, and ethyl, and R^3 , and R^5 are the same or different and are each selected from the group consisting of hydrogen, methyl, methoxy, ethoxy, methoxyethoxy and ethoxyethoxy and R^4 is methoxy, ethoxy, methoxyethoxy or ethoxyethoxy whereby R^3 , R^4 , and R^5 are not all hydrogen, and whereby when two of R^3 , R^4 , and R^5 are hydrogen, the third of R^3 , R^4 and R^5 is not methyl.

Alkyl R^1 and R^2 of formula III are suitably alkyl having up to 7 carbon atoms, preferably up to 4 carbon atoms. Thus, alkyl R may be methyl, ethyl, n-propyl, isopropyl, n-butyl or isobutyl.

Halogen R^1 and R^2 are chloro, bromo, fluoro, or iodo.

Alkoxy R^1 and R^2 are suitably alkoxy groups having up to 5 carbon atoms, preferably up to 3 carbon atoms, such as methoxy, ethoxy, n-propoxy, or isopropoxy.

Alkanoyl R^1 and R^2 have preferably up to 4 carbon atoms and are e.g. formyl, acetyl, or propionyl, preferably acetyl.

A preferred group of compounds of the general formula III are those wherein R^1 and R^2 are the same or different and are each selected from the group consisting of hydrogen, alkyl, carbomethoxy, alkoxy, and alkanoyl, whereby R^1 and R^2 are not both hydrogen, R^6 is hydrogen, and R^3 , R^4 , and R^5 are the same or different and are each selected from the group consisting of hydrogen, methyl, methoxy, and ethoxy, whereby R^3 , R^4 , and R^5 are not all hydrogen, and whereby when two of R^3 , R^4 , and R^5 are hydrogen the third of R^3 , R^4 , and R^5 is not methyl.

A second preferred group of compounds of the general formula III are those wherein R^1 and R^2 are the same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, carbethoxy, alkoxy, and alkanoyl, R^6 is selected from the group consisting of hydrogen, methyl, and ethyl, R^3 is methyl, R^4 is methoxy, and R^5 is methyl.

A third preferred group of compounds of the general formula III are those wherein R^1 and R^2 are the same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, carbethoxy, alkoxy and alkanoyl, R^6 is selected from the group consisting of hydrogen, methyl and ethyl, and R^3 is hydrogen, R^4 is methoxy and R^5 is methyl or R^3 is methyl, R^4 is methoxy and R^5 is hydrogen.

A fourth preferred group of compounds of the general formula III are those wherein R^1 and R^2 are the same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, carbethoxy, alkoxy, and alkanoyl, R^6 is selected from the group consisting of hydrogen, methyl and ethyl, R^3 and R^5 are hydrogen and R^4 is methoxy.

A fifth preferred group of compounds of the general formula III are those wherein R^1 and R^2 are the same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, carbethoxy, alkoxy, and alkanoyl, R^6 is selected from the group consisting of hydrogen, methyl and ethyl, and R^3 and R^5 are methyl and R^4 is hydrogen.

A sixth preferred group of compounds of the general formula III are those wherein R^1 and R^2 are the same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, carbe-

3

thoxy, alkoxy, and alkanoyl, R⁶ is selected from the group consisting of hydrogen, methyl and ethyl, R³ and R⁵ are hydrogen and R⁴ is ethoxy, methoxyethoxy or ethoxyethoxy.


A seventh preferred group of compounds of the general formula III are those wherein R¹ and R² are the same or different and are each selected from the group consisting of hydrogen, alkyl, halogen, carbomethoxy, alkoxy, and alkanoyl, R⁶ is selected from the group consisting of hydrogen, methyl, and ethyl, R³, R⁴, and R⁵ are all methyl.

Other groups of compounds which may be used are:

Compounds in which R¹ is hydrogen, chloro, methyl, ethyl, methoxy, acetyl, carboethoxy or carbomethoxy; R² is hydrogen or methyl; R⁶ is hydrogen, methyl or ethyl; R³ and R⁵ are methyl; and R⁴ is methoxy. Compounds in which R¹ is hydrogen, chloro, methyl, ethyl, methoxy, acetyl, carboethoxy or carbomethyl; R² is hydrogen, methyl or ethyl; R⁴ is methoxy; and R³ is methyl and R⁵ is hydrogen or R³ is hydrogen and R⁵ is methyl.


Compounds of formula III above may be prepared according to the following methods:

(a) oxidizing a compound of formula IV




wherein R¹, R², R⁶, R³, R⁴, and R⁵ have the meanings given, to the formation of a compound of formula III.

(b) reacting a compound of the formula V




wherein R¹, R², and R⁶ have the meanings given above and M is a metal selected from the group consisting of K, Na and Li, with a compound of formula VI.




wherein R³, R⁴, and R⁵ have the same meanings as given above, Z is a reactive esterified hydroxy group, to the formation of a compound of formula III;

(c) reacting a compound of the formula VII

4




wherein R¹, and R² have the same meanings as given above and Z¹ is SH or a reactive esterified hydroxy group, with a compound of the formula VIII




wherein R⁶, R³, R⁴, and R⁵ have the same meanings as given above, and Z² is a reactive esterified hydroxy group or SH, to the formation of an intermediate of formula IV above, which then is oxidized to give a compound of formula III;

(d) reacting a compound of the formula IX



wherein R¹ and R² have the same meanings as given above with a compound of the formula X



wherein R⁶, R³, R⁴, and R⁵ have the same meanings as given above, to the formation of an intermediate of formula IV above, which then is oxidized to give a compound of formula III, which compound may be converted to its therapeutically acceptable salts, if so desired.

In the reactions above, Z, Z¹, and Z² may be a reactive, esterified hydroxy group which is a hydroxy group esterified with strong, inorganic or organic acid, preferably a hydrohalogen acid, such as hydrochloric acid, hydrobromic acid, or hydroiodic acid, also sulfuric acid or a strong organic sulfonic acid as a strong aromatic acid, e.g. benzenesulfonic acid, 4-bromobenzenesulfonic acid or 4-toluenesulfonic acid.

The oxidation of the sulfur atom in the chains above to sulfinyl (S→O) takes place in the presence of an oxidizing agent selected from the group consisting of nitric acid, hydrogen peroxide, peracids, peresters, ozone, dinitrogen tetraoxide, iodosobenzene, N-halosuccinimide, 1-chlorobenzotriazole, t-butylhypochlorite, diazobicyclo-[2,2,2]-octane bromine complex, sodium metaperiodate, selenium dioxide, manganese dioxide, chromic acid, ceric ammonium nitrate, bromine, chlo-

rine, and sulfuryl chloride. The oxidation usually takes place in a solvent wherein the oxidizing agent is present in some excess in relation to the product to be oxidized.

Depending on the process conditions and the starting materials, the end product is obtained either as the free base or in the acid addition salt, both of which are included within the scope of the invention. Thus, basic, neutral or mixed salts may be obtained as well as hemi, mono, sesqui or polyhydrates. The acid addition salts of the new compounds may in a manner known per se be transformed into free base using basic agents such as alkali or by ion exchange. On the other hand, the free bases obtained may form salts with organic or inorganic acids. In the preparation of acid addition salts preferably such acids are used which form suitable therapeutically acceptable salts. Such acids include hydrohalogen acids, sulfonic, phosphoric, nitric, and perchloric acids; aliphatic, alicyclic, aromatic, heterocyclic carboxy or sulfonic acids, such as formic, acetic, propionic, succinic, glycolic, lactic, malic, tartaric, citric, ascorbic, maleic, hydroxymaleic, pyruvic, phenylacetic, benzoic, p-aminobenzoic, antranilic, p-hydroxybenzoic, salicylic or p-aminosalicylic acid, embonic, methanesulfonic, ethanesulfonic, hydroxyethanesulfonic, ethylenesulfonic, halogenbenzenesulfonic, toluenesulfonic, naphthylsulfonic or sulfanilic acids; methionine, tryptophane, lysine or arginine.

These or other salts of the new compounds, as e.g. picrates, may serve as purifying agents of the free bases obtained. Salts of the bases may be formed, separated from solution, and then the free base can be recovered from a new salt solution in a purer state. Because of the relationship between the new compounds in free base form and their salts, it will be understood that the corresponding salts are included within the scope of the invention.

Some of the new compounds may, depending on the choice of starting materials and process, be present as optical isomers or racemate, or if they contain at least two asymmetric carbon atoms, be present as an isomer mixture (racemate mixture).

The isomer mixtures (racemate mixtures) obtained may be separated into two stereoisomeric (diastereomeric) pure racemates by means of chromatography or fractional crystallization.

The racemates obtained can be separated according to known methods, e.g. recrystallization from an optically active solvent, use of microorganisms, reactions with optically active acids forming salts which can be separated, separation based on different solubilities of the diastereomers. Suitable optically active acids are the L- and D-forms of tartaric acid, di-o-tolyl-tartaric acid, malic acid, mandelic acid, camphorsulfonic acid or quinic acid. Preferably the more active part of the two antipodes is isolated.

The starting materials are known or may, if they should be new, be obtained according to processes known per se.

In clinical use the compounds of the invention are administered orally, rectally or by injection in the form of a pharmaceutical preparation which contains an active component either as a free base or as a pharmaceutically acceptable, non-toxic acid addition salt, such as hydrochloride, lactate, acetate, sulfamate, in combination with a pharmaceutically acceptable carrier. The carrier may be in the form of a solid, semisolid or liquid diluent, or a capsule. These pharmaceutical preparations are a further object of the invention. Usually the

amount of active compound is between 0.1 to 95% by weight of the preparation, between 0.5 to 20% by weight in preparations for injection and between 2 and 50% by weight in preparations for oral administration.

In the preparation of pharmaceutical preparations containing a compound of the present invention in the form of dosage units for oral administration the compound selected may be mixed with a solid, pulverulent carrier, such as lactose, saccharose, sorbitol, mannitol, starch, amylopectin, cellulose derivatives or gelatin, as well as with an antifriction agent such as magnesium stearate, calcium stearate, and polyethyleneglycol waxes. The mixture is then pressed into tablets. If coated tablets are desired, the above prepared core may be coated with a concentrated solution of sugar, which may contain gum arabic, gelatin, talc, titanium dioxide or with a lacquer dissolved in volatile organic solvent or mixture of solvents. To this coating various dyes may be added in order to distinguish among tablets with different active compounds or with different amounts of the active compound present.

Soft gelatin capsules may be prepared which capsules contain a mixture of the active compound or compounds of the invention and vegetable oil. Hard gelatin capsules may contain granules of the active compound in combination with a solid, pulverulent carrier as lactose, saccharose, sorbitol, mannitol, potato starch, corn starch, amylopectin, cellulose derivatives or gelatin.

Dosage units for rectal administration may be prepared in the form of suppositories which contain the active substance in a mixture with a neutral fat base, or they may be prepared in the form of gelatin-rectal capsules which contain the active substance in a mixture with a vegetable oil or paraffin oil.

Liquid preparations for oral administration may be prepared in the form of syrups or suspensions, e.g. solutions containing from 0.2% to 20% by weight of the active ingredient and the remainder consisting of sugar and a mixture of ethanol, water, glycerol and propylene glycol. If desired, such liquid preparations may contain colouring agents, flavouring agents, saccharin and carboxymethylcellulose as a thickening agent.

Solutions for parenteral administration by injection may be prepared as an aqueous solution of a watersoluble pharmaceutically acceptable salt of the active compound, preferably in a concentration from 0.5% to 10% by weight. These solutions may also contain stabilizing agents and/or buffering agents and may be manufactured in different dosage unit ampoules.

Pharmaceutical tablets for oral use are prepared in the following manner: The solid substances are ground or sieved to a certain particle size, and the binding agent is homogenized and suspended in a suitable solvent. The therapeutically active compounds and auxiliary agents are mixed with the binding agent solution. The resulting mixture is moistened to form a uniform suspension having the consistency of wet snow. The moistening causes the particles to aggregate slightly, and the resulting mass is pressed through a stainless steel sieve having a mesh size of approximately 1 mm. The layers of the mixture are dried in carefully controlled drying cabinets for approximately ten hours to obtain the desired particle size and consistency. The granules of the dried mixture are sieved to remove any powder. To this mixture, disintegrating, antifriction and antiadhesive agents are added. Finally, the mixture is pressed into tablets using a machine with the appropriate punches and dies to obtain the desired tablet size. The pressure applied af-

fects the size of the tablet, its strength and its ability to dissolve in water. The compression pressure used should be in the range 0.5 to 5 tons. Tablets are manufactured at the rate of 20,000 to 200,000 per hour. The tablets, especially those which are rough or bitter, may be coated with a layer of sugar or some other palatable substance. They are then packaged by machines having electronic counting devices. The different types of packages consist of glass or plastic gallipots, boxes, tubes and specific dosage adapted packages.

The typical daily dose of the active substance varies according to the individual needs and the manner of administration. In general, oral dosages range from 100 to 400 mg/day of active substance and intravenous dosages range from 5 to 20 mg/day.

The following illustrates a preferred embodiment of the invention without being limited thereto. Temperature is given in degrees Centigrade.

The starting materials in the examples found below were prepared in accordance with the following methods:

(1) a 1,2-diamino compound, such as o-phenylenediamine was reacted with potassium ethylxanthate (according to *Org. Synth.* Vol. 30, p. 56) to form a 2-mercaptobenzimidazole; (2) the compound 2-chloromethylpyridine was prepared by reacting 2-hydroxymethylpyridine with thionylchloride (according to *Arch. Pharm.* Vol. 26, pp. 448-451 (1956)); (3) the compound 2-chloromethylbenzimidazole was prepared by condensing o-phenylenediamine with chloroacetic acid.

EXAMPLE 1

28.9 g of 2-[2-(4,5-dimethyl)pyridylmethylthio]-(5-acetyl-6-methyl)-benzimidazole were dissolved in 160 ml of CHCl_3 , 24.4 g of m-chloroperbenzoic acid were added in portions while stirring and cooling to 5° C. After 10 minutes, the precipitated m-chlorobenzoic acid was filtered off. The filtrate was diluted with CH_2Cl_2 , washed with Na_2CO_3 solution, dried over Na_2SO_4 and evaporated in vacuo. The residue crystallized when diluted with CH_3CN , and 2-[2-(4,5-dimethyl)pyridylmethylsulfanyl]-(5-acetyl-6-methyl)benzimidazole was recrystallized from CH_3CN . Yield 22.3 g; m.p. 158° C.

EXAMPLES 2-30

The preparation of compounds of formula III labelled 2-26 was carried out in accordance with Example 1 above. The compounds prepared are listed in Table 1 which identifies the substituents for these compounds.

EXAMPLE 31 (METHOD C)

0.1 moles of 4,6-dimethyl-2-mercaptobenzimidazole were dissolved in 20 ml of water and 200 ml of ethanol containing 0.2 moles of sodium hydroxide. 0.1 moles of 2-chloromethyl-(3,5-dimethyl)pyridine hydrochloride

were added and the mixture was refluxed for two hours. The sodium chloride formed was filtered off and the solution was evaporated in vacuo. The residue was dissolved in acetone and was treated with active carbon. An equivalent amount of concentrated hydrochloric acid was added, whereupon the mono-hydrochloride of 2-[2-(3,5-dimethyl)pyridylmethylthio]-4,6-dimethylbenzimidazole was isolated. Yield 0.05 moles.

This compound was then oxidized in accordance with Example 1 above to give the corresponding sulfinyl compound, melting point 50°-55° C.

EXAMPLE 32 (METHOD B)

0.1 moles of 2-[Li-methylsulfanyl](5-acetyl-6-methyl)-benzimidazole were dissolved in 150 mls of benzene. 0.1 moles 2-chloro-(3,5-dimethyl)pyridine were added and the mixture was refluxed for two hours. The lithium-chloride formed was filtered off, and the solution was evaporated in vacuo. The residue was crystallized from CH_3CN , and recrystallized from the same solvent. Yield 0.82 moles of 2-[2-(3,5-dimethyl)pyridylmethylsulfanyl]-(5-acetyl-6-methyl)benzimidazole melting at 171° C.

EXAMPLE 33 (METHOD D)

23.4 g of 2-[2-(3,4,5-trimethyl)pyridylmethylthio]formic acid and 16.6 g of o-(5-acetyl-6-methyl)-phenylenediamine were boiled for 40 minutes in 100 ml of 4 N HCl. The mixture was cooled and neutralized with ammonia. The neutral solution was then extracted with ethyl acetate. The organic phase was treated with active carbon and evaporated in vacuo. The residue was dissolved in acetone whereupon an equivalent of concentrated HCl was added. The precipitated hydrochloride was filtered off after cooling and the salt was recrystallized from absolute ethanol and some ether. Yield of 2-[2-(3,4,5-trimethylpyridyl)methylthio]-(5-acetyl-6-methyl)benzimidazole was 6.5 g.

This compound was then oxidized in accordance with Example 1 above, to give the corresponding sulfinyl derivative. M.p. 190° C.

EXAMPLE 34 (METHOD C)

22.0 g of 2-mercapto-(5-acetyl-6-methyl)benzimidazole and 19.5 g of chloromethyl(4,5-dimethyl)pyridine hydrochloride were dissolved in 200 ml of 95% ethanol. 8 g of sodium hydroxide in 20 ml of water were added, whereupon the solution was refluxed for two hours. The sodium chloride formed was filtered off and the solution was evaporated in vacuo. The residue, 2-[2-(4,5-dimethyl)pyridylmethylthio]-(5-acetyl-6-methyl)benzimidazole, was recrystallized from 70% ethanol. Yield 10.6 g.

This compound was then oxidized in accordance with Example 1 above, to give the corresponding sulfinyl derivative. M.p. 158° C.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.