

# Plasma Physics and Engineering

second edition

Alexander Fridman

Drexel University, Philadelphia, USA

Lawrence A. Kennedy

University of Illinois at Chicago, USA



CRC Press

Taylor & Francis Group

Boca Raton London New York

CRC Press is an imprint of the  
Taylor & Francis Group, an **informa** business

CRC Press  
Taylor & Francis Group  
6000 Broken Sound Parkway NW, Suite 300  
Boca Raton, FL 33487-2742

© 2011 by Taylor and Francis Group, LLC  
CRC Press is an imprint of Taylor & Francis Group, an Informa business

No claim to original U.S. Government works

Printed in the United States of America on acid-free paper  
10 9 8 7 6 5 4 3 2 1

International Standard Book Number-13: 978-1-4398-1229-7 (Ebook-PDF)

This book contains information obtained from authentic and highly regarded sources. Reasonable efforts have been made to publish reliable data and information, but the author and publisher cannot assume responsibility for the validity of all materials or the consequences of their use. The authors and publishers have attempted to trace the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission to publish in this form has not been obtained. If any copyright material has not been acknowledged please write and let us know so we may rectify in any future reprint.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access [www.copyright.com](http://www.copyright.com) (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

**Trademark Notice:** Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Visit the Taylor & Francis Web site at  
<http://www.taylorandfrancis.com>

and the CRC Press Web site at



Find authenticated court documents without watermarks at [docketalarm.com](http://docketalarm.com).

be derived taking into account the ion motion

$$\frac{k^2 c^2}{\omega^2} = 1 - \frac{\omega_p^2}{\omega^2} \frac{1}{\left(1 \pm \frac{\omega_B}{\omega} - \frac{\omega_B \omega_{Bi}}{\omega^2}\right)}. \quad (6.6.42)$$

If the electromagnetic wave frequency exceeds ion-cyclotron frequency ( $\omega \gg \omega_B$ ), influence of the term  $\omega_B \omega_{Bi} / \omega^2$  in the denominator is negligible and the dispersion equation 6.6.42 coincides with Equation 6.6.39. Conversely, if the wave frequency is low ( $\omega \ll \omega_B$ ), then the term  $\omega_B \omega_{Bi} / \omega^2$  becomes dominant in the dispersion equation. In this case, the dispersion equation 6.6.42 can be rewritten as


$$\frac{c^2}{(\omega/k)^2} = 1 + \frac{\omega_p^2}{\omega_B \omega_{Bi}} = 1 + \frac{\mu_0 \rho}{B^2} \approx \frac{c^2}{v_A^2}. \quad (6.6.43)$$

This introduces again the Alfvén wave velocity  $v_A$  (Equation 6.5.17); here,  $\rho = n_e M$  is the mass density in completely ionized plasma (see Ginsburg, 1960; Ginsburg and Rukhadze, 1970).

## 6.7 Emission and Absorption of Radiation in Plasma, Continuous Spectrum

### 6.7.1 Classification of Radiation Transitions

Radiation occurs due to transitions between different energy levels of a quantum system: transition up corresponds to absorption of a quantum  $E_f - E_i = \hbar\omega$ , transition down the spectrum corresponds to emission  $E_i - E_f = \hbar\omega$  (see Section 6.7.2). From the point of classical electrodynamics, radiation is related to the nonlinear change of dipole momentum, actually with the second derivative of dipole momentum. Neither emission nor absorption of radiation is possible for free electrons. Electron collisions are necessary in this case. It will be shown in Section 6.7.4 that electron interaction with a heavy particle, ion, or neutral is able to provide emission or absorption, but electron–electron interaction cannot. It is convenient to classify different types of radiation according to the different types of electron transition from one state to another. Electron energy levels in the field of an ion as well as transitions between the energy levels are illustrated in Figure 6.32. The case when both initial and final electron states are in continuum is called the free–free transition. A free electron in this transition loses part of its kinetic energy in the Coulomb field of a positive ion or in interaction with neutrals. The emitted energy in this case is a continuum, usually infrared and called bremsstrahlung (direct translation—



**FIGURE 6.32**  
Energy levels and electron transitions induced by ion field.


Electron transition between a free state in continuum and a bound state in atom (see Figure 6.32) is usually referred to as the free-bound transition. The free-bound transitions correspond to processes of the radiative electron-ion recombination (see Section 2.3.5) and the reverse one of photoionization (see Section 2.2.6). Such kinds of transitions also could take place in electron-neutral collisions. In this case, these are related to photo-attachment and photo-detachment processes of formation and destruction of negative ions (see Section 2.4). The free-bound transitions correspond to continuum radiation. Finally, the bound-bound transitions mean transition between discrete atomic levels (see Figure 6.32) and result in emission and absorption of spectral lines. Molecular spectra are obviously much more complex than that of single atoms because of possible transitions between different vibrational and rotational levels.

### 6.7.2 Spontaneous and Stimulated Emission: Einstein Coefficients

Consider transitions between two states (upper “ $u$ ” and ground “ $0$ ”) of an atom or molecule with emission and absorption of a photon  $\hbar\omega$ , which is illustrated in Figure 6.33. The probability of a photon absorption by an atom per unit time (and, hence, atom transition “ $0$ ”  $\rightarrow$  “ $u$ ”) can be expressed as

$$P(“0”, n_\omega \rightarrow “u”, n_\omega - 1) = An_\omega. \quad (6.7.1)$$

Here,  $n_\omega$  is the number of photons;  $A$  is the Einstein coefficient, which



**FIGURE 6.33**  
Radiative transition between two energy levels.

characteristics. Similarly, the probability of atomic transition with a photon emission is

$$P("u", n_\omega \rightarrow "0", n_\omega + 1) = \frac{1}{\tau} + Bn_\omega. \tag{6.7.2}$$

Here,  $1/\tau$  is the frequency of spontaneous emission, which takes place without direct relation to external fields; Einstein coefficient  $B$  characterizes emission induced by an external electromagnetic field. The factors  $B$  and  $\tau$  as well as  $A$  depend only on atomic parameters. If the first right-hand side term in Equation 6.7.2 corresponds to spontaneous emission, the second term is related to the stimulated emission. To find relations between the Einstein coefficients  $A$ ,  $B$  and the spontaneous emission frequency  $1/\tau$ , analyze the thermodynamic equilibrium of radiation with the atomic system. In this case, the densities of atoms in lower and upper states (Figure 6.33) are related in accordance with the Boltzmann law (Equation 4.1.9) as

$$n_u = \frac{g_u}{g_0} n_0 \exp\left(-\frac{\hbar\omega}{T}\right), \tag{6.7.3}$$

where  $\hbar\omega$  is energy difference between the two states,  $g_u, g_0$  are their statistical weights. According to the Planck distribution, the average number of photons  $\bar{n}_\omega$  in one state can be determined as

$$\bar{n}_\omega = 1 / \left( \exp \frac{\hbar\omega}{T} - 1 \right). \tag{6.7.4}$$

Taking into account, the balance of photon emission and absorption (see Figure 6.33)

$$n_0 P("0", n_\omega \rightarrow "u", n_\omega - 1) = n_u P("u", n_\omega \rightarrow "0", n_\omega + 1), \tag{6.7.5}$$

which can be rewritten based on Equations 6.7.1 and 6.7.2 as

$$A + B\bar{n}_\omega = \left( \frac{1}{\tau} + B\bar{n}_\omega \right) \tag{6.7.6}$$

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.