

DONALD VOET
University of Pennsylvania
JUDITH G. VOET
Swarthmore College

BIOCHEMISTRY

SECOND EDITION



JOHN WILEY & SONS, INC.

Acquisitions Editor: *Nedah Rose*
Marketing Manager: *Catherine Faduska*
Designer: *Madelyn Lesure*
Manufacturing Manager: *Susan Stetzer*
Photo Researcher: *Hilary Newman*
Illustration Coordinator: *Edward Starr*
Illustrators: *J/B Woolsey Associates*
Production Management: *Pamela Kennedy Oborski and Suzanne Ingrao*

Cover and Part Opening Illustrations © *Irving Geis*
This book was set in 9.5/12 Times Roman by Progressive Information Technologies and printed and bound by Von Hoffman Press. The cover was printed by Lehigh Press Lithographers. Color separations by Lehigh Press Colortronics.

Cover Art: Two paintings of horse heart cytochrome *c* by Irving Geis in which the protein is illuminated by its single iron atom. On the front cover the hydrophilic side chains are drawn in green, and on the back cover the hydrophobic side chains are drawn in orange. The paintings are based on an X-ray structure by Richard Dickerson.

Recognizing the importance of preserving what has been written, it is a policy of John Wiley & Sons, Inc. to have books of enduring value published in the United States printed on acid-free paper, and we exert our best efforts to that end.

Copyright © 1995, by John Wiley & Sons, Inc.


All rights reserved. Published simultaneously in Canada.

Reproduction or translation of any part of this work beyond that permitted by Sections 107 and 108 of the 1976 United States Copyright Act without the permission of the copyright owner is unlawful. Requests for permission or further information should be addressed to the Permissions Department, John Wiley & Sons, Inc.

ISBN: 0-471-58651-X

Printed in the United States of America

10 9



12

Introduction to Enzymes

1. **Historical Perspective**
2. **Substrate Specificity**
 - A. Stereospecificity
 - B. Geometric Specificity
3. **Coenzymes**
4. **Regulation of Enzymatic Activity**
5. **A Primer of Enzyme Nomenclature**

The enormous variety of biochemical reactions that comprise life are nearly all mediated by a series of remarkable biological catalysts known as enzymes. Although enzymes are subject to the same laws of nature that govern the behavior of other substances, they differ from ordinary chemical catalysts in several important respects:

1. **Higher reaction rates:** The rates of enzymatically catalyzed reactions are typically factors of 10^6 to 10^{12} greater than those of the corresponding uncatalyzed reaction and are at least several orders of magnitude greater than those of the corresponding chemically catalyzed reactions.
2. **Milder reaction conditions:** Enzymatically catalyzed reactions occur under relatively mild conditions: temperatures below 100°C , atmospheric pressure, and nearly neutral pH's. In contrast, efficient chemical catalysis often requires elevated temperatures and pressures as well as extremes of pH.
3. **Greater reaction specificity:** Enzymes have a vastly greater degree of specificity with respect to the identities of both their substrates (reactants) and their products than do chemical catalysts; that is, enzymatic reactions

rarely have side products. For example, in the enzymatic synthesis of proteins on ribosomes (Section 30-3), polypeptides consisting of well over 1000 amino acid residues are made all but error free. Yet, in the chemical synthesis of polypeptides, side reactions and incomplete reactions presently limit the lengths of polypeptides that can be accurately produced in reasonable yields to ~10 residues (Section 6-4B).

4. **Capacity for regulation:** The catalytic activities of many enzymes vary in response to the concentrations of substances other than their substrates. The mechanisms of these regulatory processes include allosteric control, covalent modification of enzymes, and variation of the amounts of enzymes synthesized.

Consideration of these remarkable catalytic properties of enzymes leads to one of the central questions of biochemistry: *How do enzymes work?* We address this issue in the next part of the text.

In this chapter, following a historical review, we commence our study of enzymes with a discussion of two clear instances of enzyme action: one that illustrates how enzyme specificity is manifested, and a second that exemplifies the regulation of enzyme activity. These are by no means exhaustive treatments but are intended to highlight these all-important aspects of enzyme mechanism. We shall encounter numerous other examples of these phenomena in our study of metabolism (Chapters 15-26). These two expositions are interspersed with a consideration of the role of enzymatic cofactors. The chapter ends with a short synopsis of enzyme nomenclature. In Chapter 13 we take up the formalism of enzyme kinetics because the study of the rates of enzymatically catalyzed reactions provides indispensable mechanistic information. Finally, Chapter

34 is a general mechanism employed by eukaryotic cells.

1. HIST

The early history of yeast is largely that of a collection of together from various sources and considered as a single entity. Berzelius, in 1828, determined that yeast is a product of fermentation, and pointed out that it is known to catalyze the fermentation of sulfuric acid. In 1857, Louis Pasteur demonstrated that the process of fermentation is carried out by living cells. This discovery permitted the development of a more systematic method of yeast production that by 1870 had led to the commercial production of yeast. Indeed, the term "yeast" was coined in 1870 to emphasize the living nature of the yeast itself. Nevertheless, the term "yeast" obtained a new meaning in the synthesis of Section 16.

Eril Fisi-
gines can c
the formula
nity of an e
from their
chemical co
lated until
Sumner, w
ase, which
CO₂, demo
Since Sumr
ner, the pr
cepted unt
Moses Kun
when the e
and chymo
zymologica
that enzyrn
shown that
the Sectio

14 is a general discussion of the catalytic mechanisms employed by enzymes, followed by an examination of the mechanisms of several specific enzymes.

1. HISTORICAL PERSPECTIVE

The early history of enzymology, the study of enzymes, is largely that of biochemistry itself; these disciplines evolved together from nineteenth century investigations of fermentation and digestion. Research on fermentation is widely considered to have begun in 1810 with Joseph Gay-Lussac's determination that ethanol and CO_2 are the principal products of sugar decomposition by yeast. In 1835, Jacob Berzelius, in the first general theory of chemical catalysis, pointed out that an extract of malt known as diastase (now known to contain the enzyme α -amylase; Section 10-2D) catalyzes the hydrolysis of starch more efficiently than does sulfuric acid. Yet, despite the ability of mineral acids to mimic the effect of diastase, it was the inability to reproduce most other biochemical reactions in the laboratory that led Louis Pasteur, in the mid-nineteenth century, to propose that the processes of fermentation could only occur in living cells. Thus, as was common in his era, Pasteur assumed that living systems were endowed with a "vital force" that permitted them to evade the laws of nature governing inanimate matter. Others, however, notably Justus Liebig, argued that biological processes are caused by the action of chemical substances that were then known as "ferments." Indeed, the name "enzyme" (Greek: *en*, in + *zyme*, yeast) was coined in 1878 by Friedrich Wilhelm Kühne in an effort to emphasize that there is something *in* yeast, as opposed to the yeast itself, that catalyzes the reactions of fermentation. Nevertheless, it was not until 1897 that Eduard Buchner obtained a cell-free yeast extract that could carry out the synthesis of ethanol from glucose (alcoholic fermentation; Section 16-3B).

Emil Fischer's discovery, in 1894, that glycolytic enzymes can distinguish between stereoisomeric sugars led to the formulation of his **lock-and-key hypothesis**: *The specificity of an enzyme (the lock) for its substrate (the key) arises from their geometrically complementary shapes.* Yet, the chemical composition of enzymes was not firmly established until well into the twentieth century. In 1926, James Sumner, who crystallized the first enzyme, jack bean urease, which catalyzes the hydrolysis of urea to NH_3 and CO_2 , demonstrated that these crystals consist of protein. Since Sumner's preparations were somewhat impure, however, the protein nature of enzymes was not generally accepted until the mid-1930s, when John Northrop and Moses Kunitz showed that there is a direct correlation between the enzymatic activities of crystalline pepsin, trypsin, and chymotrypsin and the amounts of protein present. Enzymological experience since then has amply demonstrated that enzymes are proteins (although it has recently been shown that some species of RNA also have catalytic properties; Section 29-4B).

Although the subject of enzymology has a long history, most of our understanding of the nature and functions of enzymes is a product of the latter half of the twentieth century. Only with the advent of modern techniques for separation and analysis (Chapter 5) has the isolation and characterization of an enzyme become less than a monumental task. It was not until 1963 that the first amino acid sequence of an enzyme, that of bovine pancreatic ribonuclease A (Section 14-1A), was reported in its entirety, and not until 1965 that the first X-ray structure of an enzyme, that of hen egg white lysozyme (Section 14-2A), was elucidated. In the years since then, several thousand enzymes have been purified and characterized to at least some extent and the pace of this endeavor is rapidly accelerating.

2. SUBSTRATE SPECIFICITY

The noncovalent forces through which substrates and other molecules bind to enzymes are identical in character to the forces that dictate the conformations of the proteins themselves (Section 7-4): Both involve van der Waals, electrostatic, hydrogen bonding, and hydrophobic interactions. In general, a substrate-binding site consists of an indentation or cleft on the surface of an enzyme molecule that is complementary in shape to the substrate (geometrical complementarity). Moreover, the amino acid residues that form the binding site are arranged to interact specifically with the substrate in an attractive manner (electronic complementarity; Fig. 12-1). Molecules that differ in shape or func-




FIGURE 12-1. An enzyme-substrate complex illustrating both the geometrical and the physical complementarity between enzymes and substrates. Hydrophobic groups are represented by an h in a brown circle and dashed lines represent hydrogen bonds.

tional group distribution from the substrate cannot productively bind to the enzyme; that is, they cannot form enzyme-substrate complexes that lead to the formation of products. The substrate-binding site may, in accordance with the lock-and-key hypothesis, exist in the absence of bound substrate or it may, as suggested by the induced fit hypothesis (Section 9-4C), form about the substrate as it binds to the enzyme. *X-Ray studies indicate that the substrate-binding sites of most enzymes are largely preformed but that most of them exhibit at least some degree of induced fit upon binding substrate.*


A. Stereospecificity

Enzymes are highly specific both in binding chiral substrates and in catalyzing their reactions. This stereospecificity arises because enzymes, by virtue of their inherent chirality (proteins consist of only L-amino acids), form asymmetric active sites. For example, trypsin readily hydrolyzes polypeptides composed of L-amino acids but not those consisting of D-amino acids. Likewise, the enzymes involved with glucose metabolism (Section 16-2) are specific for D-glucose residues.

Enzymes are absolutely stereospecific in the reactions they catalyze. This was strikingly demonstrated for the case of yeast alcohol dehydrogenase (YADH) by Frank Westheimer and Birgit Vennesland. Alcohol dehydrogenase catalyzes the interconversion of ethanol and acetaldehyde according to the reaction:



The structures of NAD⁺ and NADH are presented in Fig. 12-2. Ethanol, it will be recalled, is a prochiral molecule (see Section 4-2C for a discussion of prochirality):



Ethanol's two methylene H atoms may be distinguished if the molecule is held in some sort of asymmetric jig (Fig. 12-3). The substrate-binding sites of enzymes are, of course,




FIGURE 12-2. The structures and reactions of nicotinamide adenine dinucleotide (NAD⁺) and nicotinamide adenine dinucleotide phosphate (NADP⁺). Their reduced forms are NADH and NADPH. [In the older literature they are termed diphosphopyridine nucleotide (DPN⁺) and triphosphopyridine nucleotide (TPN⁺) and their reduced forms are symbolized DPNH and TPNH.] These substances, which are collectively

referred to as the nicotinamide coenzymes or pyridine nucleotides (nicotinamide is a pyridine derivative) function, as indicated in later chapters, as intracellular carriers of reducing equivalents (electrons). Note that only the nicotinamide ring is changed in the reaction. Reduction formally involves the transfer of two hydrogen atoms (H⁺), although the actual reduction may occur via a different mechanism.

FIG
to c
bet

just
the
W
cific
serie
1. If
et

CH

No
chi
2. Up
reac

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.