

onglyza®
(saxagliptin) 5 mg tablets

For US Consumers

MENU



Register >



A MEDICATION FOR ADULTS WITH TYPE 2 DIABETES,
IN COMBINATION WITH DIET AND EXERCISE

A type 2 diabetes medication that can help you get closer to your goal

onglyza
Add an
BOOST

LEARN HOW ONGLYZA MAY HELP



Save with a \$0 co-pay offer

You may be eligible to get ONGLYZA for as little as \$0 a month for as long as your doctor prescribes.*

*Subject to eligibility. Restrictions apply. Not available for government-insured patients. Savings subject to monthly limit.

Do not take ONGLYZA if you:



Download the Doctor Discussion Guide

Talk to your doctor and find out if ONGLYZA is the right type 2 diabetes medication for you.



is can occur with ONGLYZA and may include swelling of the face, lips or throat, difficulty swallowing or breathing, swelling of the skin, hives, rash, itching, flaking, or peeling. If you have these symptoms, stop taking ONGLYZA and contact your doctor right away.



Register for support

Get ongoing prescription coverage support, savings



Important Safety Information

Do not take ONGLYZA if you:



- are allergic to any of its ingredients. Serious allergic reactions can occur with ONGLYZA and may include swelling of the face, lips or throat, difficulty swallowing or breathing, swelling of the skin, hives, rash, itching, flaking, or peeling. If you have these symptoms, stop taking ONGLYZA and contact your doctor right away.

with your kidneys. Contact your doctor right away if you have any of the following symptoms of heart failure: increasing shortness of breath or trouble breathing, especially when you lie down; an unusually fast increase in weight; swelling in the feet, ankles or legs

- When ONGLYZA is used with certain other diabetes medicines to treat high blood sugar, such as a sulfonyleurea or insulin, the risk of low blood sugar (hypoglycemia) is higher. Symptoms of low blood sugar include shaking, hunger, sweating, headache, rapid heartbeat, change in mood, and change in vision. Follow your doctor's instructions for treating low blood sugar
- Some people who take medicines called DPP-4 inhibitors like ONGLYZA, may develop joint pain that can be severe. Call your doctor if you have severe joint pain
- When ONGLYZA is used with a thiazolidinedione (TZD), such as pioglitazone or rosiglitazone, to treat high blood sugar, peripheral edema (fluid retention) may become worse. If you have symptoms of peripheral edema, such as swelling of hands, feet, or ankles, call your doctor

The most common side effects with ONGLYZA include upper respiratory tract infection, urinary tract infection, and headache.

Before you take ONGLYZA:

- Your doctor should test your blood to measure how well your kidneys are working before and during treatment with ONGLYZA. You may need a lower dose if your kidneys are not working well
- Tell your doctor if you start or stop taking other medications, including antibiotics, antifungals or HIV/AIDS medications, as your doctor may need to change your dose of ONGLYZA
- Tell your doctor if you are pregnant, breast-feeding, or plan to become pregnant or breast-feed

Approved Uses

ONGLYZA is a prescription medicine used along with diet and exercise to lower blood sugar in adults with type 2 diabetes.

ONGLYZA should not be used to treat people with type 1 diabetes or diabetic ketoacidosis (increased ketones in the blood or urine).

Please read full [Prescribing Information](#) and [Medication Guide](#) for ONGLYZA.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.FDA.gov/medwatch, or call 1-800-FDA-1088.

AZ&ME[™]

AstraZeneca Prescription Savings Program

Important Safety Information

Do not take ONGLYZA if you:

- are allergic to any of its ingredients. Serious allergic reactions can occur with ONGLYZA and may include swelling of the face, lips or throat, difficulty swallowing or breathing, swelling of the skin, hives, rash, itching, flaking, or peeling. If you have these symptoms, stop taking ONGLYZA and contact your doctor right away.



[Medication Guide](#) [Contact Us](#)[Site Map](#)[Privacy Notice](#) | [Legal Statement](#)

Important Safety Information

Do not take ONGLYZA if you:



- are allergic to any of its ingredients. Serious allergic reactions can occur with ONGLYZA and may include swelling of the face, lips or throat, difficulty swallowing or breathing, swelling of the skin, hives, rash, itching, flaking, or peeling. If you have these symptoms, stop taking ONGLYZA and contact your doctor right away.

| [Prescribing Information](#) | [Approved Uses](#) | [Med Guide](#) [Important Safety Information](#) | [Approved Uses](#)

| [Prescribing Information](#) | [Medication Guide](#)

For US Consumers

[Visit HCP Site >](#)

[Go to Healthcare Professional Site](#)

MENU

[Register »](#)




How ONGLYZA works with your body to lower blood sugar levels in type 2 diabetes management

Not all medicines for type 2 diabetes management work the same way. Find out below how ONGLYZA enhances the body’s natural ability to control blood sugar, which may help you with your type 2 diabetes management.


Important Safety Information

Do not take ONGLYZA if you are allergic to any of its ingredients. Serious allergic reactions can occur with ONGLYZA and may include swelling of the face, lips or throat, difficulty swallowing or breathing, swelling of the skin, hives, rash, itching, flaking, or peeling. If you have these symptoms, stop taking ONGLYZA and contact your doctor right away.



Serious side effects can happen to people taking ONGLYZA, including inflammation of the pancreas (pancreatitis) which may be severe and lead to death. Before taking ONGLYZA, tell your doctor if you ever had pancreatitis, gallstones, history of alcoholism, or high triglyceride levels. Stop taking



When you have type 2 diabetes, your pancreas doesn't release enough insulin or your body's cells may not use insulin properly. This can cause your blood sugar level to spike too high after meals.



ONGLYZA works by enhancing your natural ability to control blood sugar by helping the body increase the level of insulin after meals. This may result in lower after-meal blood sugar spikes.



Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.