

Yuri P. Raizer

Gas Discharge Physics

With 209 Figures



Springer

DOCKET
A L A R M

Find authenticated court documents without watermarks at docketalarm.com.

Professor Dr. Yuri P. Raizer
The Institute for Problems in Mechanics, Russian Academy of Sciences,
Vernadsky Street 101, 117526 Moscow, Russia

Editor:

Dr. John E. Allen
Department of Engineering Science, University of Oxford, Parks Road,
Oxford OX1 3PJ, United Kingdom

Translator:

Dr. Vitaly I. Kisin
24 Varga Street, Apt. 9, 117133 Moscow, Russia

This edition is based on the original second Russian edition: *Fizika gazovogo razryada*
© Nauka, Moscow 1987, 1992

1st Edition 1991
Corrected 2nd Printing 1997

ISBN 3-540-19462-2 Springer-Verlag Berlin Heidelberg New York

Library of Congress Cataloging-in-Publication Data.

Raizer, Ū. P. (Ūrii Petrovich) [Fizika gazovogo razrīada. English] Gas discharge physics / Yuri P. Raizer. p. cm. "Corr. printing 1997" – t.p. verso. Includes bibliographical references and index. ISBN 3-540-19462-2 (hardcover: alk. paper) 1. Electric discharges through gases. I. Title. QC711.R22713 1997 537.53-dc21 96-53988

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilm or in any other way, and storage in data banks. Duplication of this publication or parts thereof is permitted only under the provisions of the German Copyright Law of September 9, 1965, in its current version, and permission for use must always be obtained from Springer-Verlag. Violations are liable for prosecution under the German Copyright Law.

© Springer-Verlag Berlin Heidelberg 1991
Printed in Germany

The use of general descriptive names, registered names, trademarks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

Typesetting: Springer T_EX inhouse system
Cover design: *design & production* GmbH, Heidelberg

SPIN 10565824 54/3144 – 5 4 3 2 1 0 – Printed on acid-free paper

3. Interaction of Electrons in an Ionized Gas with Oscillating Electric Field and Electromagnetic Waves

$$\Gamma_e \nu_u - qI, \quad (2.50)$$

$$(2.51)$$

1 of the hydrodynamic flux of efficient.³ The terms containing $\ll kT_e$, are neglected; according to collisions with molecules). The last term in (2.50) denotes the resultant creation rate that ionization potential.

t the equation for the rate of

$$(2.52)$$

o time, as in (2.49).

the effects caused by *spatial* contribution of heat conduction and passing thermal energy to ion with the term proportional

in (2.22, 50–52) in the form neglected out of Γ_e . Ignoring the amount that $\mu_+ \ll \mu_e$, we obtain

$$(2.53)$$

, to the same accuracy,

$$(2.54)$$

l in Sect. 9.7.

i (2.51) for F and λ_e stem from the distribution functions and $\nu_m(\epsilon) = \frac{1}{2}$ and $\lambda_e, \frac{1}{2}$, are set equal to 2; the

3.1 The Motion of Electrons in Oscillating Fields

Both the equations of electrodynamics and the equations of motion of electrons are linear with respect to the fields E , H and the velocity v of the electron. For this reason, the *superposition principle* holds. Any periodical field can be resolved into harmonic components, so that it is sufficient to consider only the *sinusoidal* field, all the more so because one normally deals with *monochromatic* fields and waves. In the case of nonrelativistic motion, the magnetic force of the wave, $e(v/c)H$, is much less than the electric force eE . Furthermore, the amplitude of electron oscillations in discharge processes is usually small in comparison with wavelength λ . We assume, therefore, that the electron is in a spatially uniform electric field $E = E_0 \sin \omega t$, $E_0 = \text{const}$.

3.1.1 Free Oscillations

Assume that an electron moves without collisions, an assumption that is meaningful if the electron performs a large number of oscillations in the interval between collisions, $\omega \gg \nu_m$. We integrate the equation of collisionless motion,

$$m\dot{v} = -eE_0 \sin \omega t, \quad \dot{r} = v,$$

to give

$$v = \frac{eE_0}{m\omega} \cos \omega t + v_0, \quad r = \frac{eE_0}{m\omega^2} \sin \omega t + v_0 t + r_0. \quad (3.1)$$

An electron oscillates at the frequency of the field; these oscillations are superimposed onto an arbitrary translation velocity v_0 . The displacement and oscillation velocities are

$$a = \frac{eE_0}{m\omega^2}, \quad u = \frac{eE_0}{m\omega}. \quad (3.2)$$

The displacement is in phase with the field, while the velocity is out of phase by $\pi/2$. The limiting case of "collisionless" oscillations is approximately realized at optical frequencies, and also at microwave frequencies at low pressures, $p \lesssim 10$ Torr.

3.1.2 Effect of Collisions

Collisions "throw off" the phase, thereby disturbing the purely harmonic course of the electron's oscillations. A sharp change in the direction of motion after

scattering stops the electron from achieving the full range of displacement (3.2) that the applied force can produce; the electron starts oscillating anew after each collision, with a new phase and new angle relative to the instantaneous direction of velocity. In order to take this factor into account, we add the rate of loss of momentum due to collisions to the equation of motion of the "mean" electron. As in the case of constant fields (Sect. 2.1.1), we have the equation for the mean velocity:

$$m\dot{v} = -eE_0 \sin \omega t - m\nu v_m, \quad \dot{r} = v. \quad (3.3)$$

The solution of (3.3), valid after several collisions, is

$$v = \frac{eE_0}{m\sqrt{\omega^2 + \nu_m^2}} \cos(\omega t + \varphi), \quad \varphi = \arctan \frac{\nu_m}{\omega},$$

$$r = \frac{eE_0}{m\omega\sqrt{\omega^2 + \nu_m^2}} \sin(\omega t + \varphi). \quad (3.4)$$

The amplitudes of displacement and velocity of the electron are less by a factor of $\sqrt{1 + \nu_m^2/\omega^2}$ than those for free oscillations. The higher the effective collision frequency ν_m , the smaller they are (ν_m is determined by the velocity of random motion, which is much greater in discharges than the oscillation velocity; see Sect. 3.2). The displacement is shifted in phase relative to the field, the phase shift increasing from 0 to $\pi/2$ as the relative role of collisions ν_m/ω increases from 0 to ∞ .

The oscillation displacement and velocity (3.4) can always be resolved into two components, one proportional to the magnitude of the field $E = E_0 \sin \omega t$, and the other to its rate of change, $\dot{E} = \omega E_0 \cos \omega t$:

$$r = \frac{eE_0}{m(\omega^2 + \nu_m^2)} \sin \omega t + \frac{\nu_m}{\omega} \frac{eE_0}{m(\omega^2 + \nu_m^2)} \cos \omega t,$$

$$v = \frac{\omega eE_0}{m(\omega^2 + \nu_m^2)} \cos \omega t - \frac{\nu_m eE_0}{m(\omega^2 + \nu_m^2)} \sin \omega t. \quad (3.5)$$

The ratio of the components is determined by the relative role of collisions and is unambiguously related to the phase shift φ . This form of presenting the solution adds visual clarity to the results of the subsequent sections.

Expressions (3.4,5) show that the role of collisions is characterized by the ratio of the effective frequency ν_m and the circular frequency of the field $\omega = 2\pi f$, which is greater than the frequency f by nearly an order of magnitude.¹ In the limit $\nu_m^2 \ll \omega^2$, formulas (3.4,5) are close to (3.1) for free oscillations. To illustrate numerical values, consider an example of microwave radiation at frequency $f = 3$ GHz; $\lambda = 10$ cm, $\omega = 1.9 \times 10^{10} \text{ s}^{-1}$. Let $p \approx 1$ Torr, then $\nu_m \approx 3 \times 10^9 \text{ s}^{-1} \ll \omega$; $E_0 = 500 \text{ V/cm}$, roughly corresponding to the threshold

¹ When the degree of spatial uniformity of the field is evaluated, the displacement amplitude must be compared not with wavelength $\lambda = c/f$, but with $\tilde{\lambda} = \lambda/2\pi$: $a/\tilde{\lambda} = eE_0/m\omega^2 \tilde{\lambda} = u/c$.

all range of displacement (3.2) starts oscillating anew after each collision, returning to the instantaneous direction of motion. In addition, we add the rate of loss of motion of the "mean" electron. We have the equation for the mean

$$(3.3)$$

ions, is

$$(3.4)$$

of the electron are less by a factor of ν_m/ω . The higher the effective collision frequency, the more the phase shift determined by the velocity of the electron relative to the field, the phase of collisions ν_m/ω increases

can always be resolved into a drift velocity v_d and an oscillation velocity v_0 of the field $E = E_0 \sin \omega t$,

$$(3.5)$$

The relative role of collisions is characterized by the parameter ν_m/ω . This form of presenting the subsequent sections.

Motion is characterized by the frequency of the field $\omega = 2\pi f$, an order of magnitude.¹ In (3.1) for free oscillations.

of microwave radiation at $\nu_m \approx 10^9 \text{ s}^{-1}$. Let $p \approx 1 \text{ Torr}$, then corresponding to the threshold

the displacement amplitude must be $a/\lambda = eE_0/m\omega^2 \lambda = u/c$.

of microwave breakdown at such pressures. Formulas (3.2) show that $a = 2.5 \times 10^{-3} \text{ cm}$, $u = 4.7 \times 10^7 \text{ cm/s}$. We find that $a \ll \lambda = 1.6 \text{ cm}$, that is, the field in the electromagnetic wave is "uniform".

3.1.3 Drift Oscillations

In the limit of very frequent collisions or relatively low frequencies, $\nu_m^2 \gg \omega^2$, the oscillation velocity drops to

$$v \approx -\frac{eE_0}{m\nu_m} \sin \omega t = -\frac{eE(t)}{m\nu_m} = -\mu_e E(t) = v_d(t). \quad (3.6)$$

At each moment of time, the oscillation velocity coincides with the drift velocity that corresponds to the field vector at this moment. For brevity, we refer to such oscillations in the *mobility regime* as *drift oscillations*.

An electron behaves as it would in a constant field, responding to relatively slow changes of the field. Its displacement,

$$r \approx A \cos \omega t, \quad A = \frac{eE_0}{m\nu_m\omega} = \frac{\mu_e E_0}{\omega}, \quad (3.7)$$

has an amplitude A less than that of free oscillations in the same field by a factor of $\nu_m/\omega \gg 1$.

The oscillations of electrons in rf fields (and of course, at lower frequencies) are of drift type. For example, the collision frequency at $f \approx 10 \text{ MHz}$, $\nu_m \approx 3 \times 10^9 \text{ ps}^{-1}$, exceeds $\omega \sim 10^8 \text{ s}^{-1}$ even at fairly low pressures of $p \sim 0.03 \text{ Torr}$. In order to maintain a low-pressure, weakly ionized plasma by an rf field, one usually needs the values of E_0/p of the same order as E/p in a constant field. Therefore, at $f \approx 10 \text{ MHz}$ and $E_0/p \approx 10 \text{ V}/(\text{cm} \cdot \text{Torr})$, we have $A \sim 0.1 \text{ cm}$ regardless of pressure.

3.2 Electron Energy

3.2.1 Collisionless Motion

If collisions do not occur, the field does no work, on the average, on an electron; indeed, (3.1) implies that

$$\langle -e\mathbf{E} \cdot \mathbf{v} \rangle = -\frac{eE_0^2}{m\omega} \langle \sin \omega t \cos \omega t \rangle - eE_0 \cdot v_0 \langle \sin \omega t \rangle = 0,$$

where angle brackets denote time averaging.

The electric field pumps up the motion of the electron only once, when it is switched on; then the electron's energy $mv^2/2$ pulsates but remains unchanged on the average. The time averaged energy $\langle mv^2/2 \rangle$ is made up of the energy of translational motion $mv_0^2/2$, corresponding to the mean velocity $v_0 = \langle v(t) \rangle$, and that of oscillations. In the case of free oscillations, the latter energy is

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.