



JOURNAL  
of the  
OPTICAL SOCIETY  
of  
AMERICA



JUNE

1980

VOLUME 70

NUMBER 6

ASML 1019

# JOURNAL OF THE OPTICAL SOCIETY OF AMERICA

Joseph W. Goodman, Editor

Honmai Goodman, Assistant to the Editor

## Associate Editors

(1981)		(1983)		(1985)	
Term Ending 31 December 1981		Term Ending 31 December 1983		Term Ending 31 December 1985	
Darrell E. Burch	C. E. Moore Sitterly	David E. Aspnes	Thomas Hirschfeld	Kenneth L. Andrew	Leo M. Hurvich
David A. DeWolf	David H. Rank	William M. Benesch	Karl A. Stetson	Stanley S. Ballard	Akira Ishimaru
Georg H. Hass	James A. R. Samson	William B. Bridges	William Streifer	Richard G. Brewer	B. E. A. Saleh
Robert E. Hufnagel	Richard Tousey	Steven F. Clifford	Adriaan Walther	William H. Carter	Murray Sargent III
R. Clark Jones	A. Francis Turner	B. Roy Frieden	James C. Wyant	Henry M. Crosswhite	Jeffrey H. Shapiro
David L. MacAdam	Dudley Williams	Nicholas George		David L. Fried	Gerald Westheimer
				Carl W. Helstrom	Gunter W. Wyszecki

## Panel of Topical Advisors

William B. Bridges, *Chairman*  
 Robert M. Boynton, *Color and Vision*  
 John C. Dainty, *Holography and Information Processing*  
 Sumner P. Davis, *Spectroscopy*

Freeman F. Hall, Jr., *Atmospheric Optics*  
 Peter Kaiser, *Fiber and Integrated Optics*  
 James E. Pearson, *Lasers and Electro-Optics*  
 William H. Price, *Optical Engineering*

Editorial Staff at AIP: Lawrence Feinberg, *Journal Supervisor*; William L. Siletti, *Senior Copy Editor*

Published monthly by the American Institute of Physics for the Optical Society of America, the *Journal of the Optical Society of America* contains papers and Letters that contribute significant new knowledge or understanding of any optical phenomenon, principle, or method. Reports of officers and committees of the Society are also included.

Submit manuscripts and all other materials for publication, including books for review, to

Dr. Joseph W. Goodman, Editor  
*Journal of the Optical Society of America*  
 570 University Terrace  
 Los Altos, California 94022

Submission is a representation that the manuscript has not been published previously or currently submitted for publication elsewhere. The manuscript should be accompanied by a statement transferring copyright from the authors (or their employers—whoever holds the copyright) to the Optical Society of America; a form for copyright transfer is occasionally printed in the back of this journal and is available from the Editor's office or AIP.

**Manuscripts** must be prepared in accordance with the requirements of the journal which are summarized on page (iii) of the January issue of the *Journal of the Optical Society of America* of this year. Authors should consult the Style Manual of AIP which can be obtained (for \$7.50 prepaid) by writing to the American Institute of Physics, 335 East 45th Street, New York, New York 10017. Manuscripts submitted in a form which is not appropriate for the Journal will be returned to the authors.

**Publication Charge:** Author's institutions are expected to pay a publication charge of \$80.00 per printed page. Failure to honor the publication charge will usually delay publication.

**AIP's Physics Auxiliary Publication Service (PAPS)** is a low-cost depository for material which is part of and supplementary to a published paper, but is too long to be included in the Journal; inquire of the Editor.

**Proof and all correspondence** concerning papers in the process of publication should be addressed as follows: Supervisor, Editorial Department, American Institute of Physics, 335 East 45th Street, New York, N.Y. 10017. Reference must be to title, author, journal, and scheduled date of publication. A limited number of alterations in proof are unavoidable, but the cost of making extensive alterations after the article has been set in type will be charged to the author.

**Copyright 1980, Optical Society of America.** Individual readers of this journal and nonprofit libraries acting for them are freely permitted to make fair use of the material in it, such as to copy an article for use in teaching or research. The code that appears on the first page of each article in this journal gives the per-article copying fee for each copy of the article made beyond the free copying permitted under Sections 107 and 108 of the U.S. Copyright Law. This fee should be paid through the Copyright Clearance Center, Inc., Box 765, Schenectady, New York 12301. The same fees and procedures are applicable to articles published in previous volumes of this journal. Permission is granted to quote excerpts from articles in this journal in scientific works with the customary acknowledgement of the source, including the author's name and the journal name, volume, page and year. Reproduction of figures and tables is likewise permitted in other articles and books, provided that the same information is printed with them and notification is given to the Optical Society of America. Republication or systematic or multiple reproduction of any material in this journal (including abstracts) is permitted only under license from the Optical Society of America; in addition, the Optical Society will require that permission also be obtained from one of the authors. Address inquiries and notices to the Executive Director, Optical Society of America, 1816 Jefferson Place, N.W., Washington, D. C. 20036. In the case of authors who are employees of the United States government or of its contractors or grantees, the Optical Society of America recognizes the right of the United States government to retain a nonexclusive, royalty-free license to use the author's copyrighted article for United States government purposes.



# Eleventh International Quantum Electronics Conference

## Session A

Monday, June 23, 1980

8:30 A.M. Independence Room

Laser Spectroscopy I

Chairman: P. F. Liao

**A.1. Doubly Excited Alkaline Earth Atoms\*** (Invited), T. F. GALLAGHER, K. A. SAFINYA, AND W. SANDNER, *SRI International, Menlo Park, CA*, AND W. E. COOKE, *University of Southern California, Los Angeles, California*. (30 min.)

We describe recent experiments to probe the properties of doubly excited autoionizing atoms using a laser spectroscopic technique. The physical basis of the excitation process as well as experiments to probe the autoionization process are described.

\* This work supported by AFOSR, NSF, and DOE.

**A.2. Rydberg Atom Masers** (Invited), S. HAROCHE, C. FABRE, P. GOY, M. GROSS, AND B. MOI, *Laboratoire de Physique de l'Ecole Normale Supérieure, Paris, France*. (30 min.)

Using optically pumped alkali Rydberg atoms as the active medium, we have recently developed new types of pulsed maser sources with unusual characteristics. Due to the giant electric dipole matrix elements of the Rydberg states, these masers have extremely low inversion thresholds—several orders of magnitude smaller than those of conventional masers—and “microscopic” energy outputs. Very sensitive detection procedures have to be used in order to observe their emission. An indirect detection method consists in studying the field ionization characteristics of the atoms, which is strongly modified when maser action occurs. Direct detection of the tiny microwave bursts has also been achieved by a heterodyne technique with a very sensitive Schottky diode mixer. Using both methods, we have made a detailed study of the emission characteristics (typical number of inverted atoms at threshold  $\sim 10^4$ ; pulse energy  $\sim 1$  to  $10$  eV; peak emission power  $\sim 10^{-12}$  W; pulse duration  $\sim 0.5$   $\mu$ s).

Owing to the large number of energy levels near the ionization limit, these new masers can be operated at many wavelengths ranging from the centimeter to the submillimeter range. In fact, the source should continuously evolve from the maser to the laser case when the binding energy of the levels involved in the emission is increased. These new coherent sources are bound to have very interesting applications in fundamental physics (study of superradiance of very small systems of atoms) and for the technology of millimeter and submillimeter wave detectors as well.

**A.3. Singlet-Triplet Mixing by Hyperfine Interactions in  $^3\text{He}$** , R. R. FREEMAN, P. F. LIAO, R. PANOCK, AND L. M. HUMPHREY, *Bell Telephone Laboratories, Holmdel, NJ 07733* (15 min.)

A complete determination of the hyperfine structure of the  $2^3P$  and  $3^3D$  states of  $^3\text{He}$  through analysis of Doppler-free intermodulated fluorescence spectra is reported. We find the structure of the  $3^3D$  state to be significantly modified by singlet-triplet mixing which is induced by hyperfine interactions. The hyperfine interaction is dominated by the Fermi contact interaction of the inner  $1s$  open shell electron with the nucleus and therefore does not decrease for higher lying states. Hence, unlike the case of one electronlike spectra (e.g., alkali atoms) or multielectron atoms with zero spin (e.g.,  $^4\text{He}$ ) the high Rydberg states of  $^3\text{He}$  will have their electronic structures completely dominated by the hyperfine interaction. In particular, the hyperfine induced singlet-triplet mixing for  $^3\text{He}$  will increase rapidly with increasing principal quantum number  $n$ . Our results are in good agreement with theoretical calculations of the hyperfine interaction.

In Fig. 1 we show a portion of our spectrum which contains transitions associated with the  $2^3P_{1,2}$  levels to the  $3^3D_{1,2,3}$  states of  $^3\text{He}$ . These states were obtained in a dc discharge tube operated with 0.8 Torr of  $^3\text{He}$ . The tube is probed with two counterpropagating tunable laser beams which are chopped at different frequencies. By monitoring fluores-

cence at the difference frequency we obtain the Doppler-free spectrum shown in Fig. 1. This spectrum is fit to a parametrized hyperfine Hamiltonian and the calculated resonance positions and line strength from the fit are shown in the figure. We find the majority of the interaction is due to the Fermi contact term of the  $1s$  electron and that this term is nearly the same for the  $2P$  and  $3D$  states as expected. This term produces sizeable singlet-triplet mixing which must be included to correctly give the structure. The dotted lines show resonance positions if one neglects this mixing.

Because the hyperfine interaction is essentially constant, our results, along with published fine structure measurements allow a precise determination of the structure of all higher lying states in  $^3\text{He}$ . We find, for example, we can reproduce the two-photon spectra recently obtained by Giacobino *et al.*<sup>1</sup> and also predict the hyperfine splittings of the  $n\ ^1D_2$  states observed in level crossing experiments.<sup>2</sup> In Table I we give our calculated values for these splittings and the measured experimental values. There is excellent agreement. As one of the simplest atoms, helium is amenable to accurate calculations. In Table I we also include the results of a theory based on hydrogenic electronic wave functions and good agreement with our calculations is again found.

In conclusion we have made a determination of the hyperfine interaction in  $^3\text{He}$ . This determination shows important singlet-triplet mixing effects which will dominate the




FIG. 1. Portion of Doppler-free spectrum of  $2^3D$ - $3^3D$  transitions in  $^3\text{He}$ . The levels are marked by  $(2^3P)F$ - $(2^3D)F'$ . The upper trace is the transmission of an interferometer having an FSR = 122.4 MHz. The solid lines show the calculated positions and intensities including singlet-triplet mixing. The dotted lines show the calculated positions if singlet-triplet mixing is ignored.



$\text{cm}^{-1}$ . The magnitude of the cross section for collisional deexcitation by spontaneous emissions,  $\sigma_S$ , was determined by measuring the total number of signal photons integrated over the emission bandwidth and using, for the number of detected photons,

$$N[\text{Ba}(5d^1D_2)]N[\text{Ti}(6p^2P_{3/2}^0)]\sigma_S\bar{V}\tau V_0\zeta,$$

where  $N[\text{Ba}(5d^1D_2)]$  and  $N[\text{Ti}(6p^2P_{3/2}^0)]$  are the number densities for the initial storage levels,  $\bar{V}$  is the mean velocity of collision,  $\tau$  is the effective radiating time,  $V_0$  is the effective radiating volume, and  $\zeta$  is the ratio of detected to generated photons. In this manner we obtain a measured value for the cross section for dipole-quadrupole collisional deexcitation of  $\sigma_S = 1.5 \times 10^{-22} \text{ cm}^2$ , with an overall experimental uncertainty of approximately a factor of 7.

The results of this experiment have application to the construction of low-gain, high-energy storage media and to the spectroscopic study of the interaction potentials of colliding atoms.

<sup>1</sup>S. E. Harris and J. C. White, *IEEE J. Quantum Electron.* **13**, 972 (1977).

<sup>2</sup>J. C. White, G. A. Zdasiuk, J. F. Young, and S. E. Harris, *Phys. Rev. Lett.* **41**, 1709 (1978); **42**, 480(E) (1979).

<sup>3</sup>W. R. Green, M. D. Wright, J. Lukasik, J. F. Young, and S. E. Harris, *Opt. Lett.* **4**, 265 (1979).

### J.8. Sodium Plasmas Produced by Milliwatt cw Laser Irradiation,\* M. E. KOCH, K. K. VERMA, AND W. C. STWALLEY, Iowa Laser Facility and Departments of Chemistry and Physics, University of Iowa, Iowa City, IA 52242. (15 min.)

There are a variety of reports of significant laser-produced ionization of alkali metal vapors using pulsed lasers (in  $\text{Li}^1$  and in  $\text{Na}^2$ ) and using cw lasers (in  $\text{Cs}^3$  and in  $\text{Na}^4$ ). (See also Ref. 5 for additional background.) While all of this work is quite interesting, much of it involved resonance lines<sup>1,2,4</sup> and so is perhaps not terribly surprising. The other work<sup>3</sup> on the other hand, involves transitions between a radiatively trapped upper level of an alkali-metal resonance line and a more highly excited level which can associatively ionize to form  $M_2^+$ . In principle, then, this cw plasma may contain a concentration of  $M_2^+$  which is quite nonequilibrium. Moreover, in contrast to a discharge where  $M_2^+$  may be rapidly destroyed by photodissociation, the  $M_2^+$  may be stable with respect to laser and other light (e.g., near-resonance lines) found in the laser-produced plasma.

With this in mind, we have irradiated a sodium heat pipe (typically at 10 Torr), using a focused cw dye laser at 5688.2 or 5682.6 Å ( $3p \rightarrow 4d$ ), and also reproduced the Cs result<sup>3</sup> at 6010 Å. Unlike Cs,<sup>3</sup> where a strong atomic ion-electron radiative recombination continuum is seen, we see no significant spectroscopic evidence for atomic ions in our Na plasma. We feel this is because in Cs, at the upper level of transition studied by Tam and Happer, the channel of ion pair formations ( $\text{Cs}^{**} + \text{Cs} \rightarrow \text{Cs}^+ + \text{Cs}^-$ ) is available in addition to associative ionization ( $\text{Cs}^{**} + \text{Cs} \rightarrow \text{Cs}_2^+ + e^-$ ). However, for  $\text{Na}^{**} = \text{Na}(4d)$ , only associative ionization can occur energetically, so we have produced essentially a

molecular ion plasma. Also we note that this plasma can be produced at quite low power ( $\sim 2 \text{ mW}$  focused broadband laser light at 5688.2 Å!) and we are currently examining the energy balance in detail.

We have obtained spectra of this plasma in the 2000–9000 Å region. The interpretation of this spectrum is still not completely clear. The various atomic lines seen can be understood in terms of  $\text{Na}(4d)$ – $\text{Na}$  and  $\text{Na}(3p)$ – $\text{Na}(3p)$  collisions<sup>6</sup> and the process (dissociative recombination):  $\text{Na}_2^+ + e^- \rightarrow \text{Na}^{**} + \text{Na}$ , where  $\text{Na}^{**}$  is a highly excited Na atom (e.g.,  $4d$  or  $5s$ ). The structure seen near the exciting line and to the red is presumably molecular fluorescence and  $D$  line absorption. We see five broad features at  $\sim 3650$ ,  $3780$ ,  $4350$ ,  $4520$ , and  $8000 \text{ Å}$  which remain to be explained.

We have examined the 4200–4700 Å region under high resolution and find the structure in that region to be a continuum, not densely spaced lines. A possible explanation is that these continua represent the processes  $\text{Na}_2^+ + e^- \rightarrow \text{Na}_2 + h\nu$ , where  $\text{Na}_2^+$  is an excited state of  $\text{Na}_2$ . The occurrence of such molecular ion-electron radiative recombination has never been previously established, although the atomic form is well known. The features we see peaking at 4350 and 4520 Å have been observed in other ways, e.g., in discharges,<sup>7–9</sup> in  $\text{Ar}^+$  laser UV-line irradiation of the  $\text{Na}_2 C \leftarrow X$  bands,<sup>10</sup> in two-photon  $\text{Na}_2$  excitation,<sup>11</sup> and in cw and pulsed dye-laser excitation at the  $\text{Na } D$  lines.<sup>12–14</sup> Similar features occur in K, Rb, and Cs.<sup>8,15,16</sup> Several explanations have been proposed involving free-free, free-bound, or bound-free processes. Note that the radiative recombination discussed above can be cast in “bound-free” form when a high molecular Rydberg state  $\text{Na}_2^+$  is formed as a resonance in  $e^-$ – $\text{Na}_2^+$  scattering. Since many of the potential energy curves of  $\text{Na}_2$  are fairly well known, e.g., from high quality *ab initio* calculations and a variety of recent experiments, we are carrying out explicit calculations of a number of these alternatives. We also have and will continue to carry out simultaneous ionization detection to attempt to resolve the origin of the 4350 and 4520 Å continua. Finally we note that some mechanisms suggest these bands might be made into a powerful violet laser with limited tunability.

The 8000 Å feature almost certainly corresponds to the

$$A^1\Sigma_u^+ - X^1\Sigma_g^+$$

satellite band<sup>17,18</sup> and has previously been observed in a Na (or other alkali) discharge.<sup>8,18,19</sup> There is a continuum overlaid with many discrete lines. The discrete lines, however, extend through all parts of the laser path while the continuum is concentrated in the central “white” region near the focus where the 4350 and 4520 Å features appear.

The 3650 and 3780 Å features have apparently not been previously reported. They also appear continuous and possible explanations for them are similar to those mentioned above for the 4350 and 4520 Å bands.

We are currently examining these spectra and extending them in a variety of ways, with

emphasis on obtaining microscopic understanding of the plasma formation process.

\*Supported by the National Aeronautics and Space Administration and the National Science Foundation.

<sup>1</sup>T. J. McIlrath and T. B. Lucatorto, *Phys. Rev. Lett.* **38**, 1390 (1977).

<sup>2</sup>T. B. Lucatorto and T. J. McIlrath, *Phys. Rev. Lett.* **37**, 428 (1976).

<sup>3</sup>A. Tam and W. Happer, *Opt. Commun.* **21**, 403 (1977).

<sup>4</sup>G. H. Bearman and J. J. Leventhal, *Phys. Rev. Lett.* **41**, 1227 (1978); **41**, 1759(E) (1978).

<sup>5</sup>G. S. Hurst, M. G. Payne, S. D. Kramer, and J. P. Young, *Rev. Mod. Phys.* **51**, 767 (1979).

<sup>6</sup>M. Allegrini, G. Alzetta, A. Kopystynska, L. Moi, and G. Orriols, *Opt. Commun.* **19**, 96 (1966).

<sup>7</sup>H. Bartels, *Z. Physik* **73**, 203 (1932).

<sup>8</sup>K. Schmidt *Proceedings of the Sixth International Conference on Ionization Phenomena in Gases* (Paris, 1963), Vol. 3, p. 323.

<sup>9</sup>J. J. de Groot and J. A. J. M. van Vliet, *J. Phys. D* **8**, 651 (1975).

<sup>10</sup>J. P. Woerdman, *Opt. Commun.* **26**, 216 (1978).

<sup>11</sup>J. P. Woerdman, *Chem. Phys. Lett.* **43**, 279 (1976).

<sup>12</sup>M. Allegrini, G. Alzetta, A. Kopystynska, L. Moi, and G. Orriols, *Opt. Commun.* **22**, 329 (1977).

<sup>13</sup>A. Kopystynska and P. Kowalczyk, *Opt. Commun.* **25**, 351 (1978).

<sup>14</sup>A. Kopystynska and P. Kowalczyk, *Opt. Commun.* **28**, 78 (1979).

<sup>15</sup>M. M. Rebeck and J. M. Vaughan, *J. Phys. B* **4**, 258 (1971).

<sup>16</sup>J. M. Brom and H. P. Broida, *J. Chem. Phys.* **61**, 982 (1974).

<sup>17</sup>W. C. Stwalley, “Laser Manipulation of Metallic Vapors”, *Radiation Energy Conversion in Space*, edited by K. W. Billman, Vol. 61 of Progress in Astronautics and Aeronautics (1978), p. 593–601.

<sup>18</sup>L. K. Lam, A. Gallagher, and M. M. Hessel, *J. Chem. Phys.* **66**, 3550 (1977).

<sup>19</sup>P. P. Sorokin and J. R. Lankard, *J. Chem. Phys.* **55**, 3810 (1971).

### J.9. Laser-Induced Penning/Associative Ionization in Crossed Atomic Beams, P. POLAK-DINGELS, J.-F. DELPECH, AND J. WEINER, Department of Chemistry, University of Maryland, College Park, MD 20742. (15 min.)

Laser-switched or laser-modified collisions are the object of intensive theoretical and experimental study because they offer the possibility of controlling the relative probabilities of competing inelastic and reactive exit channels. The influence of the laser field is to modify the electronic states of the system during a collisional encounter. Laser-induced collisions are characterized by atomic-field interactions which are nonresonant with respect to dipole-allowed transitions of the separated collision partners. We discuss here new results on Penning/associative ionization of Na/Na collisions in the presence of optical field power densities of  $\approx 10^7 \text{ W/cm}^2$ .

The experimental set-up is as follows. Two alkali atomic-beam sources are mounted on a multiported vacuum chamber at right angles in the horizontal plane. Two laser beams enter from opposite ports and overlap at the interaction region with an angle of nearly  $180^\circ$ . The light sources are flashlamp pumped tunable dye lasers synchronized together and with a box car integrator/amplifier used to record the ion signal. A quadrupole mass filter, mounted above the