

JBT 00302

The formation of daptomycin by supplying decanoic acid to *Streptomyces roseosporus* cultures producing the antibiotic complex A21978C

F.M. Huber, R.L. Pieper and A.J. Tietz

Antibiotic Development Division, Eli Lilly and Company, Indianapolis, Indiana, U.S.A.

(Received 23 November 1987; accepted 9 January 1988)

Summary

Antibiotic substance A21978C is a complex of compounds having a common cyclic polypeptide nucleus and different fatty acid sidechains. Daptomycin is a semi-synthetic antimicrobial substance derived from the A21978C complex by a very elaborate chemical process. To obtain the daptomycin, the A21978C complex was first isolated from culture filtrates of *Streptomyces roseosporus* by several resin procedures. The resulting material was then 'blocked' and added to an *Actinoplanes* culture for deacylation. The protected A21978C nucleus was subsequently isolated by the same procedure as the parent complex and reacylated with the desired fatty acid (decanoic acid). The acylated compound was deblocked to yield daptomycin. This report describes the experimentation undertaken to establish a strategy to supply a very toxic precursor to cultures of *S. roseosporus* and, thereby, produce daptomycin biosynthetically.

Daptomycin; A21978C; Precursing

Introduction

The production of the antibiotic complex A21978C, by *Streptomyces roseosporus* NRRL11379 was originally described by Hamill and Hoehn (1980) and shown to be

Correspondence to: F.M. Huber, Antibiotic Development Division, Eli Lilly and Company, Indianapolis, Indiana, U.S.A.

Materials and Methods

The organism used in this study was a mutant derived from *S. roseosporus* NRRL 11379 and designated A21978.65. The inoculum development stages prior to the final reactor have been previously described (Huber et al., 1987). The medium used for the production of the A21978C antibiotic complex contained soybean flour (2.0%, w/v), $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (0.06%), dextrose (0.75%), potato dextrin (3.0%, w/v), cane molasses (0.25%, w/v), SAG471 silicone antifoam (0.02%, w/v) and water. The medium was adjusted to pH 7.0 with NaOH and sterilized for 45 min at 121 °C in a stirred reactor. The reactor was made of 304 stainless steel with an operating volume of 120 l. The reactor had four equally spaced vertical baffles, two impellers with six open turbine blades, and a single open-tube sparger. The reactor was inoculated with 400 ml of the secondary inoculum development stage. The culture was then agitated at 200–400 rpm, aerated at 1 v/v/m and maintained at 30 °C.

Biological mass was estimated by mycelial volume and dry weight. Mycelial volume (MV) was determined by centrifuging a known amount of medium at $1200 \times g$ for 10 min and noting the volume of the insolubles in relationship to the total volume of the sample. The dry weight of the mass was estimated by transferring the pellet in distilled water to a tared aluminum weight boat and drying to constant weight at 29 °C. Respiratory activity was estimated by analyzing the exit gas stream from the reactors by a Perkin Elmer mass spectrometer.

Glucose was determined by the Trinder method using glucose oxidase/peroxidase (Biodynamics/bmc, Indianapolis, IN). Total carbohydrate was estimated by the anthrone method (Morris, 1948). Inorganic phosphate and ammonia nitrogen were automatically determined by the 'Industrial Method No. 93-70W' (Technicon Industrial Systems, Tarrytown, New York).

The concentration of the various components of the A21978C complex were estimated by HPLC. The analytical HPLC column was from ES Industries (C18, 10 cm \times 4.6 mm). The mobile phase consisted of 1% ammonium dihydrogen phosphate and acetonitrile (64:36, v/v).

Results and Discussion

The data presented in Fig. 2 and Table 1 are characteristic of the initial experiments with *S. roseosporus*. The A21978C complex increased with time, but no daptomycin was formed. Oxygen demand reached a maximum at 30 h and then decreased. It was possible that although there was a relatively high concentration of polymeric carbohydrate in the broth, some of that carbon might not be readily available to the microorganism. Thus, more dextrin 700 (1%) was added at 73 h and a slight increase in oxygen demand was observed. After 75 h the oxygen demand decayed further to a steady level at approximately 100 h. Much like the oxygen demand pattern, the mycelial volume also decreased significantly after a maximum had occurred early in the process. It was possible that only the cellular system for catabolizing carbohydrates was decaying. If that was true, the culture could still




Fig. 2. Oxygen demand in original complex medium.

readily catabolize other substances (i.e. lipids, amino acids). If such a substance was toxic when in excess, it could be fed at a rate not to exceed the maximum rate for its oxidation. Under those conditions the toxic substance would never accumulate in the culture. Decanoic acid, a very toxic fatty acid, is the desired sidechain for the production of daptomycin. If decanoic acid was supplied to *S. roseosporus* so as not to accumulate, it might be possible to 'precurse' the A21978C complex and produce daptomycin biosynthetically. The latter achievement would save much processing chemistry. Some antibiotics have been successfully 'precursed' (Higuchi et al., 1946; Moyer and Coghill, 1947; Wolf and Arnstein, 1960).

Decanoic acid is a solid at the temperature of cultivation (30°C). In order to conveniently add decanoic acid to the stirred reactor, a metabolizable solvent was sought. The original solution was a 5% solution of decanoic acid in an equal mixture of ethanol and water. The resulting solution was fed to the culture at 50 ml h⁻¹ beginning at 40 h post inoculation. The information gained in these experiments is presented in Fig. 3 and Tables 2 and 3. The data clearly indicate that daptomycin could be produced biosynthetically. The concentration of daptomycin in the

TABLE 1

CHARACTERISTICS OF CULTURES OF *S. roseosporus* WITH 1% DEXTRIN 700 ADDED AT 73 h

h	pH	MV (%)	Total carbohydrate (mg ml ⁻¹)	Glucose (mg ml ⁻¹)	PO ₄ -P (μg ml ⁻¹)	NH ₃ -N (μg ml ⁻¹)	A21978C complex (μg ml ⁻¹)
0	6.5	6.0	48	6	14.0	63	
18	5.5	8.7	44	5	2.0	10	
42	7.1	20.0	29	0	0.4	203	
66	6.6	18.7	24	0	11.0	146	293
90	6.0	13.3	29	0	11.0	104	347
114	5.9	10.6	25	0	5.0	100	387
138	5.9	11.3	13	0	3.0	100	458




Fig. 3. Changes in oxygen consumption during the initial decanoic acid supplemented culture.

A21978C complex approximated 34%. The time the feed was initiated was clearly defined by increased oxygen demand. That the oxygen demand continued to decrease during the remainder of the process suggested that further increases in feed rates were necessary. The decrease in oxygen demand prior to the feed indicated

TABLE 2

MEDIUM CHARACTERISTICS DURING INITIAL DECANOIC ACID FED A21978C PROCESS

h	pH	MV (%)	Total CH ₂ O (mg ml ⁻¹)	Glucose (mg ml ⁻¹)	PO ₄ -P (μg ml ⁻¹)	NH ₃ -N (μg ml ⁻¹)	Total A21978C-like substance (μg ml ⁻¹)
0	6.3	7	44	7	19	71	
16	5.9	10	40	5	3	32	
40	7.35	19	34	0	1	210	
64	6.9	12	29	1	11	204	200
88	6.3	11	27	3	10	156	306
112	6.2	10	24	6	4	137	399
136	6.0	8	30	9	4	142	364

TABLE 3

FACTOR COMPOSITION AND CYCLE TIME DURING INITIAL DECANOIC ACID FED PROCESS

h	Factors (μg ml ⁻¹)				
	C ₁	C ₂	C ₅	C ₃	LY146032
64	60	71	0	33	36
88	73	105	16	40	72
112	82	133	22	54	108
136	72	109	19	42	122

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.