

Effects of single and repeated doses of omeprazole on gastric acid and pepsin secretion in man

C W HOWDEN, J A H FORREST, AND J L REID

From the University Department of Materia Medica and Gastroenterology Unit, Stobhill General Hospital, Glasgow

SUMMARY The effects of omeprazole, a substituted benzimidazole, on gastric acid and pepsin secretion have been studied in twelve healthy subjects. From six to eight hours after a single oral dose of 30 mg, there was a 66% reduction in basal acid output, and a 71.2% reduction in pentagastrin stimulated acid output. A single dose of 60 mg produced a 91.7% reduction in basal acid output and a 95.3% reduction in pentagastrin stimulated acid output. After seven days treatment with 30 or 60 mg daily, there was almost 100% inhibition of both basal and pentagastrin stimulated acid output. Omeprazole did not significantly affect pepsin secretion which is in keeping with its proposed mode of action, as an inhibitor of the H⁺/K⁺-ATPase enzyme on the secretory membrane of the parietal cell. There were no side effects after omeprazole either with single or repeated dosing.

The substituted benzimidazoles are new agents which are potent inhibitors of gastric acid secretion. They act by selective, non-competitive inhibition of the H⁺/K⁺-ATPase enzyme in the parietal cell.¹ This enzyme is the active transport mechanism for hydrogen ion secretion in the stomach. Although one of these agents, omeprazole, has been shown to

suppress basal and pentagastrin stimulated acid secretion after a single dose,² little is known about its effects after repeated dosing. In addition, previous studies have used a buffered suspension of the drug² as it is partially inactivated by gastric acid, or have looked at its acid inhibitory effect 24 hours after a dose.³

We have studied the effects of two different doses of omeprazole on basal and pentagastrin stimulated acid and pepsin secretion after a single dose and after seven days of treatment. Capsules of enteric-coated omeprazole granules were used, and the effects studied around the time of expected maximum acid inhibition.

Methods

SUBJECTS

Twelve healthy male subjects, mean age 25.2 years

Address for correspondence: Dr C W Howden, Department of Materia Medica, Stobhill General Hospital, Glasgow G21 3UW, Scotland.

Received for publication 29 July 1983

of placebo followed by 30 mg or 60 mg of omeprazole daily for seven days. On each occasion, subjects arrived at the laboratory around 0830 hours and received a standard light breakfast after drug administration. They remained fasted for the next 12 hours. At around 1415 hours, a nasogastric tube was inserted into the stomach and its position checked by radiograph. From 1430 hours, gastric juice was aspirated continuously through the tube and minute samples for analysis were obtained. During the next 12 hours, gastric juice was

collected into one of 100 ml syringes. Each subject received one dose of omeprazole orally as a single dose, after three to five days by either 30 mg or 60 mg. Subjects attended the Clinical Research Laboratory on three occasions: day 0 (placebo); day 1 (omeprazole) and day seven (seventh day of treatment). The study design was identical for both groups. Subjects arrived fasted at around 0830 hours and were given capsules containing either 30 mg omeprazole granules or placebo. Breakfast was given 15 minutes after drug administration and the subject remained fasted for the remainder of the study day. At 1415 hours (or 5½ hours postdose) a size 14 gastric tube was passed and it was flushed with a water recovery tube until 1530 hours - that is, for 1½ hours postdose - gastric juice was aspirated and collected as four 15 ml samples. At 1600 hours, measurement of basal acid output was made. Each subject received a single dose of pentagastrin (1.2 µg/kg/h),

continuously aspirated and collected as four 15 minute samples for determination of plateau acid output.

Basal acid output was defined as the acid produced in the second half of the basal hour multiplied by a factor of two,⁵ and plateau acid output as the acid produced in the two consecutive highest 15 minute periods of the second hour multiplied by a factor of two. An intravenous infusion of pentagastrin was used to give a constant rate of stimulation, and the dose selected was designed to give a near maximal stimulus without producing systemic side effects.⁶ Acid output was determined by titration to pH 7.0 with 0.1M sodium hydroxide. A small aliquot of gastric juice was taken from each sample for estimation of pepsin secretion by the method of Berstad.⁷ Results are expressed as mean \pm standard deviation. Statistical comparisons were made using the Mann-Whitney U test.

Results

Basal and pentagastrin stimulated acid output after placebo were similar in the two groups. A single dose of 30 mg produced a 66% reduction in basal acid output, and a 71.2% reduction in plateau acid output ($p < 0.01$). The volume of gastric juice was reduced by 30.8% in the basal hour ($p < 0.05$) and by 47.1% in response to pentagastrin ($p < 0.01$). Basal pepsin secretion was not significantly altered although there was an apparently significant ($p < 0.05$) rise in pepsin secretion in the pentagastrin stimulated hour. After seven days of omeprazole 30 mg daily, basal acid output was reduced by 99.8% ($p < 0.05$) and plateau acid output by 98.4% ($p < 0.01$) when compared with values following placebo. Five of the six subjects produced no acid in the basal hour and three remained achlorhydric in response to pentagastrin. The volume of gastric juice

at the end of the seven day course was reduced by 54.9% ($p < 0.01$) and 79.2% ($p < 0.01$) in the basal and pentagastrin stimulated hours respectively. There were no significant changes in pepsin secretion either basally or after pentagastrin (Table 1).

A single 60 mg dose of omeprazole had a much greater effect on acid output reducing basal acid output and plateau acid output by 91.7% ($p < 0.05$) and 95.3% ($p < 0.01$) respectively. Volumes of gastric juice secreted in the basal and pentagastrin stimulated hours were reduced by 59.2% ($p < 0.01$) and 79.5% ($p < 0.01$) respectively. There was no significant change in pepsin secretion. After seven days treatment with omeprazole 60 mg daily, basal acid output was reduced by 99.1% ($p < 0.01$) and plateau acid output by 99.0% ($p < 0.01$). Volumes of gastric juice were reduced by 62.3% ($p < 0.01$) in the basal hour and by 83.8% ($p < 0.01$) in the pentagastrin stimulated hours. Pepsin secretion did not significantly change (Table 2).

No side effects were encountered with omeprazole. Each subject kept a diary for the duration of the study in which they were asked to note any adverse events and none was recorded. Subjects were also specifically asked about side effects on each of the three study days and, again, none was reported. Standard biochemical and haematological indices were measured before inclusion in the study, and were repeated within five days of completion of the study. No drug related abnormalities were found.

Discussion

These data show that omeprazole, in doses of 30 or 60 mg daily, is a potent inhibitor of gastric acid secretion, when this is assessed around the time of expected maximum effect. It has no significant effect

Table 1 Acid and pepsin output and volume of gastric juice after placebo, first dose of 30 mg omeprazole and seventh dose of 30 mg omeprazole. ($n=6$; mean \pm SD)

	Basal			Stimulated		
	Acid output (mmol/h)	Pepsin output (mg/h)	Volume (ml)	Acid output (mmol/h)	Pepsin output (mg/h)	Volume (ml)
Placebo	4.30 ± 3.37	22.2 ± 16.4	99.7 ± 34.3	35.41 ± 5.45	5.4 ± 2.9	302.0 ± 42.7
First dose	1.53 ± 1.91	17.8 ± 12.2	69.0 $\pm 22.6^*$	10.24 $\pm 10.06^\dagger$	36.8 $\pm 46.1^*$	159.7 $\pm 66.8^\dagger$
Seventh dose	0.01 $\pm 0.02^*$	5.2 ± 4.1	45.0 $\pm 12.0^\dagger$	0.58 $\pm 0.99^\dagger$	18.3 ± 18.8	62.7 $\pm 19.3^\dagger$

* $p < 0.05$. † $p < 0.01$

Table 2 Acid and pepsin output and volume of gastric juice after placebo, first dose of 60 mg omeprazole and seventh dose of 60 mg omeprazole. (n=6; mean \pm SD)

	Basal			Stimulated		
	Acid output (mmol/h)	Pepsin output (mg/h)	Volume (ml)	Acid output (mmol/h)	Pepsin output (mg/h)	Volume (ml)
Placebo	4.56 \pm 3.19	11.8 \pm 7.5	96.3 \pm 25.6	37.11 \pm 10.64	5.5 \pm 6.2	297.3 \pm 62.5
First dose	0.38 \pm 0.49*	13.1 \pm 17.1	39.3 \pm 14.0†	1.74 \pm 2.28†	35.2 \pm 18.8	61.0 \pm 25.2†
Seventh dose	0.04 \pm 0.09†	6.0 \pm 5.2	36.3 \pm 8.4†	0.37 \pm 0.20†	24.6 \pm 18.5	48.3 \pm 10.2†

* p<0.05. † p<0.01

on pepsin secretion. This is consistent with the proposed mechanism and site of action of the drug.

Sixty milligrams of omeprazole produced a greater inhibitory effect than 30 mg after a single dose. This is in agreement with previous work which has shown a prolonged inhibition of gastric acid secretion which is dose-dependent.² After seven days of treatment, there was no difference between the two dose levels with the majority of our subjects being achlorhydric – that is, pH of gastric juice >7.0, in the basal hour, and the response to pentagastrin being inhibited by around 99%. This would be consistent with an increasing response with time as a consequence of accumulation of the effect of omeprazole, which for a single dose normally exceeds 24 hours.

The reduction in the volume of gastric juice secreted after omeprazole was not of the same magnitude as the reduction in acid output. There was no concomitant reduction in pepsin secretion. These observations suggest that other components of the gastric secretion are relatively unaltered by omeprazole, and give further evidence for the drug's highly specific site and mode of action. The lack of reduction in pepsin secretion is unlikely to be of clinical importance as pepsin would be biologically inactive in the absence of intragastric acid.

It is of considerable interest that the drug had no observed haematological, biochemical, or subjective side effects, despite its extremely powerful action on gastric acid secretion. Omeprazole has now been given to patients with the Zollinger-Ellison syndrome^{8,9} with good therapeutic effect and without side effects. Clearly, a degree of caution is necessary with any new agent, but clinical trials in patients with active peptic ulceration seem indicated. The optimum dose for ulcer healing remains to be found but our studies indicate that even 30 mg omeprazole daily produces a degree of

inhibition of acid secretion in excess of that seen with H₂ receptor antagonists.^{10,11} The H₂ receptor antagonists do not have an increasing effect with time.

Although most of our subjects had basal achlorhydria after seven days treatment, we were assessing basal secretion around the time of the drug's maximal inhibitory effect. Other workers³ have found degrees of acid inhibition of the order of 30% 24 hours after a single dose, and have commented that the acid inhibitory effect reached a peak after five days treatment. It seems unlikely, therefore, that prolonged periods of achlorhydria would occur on long term treatment.

Omeprazole is a powerful inhibitor of gastric acid secretion in man, and a potentially useful agent in peptic ulcer.

This work has been presented in part to the Medical Research Society in January 1983 (*Clin Sci* 1983; **64**: 74p) and to the British Society of Gastroenterology in April 1983 (*Gut* 1983; **24**: A498).

Capsules of omeprazole and placebo were supplied by Astra Pharmaceuticals.

References

- 1 Fellenius E, Berglindh T, Sachs G *et al.* Substituted benzimidazoles inhibit gastric secretion by blocking (H⁺ + K⁺) ATPase. *Nature* 1981; **290**: 159–61.
- 2 Lind T, Cederberg C, Ekenved G *et al.* Effect of omeprazole – a gastric proton pump inhibitor – on pentagastrin stimulated acid secretion in man. *Gut* 1983; **24**: 270–6.
- 3 Muller P, Dammann HG, Seitz H *et al.* Effect of

- repeated once daily, oral omeprazole on gastric secretion. *Lancet* 1983; **1**: 66
- 4 Hassan M, Hobsley M. Positioning of subject and of nasogastric tube during a gastric secretion study. *Br Med J* 1970; **1**: 458-60.
 - 5 Faber RG, Hobsley M. Basal gastric secretion: reproducibility and relationship with duodenal ulceration. *Gut* 1977; **18**: 57-63.
 - 6 Mason MC, Giles GR, Clark CG. Continuous intravenous pentagastrin as a stimulant of maximal gastric acid secretion. *Gut* 1969; **10**: 34-8.
 - 7 Berstad A. A modified haemaglobin substrate method for the estimation of pepsin in gastric juice. *Scand J Gastroenterol* 1970; **5**: 343-8.
 - 8 Blachi A, Delchier JC, Soule JC *et al.* Control of acute Zollinger-Ellison syndrome with intravenous omeprazole. *Lancet* 1982; **2**: 1223-4.
 - 9 Oberg K, Lindstrom H. Reduction of gastric hypersecretion in Zollinger-Ellison syndrome with omeprazole. *Lancet* 1983; **1**: 66-7.
 - 10 Burland WL, Duncan WAM, Hesselbo T *et al.* Pharmacological evaluation of cimetidine, a new histamine H₂-receptor antagonist, in healthy man. *Br J Clin Pharmacol* 1975; **2**: 481-6.
 - 11 Peden NR, Saunders JHB, Wormsley KG. Inhibition of pentagastrin-stimulated and nocturnal gastric secretion by ranitidine, a new H₂-receptor antagonist. *Lancet* 1979; **1**: 690-2.