



US008603506B2

(12) **United States Patent**
Ashley

(10) **Patent No.:** **US 8,603,506 B2**
(45) **Date of Patent:** ***Dec. 10, 2013**

(54) **METHODS OF TREATING ACNE**

FOREIGN PATENT DOCUMENTS

- (75) Inventor: **Robert A. Ashley**, Newtown, PA (US)
- (73) Assignee: **Galderma Laboratories, Inc.**, Fort Worth, TX (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 70 days.

This patent is subject to a terminal disclaimer.

EP	0 410 099 A1	1/1991
JP	A-5-509288	12/1993
JP	A-91-88617	7/1997
JP	A-2001-72660	3/2001
WO	WO 83/00628	3/1983
WO	WO 92/03133	3/1992
WO	WO 98/05340	2/1998
WO	WO 98/08480	3/1998
WO	WO 99/58131	11/1999
WO	WO 00/15235	3/2000
WO	WO 00/28983	5/2000
WO	WO 00/07601	2/2001
WO	WO 00/18230	4/2011

(21) Appl. No.: **13/277,789**

(22) Filed: **Oct. 20, 2011**

(65) **Prior Publication Data**

US 2012/0108552 A1 May 3, 2012

Related U.S. Application Data

- (60) Continuation of application No. 11/876,478, filed on Oct. 22, 2007, now Pat. No. 8,052,983, which is a continuation of application No. 10/757,656, filed on Jan. 14, 2004, now abandoned, which is a division of application No. 10/117,709, filed on Apr. 5, 2002, now Pat. No. 7,211,267.
- (60) Provisional application No. 60/325,489, filed on Sep. 26, 2001, provisional application No. 60/281,916, filed on Apr. 5, 2001.

- (51) **Int. Cl.**
A61K 8/02 (2006.01)
A61K 31/74 (2006.01)
- (52) **U.S. Cl.**
USPC **424/401; 424/78.05**
- (58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,704,383	A	11/1987	McNamara et al.	
5,045,538	A	9/1991	Schneider et al.	
5,122,519	A	6/1992	Ritter	
5,260,292	A	11/1993	Robinson et al.	
5,300,304	A *	4/1994	Sheth et al.	424/490
5,532,227	A	7/1996	Golub et al.	
5,674,539	A	10/1997	Tomas et al.	
5,827,840	A	10/1998	Ramamurthy et al.	
5,908,838	A	6/1999	Gans	
5,998,390	A	12/1999	Ramamurthy et al.	
6,294,200	B1	9/2001	Conte et al.	
6,365,623	B1 *	4/2002	Perricone	514/448
6,455,583	B1 *	9/2002	Pflugfelder et al.	514/528
6,673,843	B2	1/2004	Arbiser	
7,014,858	B2	3/2006	Ashley	
2002/0165220	A1 *	11/2002	Heesch	514/210.09
2003/0082120	A1	5/2003	Milstein	
2003/0148945	A1	8/2003	McNicol et al.	
2005/0220222	A1	10/2005	Amin et al.	

OTHER PUBLICATIONS

Akamatsu, et al. "Effect of Keigai-Rengyo-To, a Japanese Kampo Medicine, on Neutrophil Functions: a Possible Mechanism of Action of Keigai-Rengyo-To in Acne," The Journal of International Medical Research, 25: 255-265 (1997).

Baer, et al. "High-Dose Tetracycline Therapy in Severe Acne," Arch Dermatol, 112:479-481 (Apr. 1976).

Cheryl Guttman, "Emerging resistance changes face of antibiotic therapy for acne," Dermatology Times, Jan. 2001, p. 22.

Hirohiko Akamatsu, Maki Asada, Jinro Komura, Yasuo Asada, and Yukie Niwa, "Effect of Doxycycline on the Generation of Reactive Oxygen Species: A Possible Mechanism of Action of Acne Therapy with Doxycycline," Acta Derm Venereol (Stockh) 72:178-178 (1992).

Bodokh, Y. Jacomet, J. Ph. Lacour and J.P. Ortonne, "Minocycline Induces an Increase in the Number of Excreting Pilosebaceous Follicles in Acne Vulgaris," Acta Derm Venereol (Stockh), 77:255-259 (1997).

W. J. Cunliffe, M.D., F.R.C.P., "Evolution of a Strategy for the Treatment of Acne," J Am Acad Dermatol, 16:591-9 (1987).

E. Anne Eady, Eileen Ingham, Christina E. Walters, Jonathan H. Cove, and William J. Cunliffe, "Modulation of Comedonal Levels of Interleukin-1 in Acne Patients Treated with Tetracyclines," J. Invest Dermatol, 101:86-91 (1993).

Boni E. Elewski, M.D., Beth A.J. Lamb, W. Mitchell Sams, Jr., M.D., and W. Ray Gammon, M.D., "In Vivo Suppression of Neutrophil Chemotaxis by Systemically and Topically Administered Tetracycline," J Am Acad Dermatol, 8:807-812 (1983).

Nancy B. Esterly, M.D., Nancy L. Furey, M.D., and Lillian E. Flanagan, B.S., "The Effect of Antimicrobial Agents on Leukocyte Chemotaxis," The Journal of Investigative Dermatology, 70(1):51-55 (1978).

Sainte-Marie, I. Tenaud, O. Jumbou and B. Dréno, "Minocycline Modulation of Alpha-MSH Production by Keratinocytes In vitro," Acta Derm Venereol 79:265-267 (1999).

Hoshiki Miyachi, M.D., Akira Yoshioka, M.D., Sadao Imamura, M.D., and Yukie Niwa, M.D., "Effect of Antibiotics on the Generation of Reactive Oxygen Species," J Invest Dermatol, 86(4):449-453 (1986).

Gerd Plewig, M.D., and Erwin Schöpf, M.D., "Anti-Inflammatory Effects of Antimicrobial Agents: An In Vivo Study," The Journal of Investigative Dermatology, 65:532-536 (1975).

(Continued)

Primary Examiner — Susan Tran
(74) *Attorney, Agent, or Firm* — Hoffmann & Baron, LLP

(57) **ABSTRACT**

A method of treating acne in a human in need thereof comprising administering systemically to said human a tetracycline compound in an amount that is effective to reduce acne and has substantially no antibiotic activity and a bisphosphonate compound.

Dr. Reddy's Laboratories, Ltd., et al.
v.
Galderma Laboratories, Inc.
IPR2015-_____

20 Claims, 1 Drawing



(12) **United States Patent**
Ashley

(10) **Patent No.:** **US 8,603,506 B2**
(45) **Date of Patent:** ***Dec. 10, 2013**

(54) **METHODS OF TREATING ACNE**

FOREIGN PATENT DOCUMENTS

- (75) Inventor: **Robert A. Ashley**, Newtown, PA (US)
- (73) Assignee: **Galderma Laboratories, Inc.**, Fort Worth, TX (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 70 days.

This patent is subject to a terminal disclaimer.

EP	0 410 099 A1	1/1991
JP	A-5-509288	12/1993
JP	A-91-88617	7/1997
JP	A-2001-72660	3/2001
WO	WO 83/00628	3/1983
WO	WO 92/03133	3/1992
WO	WO 98/05340	2/1998
WO	WO 98/08480	3/1998
WO	WO 99/58131	11/1999
WO	WO 00/15235	3/2000
WO	WO 00/28983	5/2000
WO	WO 00/07601	2/2001
WO	WO 00/18230	4/2011

(21) Appl. No.: **13/277,789**

OTHER PUBLICATIONS

(22) Filed: **Oct. 20, 2011**

- Akamatsu, et al. "Effect of Keigai-Rengyo-To, a Japanese Kampo Medicine, on Neutrophil Functions: a Possible Mechanism of Action of Keigai-Rengyo-To in Acne," The Journal of International Medical Research, 25: 255-265 (1997).
- Baer, et al. "High-Dose Tetracycline Therapy in Severe Acne," Arch Dermatol, 112:479-481 (Apr. 1976).
- Cheryl Guttman, "Emerging resistance changes face of antibiotic therapy for acne," Dermatology Times, Jan. 2001, p. 22.
- Hirohiko Akamatsu, Maki Asada, Jinro Komura, Yasuo Asada, and Yukie Niwa, "Effect of Doxycycline on the Generation of Reactive Oxygen Species: A Possible Mechanism of Action of Acne Therapy with Doxycycline," Acta Derm Venereol (Stockh) 72:178-178 (1992).
- Bodokh, Y. Jacomet, J. Ph. Lacour and J.P. Ortonne, "Minocycline Induces an Increase in the Number of Excreting Pilosebaceous Follicles in Acne Vulgaris," Acta Derm Venereol (Stockh), 77:255-259 (1997).
- W. J. Cunliffe, M.D., F.R.C.P., "Evolution of a Strategy for the Treatment of Acne," J Am Acad Dermatol, 16:591-9 (1987).
- E. Anne Eady, Eileen Ingham, Christina E. Walters, Jonathan H. Cove, and William J. Cunliffe, "Modulation of Comedonal Levels of Interleukin-1 in Acne Patients Treated with Tetracyclines," J Invest Dermatol, 101:86-91 (1993).
- Boni E. Elewski, M.D., Beth A.J. Lamb, W. Mitchell Sams, Jr., M.D., and W. Ray Gammon, M.D., "In Vivo Suppression of Neutrophil Chemotaxis by Systemically and Topically Administered Tetracycline," J Am Acad Dermatol, 8:807-812 (1983).
- Nancy B. Esterly, M.D., Nancy L. Furey, M.D., and Lillian E. Flanagan, B.S., "The Effect of Antimicrobial Agents on Leukocyte Chemotaxis," The Journal of Investigative Dermatology, 70(1):51-55 (1978).
- Sainte-Marie, I. Tenaud, O. Jumbou and B. Dréno, "Minocycline Modulation of Alpha-MSH Production by Keratinocytes In vitro," Acta Derm Venereol 79:265-267 (1999).
- Hoshiki Miyachi, M.D., Akira Yoshioka, M.D., Sadao Imamura, M.D., and Yukie Niwa, M.D., "Effect of Antibiotics on the Generation of Reactive Oxygen Species," J Invest Dermatol, 86(4):449-453 (1986).
- Gerd Plewig, M.D., and Erwin Schöpf, M.D., "Anti-Inflammatory Effects of Antimicrobial Agents: An In Vivo Study," The Journal of Investigative Dermatology, 65:532-536 (1975).

(65) **Prior Publication Data**

US 2012/0108552 A1 May 3, 2012

Related U.S. Application Data

- (60) Continuation of application No. 11/876,478, filed on Oct. 22, 2007, now Pat. No. 8,052,983, which is a continuation of application No. 10/757,656, filed on Jan. 14, 2004, now abandoned, which is a division of application No. 10/117,709, filed on Apr. 5, 2002, now Pat. No. 7,211,267.
- (60) Provisional application No. 60/325,489, filed on Sep. 26, 2001, provisional application No. 60/281,916, filed on Apr. 5, 2001.

(51) **Int. Cl.**

A61K 8/02 (2006.01)
A61K 31/74 (2006.01)

(52) **U.S. Cl.**

USPC **424/401**; 424/78.05

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,704,383	A	11/1987	McNamara et al.	
5,045,538	A	9/1991	Schneider et al.	
5,122,519	A	6/1992	Ritter	
5,260,292	A	11/1993	Robinson et al.	
5,300,304	A *	4/1994	Sheth et al.	424/490
5,532,227	A	7/1996	Golub et al.	
5,674,539	A	10/1997	Tomas et al.	
5,827,840	A	10/1998	Ramamurthy et al.	
5,908,838	A	6/1999	Gans	
5,998,390	A	12/1999	Ramamurthy et al.	
6,294,200	B1	9/2001	Conte et al.	
6,365,623	B1 *	4/2002	Perricone	514/448
6,455,583	B1 *	9/2002	Pflugfelder et al.	514/528
6,673,843	B2	1/2004	Arbiser	
7,014,858	B2	3/2006	Ashley	
2002/0165220	A1 *	11/2002	Heesch	514/210.09
2003/0082120	A1	5/2003	Milstein	

(Continued)

Primary Examiner — Susan Tran
(74) *Attorney, Agent, or Firm* — Hoffmann & Baron, LLP

(57) **ABSTRACT**

A method of treating acne in a human in need thereof comprising administering systemically to said human a tetracycline compound in an amount that is effective to treat acne but has substantially no antibiotic activity, without administering a bisphosphonate compound.

(56)

References Cited

OTHER PUBLICATIONS

M. Toyoda and M. Morohashi, "An Overview of Topical Antibiotics for Acne Treatment," *Dermatology*, 196:130-134 (1998).

Sheila E. Unkles, and Curtis G. Gemmell, "Effect of Clindamycin, Erythromycin, Lincomycin, and Tetracycline on Growth and Extracellular Lipase Production by Propionibacteria In Vitro," *Antimicrobial Agents and Chemotherapy*, 21:39-43 (1982).

G.F. Webster, K.J. McGinley, and J.J. Leyden, "Inhibition of Lipase Production in Propionibacterium acnes by Sub-Minimal-Inhibitory Concentrations of Tetracycline and Erythromycin," *British Journal of Dermatology*, 104:453-457 (1981).

Guy F. Webster, M.D., Ph.D., Susan M. Toso, M.S., and Lutz Hegemann, M.D., Ph.D., "Inhibition of a Model of In Vitro Granuloma Formation by Tetracyclines and Ciprofloxacin," *Arch Dermatol.*, 130:748-752 (1994).

Reynold C. Wong, M.D., Sewon Kang, M.P.H., Jan L. Heezen, L.P. N., John J. Voorhees, M.D., and Charles N. Ellis, M.D., "Oral Ibuprofen and Tetracycline for the Treatment of Acne Vulgaris," *J Am Acad Dermatol*, 11:1076-1081 (1984).

Kenneth S. Kornman and Edward H. Karl, "The Effect of Long-Term Low-Dose Tetracycline Therapy on the Subgingival Microflora in Refractory Adult Periodontitis," *J. Periodontol.*, 53(10):604-610 (Oct. 1982).

Bikowski, J.B., "Treatment of rosacea with doxycycline monohydrate," *Curtis*. 2000 Aug., 66(2):149-152.

Jimenez-Acosta, "Response to tetracycline of telangiectasias in male hemophiliac with human immunodeficiency virus infection," *J. Am. Acad. Dermatol.* Aug. 1988, 19(2 Pt. 1):369-379.

Torresani, C., "Clarithromycin versus doxycycline in the treatment of rosacea," *Int. J. Clin. Dermatol.* Dec. 1997, 36(12):942-946.

McClellan, K.J., "Topical Metronidazole. A review of its use in rosacea," *Am. J. Clin. Dermatol.* May-Jun. 2000, 1(3):191-199.

Quarterman, M.J., "Ocular Rosacea. Signs, symptoms and tear studies before and after treatment with doxycycline," *Arch. Dermatol.* Jan. 1997, 133(1):49-54.

Skidmore et al., "Effects of Subantimicrobial-Dose Doxycycline in the Treatment of Moderate Acne," *Archives of Dermatology* 139:459-464 (Apr. 2003), XP009047590.

Akamatsu, et al. "Effects of subminimal inhibitory concentrations of minocycline on neutrophil chemotactic factor production in comedonal bacteria, neutrophil phagocytosis and oxygen metabolism." *Arch Dermatol Res* 283:524-528 (1991).

Bikowski, et al. "Treatment of rosacea with doxycycline monohydrate" *Cutis*, 66:149-152 (Aug 2000).

Golub, et al. "Tetracyclines inhibit connective tissue breakdown: New therapeutic implications for an old family of drugs" *Critical Reviews in Oral Biology and Medicine*, 2(2):297-322 (1991).

Illig "Positive side effects of antibiotic and antimicrobial substances in therapy" *Infection* 7 (Suppl. 6): S 584-588 (1979) (English translation. Original document in German.).

Knight, et al. "A follow-up of tetracycline-treated rosacea" *British Journal of Dermatology* 93:577-580 (1975).

Marks, et al. "Comparative effectiveness of tetracycline and ampicillin in rosacea" *The Lancet*, 1049-1052 (Nov. 13, 1971).

Millar, et al. "A general practice study investigating the effect of minocycline (Minocin) 50 mg bd for 12 weeks in the treatment of acne vulgaris" *The British Journal of Clinical Practice* 41(8):882-886 (Aug. 1987).


Plewig, et al. *Acne: Morphogenesis and Treatment*, Springer-Verlag 297-301 (1975).

Webster, et al. "Suppression of Polymorphonuclear Leukocyte Chemotactic Factor Production in Propionibacterium acnes by Sub-minimal Inhibitory Concentrations of Tetracycline, Ampicillin, Minocycline, and Erythromycin" *Antimicrobial Agents and Chemotherapy* 21(5):770-772 (1982).

The Research Foundation of State University of New York; New York University; Galderma Laboratories, Inc; Galderma Laboratories, L.P.; and *Supernus Pharmaceuticals v. Mylan Pharmaceuticals Inc.* (Civ. No. 09-184-LPS) and *Mylan Pharmaceuticals Inc. v. The Research Foundation of State University of New York; New York University; Galderma Laboratories, Inc; Galderma Laboratories, L.P.; and Supernus Pharmaceuticals* (Civ. No. 10-892-LPS), Opinion, U.S. District Court for the District of Delaware, Case 1:09-cv-00184-LPS, Document 278, filed Aug. 26, 2011, pp. 1-77.

* cited by examiner

PHOTOOTOXICITY INDEX



Exh.

1

METHODS OF TREATING ACNE

CROSS-REFERENCE TO RELATED APPLICATION

This application is a continuation of U.S. application Ser. No. 11/876,478, filed on Oct. 22, 2007 now U.S. Pat. No. 8,052,983, allowed, which is a continuation of U.S. application Ser. No. 10/757,656, filed on Jan. 14, 2004, abandoned, which is a divisional application of U.S. application Ser. No. 10/117,709, filed on Apr. 5, 2002, now U.S. Pat. No. 7,211,267, which claims the benefit of U.S. Provisional Application No. 60/281,916, filed Apr. 5, 2001, and U.S. Provisional Application No. 60/325,489, filed Sep. 26, 2001. All of the aforementioned applications are incorporated herein by reference in their entireties.

BACKGROUND OF THE INVENTION

Acne is a common disease characterized by various types of lesions. The areas affected typically are areas of the skin where sebaceous glands are largest, most numerous, and most active. The lesions associated with acne are usually categorized as either non-inflammatory or inflammatory.


Non-inflammatory lesions include comedones. Comedones appear in two forms, open and closed. Comedones are thought to arise from abnormal follicular differentiation. Instead of undergoing shedding and discharge through the follicular orifice, abnormal desquamated cells (keratinocytes) become unusually cohesive, forming a micro-comedo or a microscopic hyperkeratotic plug in the follicular canal. The progressive accumulation of these microcomedones lead to visible comedones.

In its mildest form, acne is a more or less superficial disorder characterized by slight, spotty skin irritations. In such cases, ordinary skin hygiene is typically a satisfactory treatment. In the more inflammatory types of acne, however, pustules; infected cysts; and in extreme cases, canalizing, inflamed and infected sacs appear. Without effective treatment, these lesions may become extensive and leave permanent, disfiguring scars.

Microorganisms, especially *Propionibacterium acnes*, are strongly implicated in the pathogenesis of acne. The microorganisms are thought to release microbial mediators of inflammation into the dermis or trigger the release of cytokines from ductal keratinocytes.

Accordingly, the efficacy of antibiotics in treating acne is thought to be due, in significant part, to the direct inhibitory effect of the antibiotics on the growth and metabolism of these microorganisms. Systemically-administered tetracycline antibiotics, especially minocycline hydrochloride, are particularly effective in treating acne.


The tetracyclines are a class of compounds of which tetracycline is the parent compound. Tetracycline has the following general structure:



Structure A

2

The numbering system of the multiple ring nucleus is as follows:



Structure B

Tetracycline, as well as the 5-hydroxy (oxytetracycline, e.g. Terramycin) and 7-chloro (chlorotetracycline, e.g. Aureomycin) derivatives, exist in nature, and are all well known antibiotics. Semisynthetic derivatives such as 7-dimethylaminotetracycline (minocycline) and 6 α -deoxy-5-hydroxytetracycline (doxycycline) are also known tetracycline antibiotics. Natural tetracyclines may be modified without losing their antibiotic properties, although certain elements of the structure must be retained to do so.

In addition to the direct antibiotic activity of tetracyclines, further activities of antibiotic tetracyclines have been investigated for possible therapeutic effects on acne.

For example, a study by Elewski et al., *J. Amer. Acad. Dermatol.*, 8:807-812 (1983) suggests that acne therapy, consisting of orally-administered tetracycline at a total daily dose of 1000 mg, may have therapeutic anti-inflammatory effects in addition to antibiotic effects. In particular, it was found that the anti-inflammatory effect of tetracycline was, at least in part, due to inhibition of neutrophil chemotaxis induced by bacterial chemotactic factors.

A more recent study, performed by Eady et al., *J. Invest. Dermatol.*, 101:86-91 (1993), evaluated the effects of oral minocycline or tetracycline therapy on the cytokine and microflora content of open comedones in acne patients. The total daily dose of minocycline administered was 100 mg. The total daily dose of tetracycline administered was 1000 mg.

Eady et al. found that the therapies upregulated the production of bioactive IL-1 α -like material and immunochemical IL-1 β . IL-1 is considered to be a pro-inflammatory cytokine.

Accordingly to Eady et al., no overall decrease in the numbers of propionibacteria/mg of comedonal material was found. It is important to note, however, that the numbers of propionibacteria/mg of comedonal material are not expected to decrease in response to antibiotic therapy. Since the bacteria within comedones are encapsulated by the follicle, they are not susceptible to antibiotic treatment.

Another possible activity of tetracyclines in acne therapy was investigated by Bodokh, I., et al., *Acta. Derm. Venerol.*, 77:255-259 (1997). Their study was designed to evaluate the action of minocycline on sebaceous excretion in acne patients. A 100 mg daily dose of minocycline was administered. A subclinical increase in seborrhoea was reported. The authors propose that minocycline induces an increase in seborrhoea via a reduction in ductal obstruction. The mechanism by which the ductal obstruction is reduced is proposed to be a reduction in ductal irritation. The authors suggest that the reduction of ductal irritation is due to minocycline's direct effect on *P. acnes*, or minocycline's effect on the lipase produced by *P. acnes*.

Bodokh et al. also found that during treatment no correlation exists between seborrhoea intensity and clinical severity of acne. The authors state that the lack of correlation shows that seborrhoea is pathogenic because it is the "culture medium" of *P. acnes*. Thus, it can be concluded that the

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.