

## A novel mechanism of action of tetracyclines: Effects on nitric oxide synthases

ASHOK R. AMIN\*†‡§¶, MUKUNDAN G. ATTUR\*, GEETA D. THAKKER\*, PRAKASH D. PATEL\*, PRANAV R. VYAS\*, RAJESH N. PATEL\*, INDRAVADAN R. PATEL\*, AND STEVEN B. ABRAMSON\*†

\*Department of Rheumatology, Hospital for Joint Diseases, New York, NY 10003; and Departments of †Medicine and ‡Pathology, and §Kaplan Cancer Research Center, New York University Medical Center, New York, NY 10016

Communicated by H. Sherwood Lawrence, New York University Medical Center, New York, NY, September 23, 1996 (received for review April 10, 1996)

**ABSTRACT** Tetracyclines have recently been shown to have “chondroprotective” effects in inflammatory arthritides in animal models. Since nitric oxide (NO) is spontaneously released from human cartilage affected by osteoarthritis (OA) or rheumatoid arthritis in quantities sufficient to cause cartilage damage, we evaluated the effect of tetracyclines on the expression and function of human OA-affected nitric oxide synthase (OA-NOS) and rodent inducible NOS (iNOS). Among the tetracycline group of compounds, doxycycline > minocycline blocked and reversed both spontaneous and interleukin 1 $\beta$ -induced OA-NOS activity in *ex vivo* conditions. Similarly, minocycline  $\geq$  doxycycline inhibited both lipopolysaccharide- and interferon- $\gamma$ -stimulated iNOS in RAW 264.7 cells *in vitro*, as assessed by nitrite accumulation. Although both these enzyme isoforms could be inhibited by doxycycline and minocycline, their susceptibility to each of these drugs was distinct. Unlike acetylating agents or competitive inhibitors of L-arginine that directly inhibit the specific activity of NOS, doxycycline or minocycline has no significant effect on the specific activity of iNOS in cell-free extracts. The mechanism of action of these drugs on murine iNOS expression was found to be, at least in part, at the level of RNA expression and translation of the enzyme, which would account for the decreased iNOS protein and activity of the enzyme. Tetracyclines had no significant effect on the levels of mRNA for  $\beta$ -actin and glyceraldehyde-3-phosphate dehydrogenase nor on levels of protein of  $\beta$ -actin and cyclooxygenase 2 expression. These studies indicate that a novel mechanism of action of tetracyclines is to inhibit the expression of NOS. Since the overproduction of NO has been implicated in the pathogenesis of arthritis, as well as other inflammatory diseases, these observations suggest that tetracyclines should be evaluated as potential therapeutic modulators of NO for various pathological conditions.

Nitric oxide (NO), a multifunctional mediator produced by and acting on various cells, participates in inflammatory and autoimmune-mediated tissue destruction. NO is produced by a family of ubiquitous enzymes, nitric oxide synthases (NOSs). The overexpression of NOS in a variety of inflammatory tissues has led many to conclude that the modulation of NO synthesis and action could represent a new approach to the treatment of inflammatory and autoimmune conditions (1, 2). Where examined, NO formation is found to be increased in autoimmune diseases [rheumatoid arthritis (RA), systemic lupus erythematosus, ulcerative colitis, and Crohn disease], and several classic inflammatory symptoms (erythema and vascular leakiness) are reversed by NOS inhibitors (2–4). The most compelling evidence for NO as a mediator of tissue injury

has been in arthritis, based on studies carried out in animal models (5, 6), human osteoarthritis (OA) (7), and RA (8).

We have recently observed that human OA-affected cartilage can spontaneously release NO under *ex vivo* conditions in quantities sufficient to cause cartilage damage (7). The human OA-affected NOS (OA-NOS) is overexpressed in OA-affected cartilage and not detectable in normal cartilage. The inducible OA-NOS has properties similar to neuronal NOS (based on its molecular weight and antibody cross-reactivity among anti-NOS antibodies) and the 133-kDa inducible NOS [iNOS; sensitive to NF- $\kappa$ B and cycloheximide, up-regulated by interleukin (IL) 1 $\beta$ , tumor necrosis factor  $\alpha$ , and lipopolysaccharide (LPS)]. NO is known to potentiate matrix degradation, which includes inhibition of proteoglycan and collagen type II synthesis (9, 11) and up-regulation of metalloprotease activity (10).

Doxycycline and minocycline are members of the tetracycline family of broad-spectrum antibiotics. During recent years, it has been established that tetracyclines, which are rapidly absorbed and have a prolonged half-life, exert biological effects independent of their antimicrobial activity (11–13). Such effects include inhibition of matrix metalloproteases (MMPs) [including collagenase (MMP-1), gelatinase (MMP-2), and stromelysin (MMP-3) activity] and prevention of pathogenic tissue destruction (11). Furthermore, recent studies have also suggested that tetracyclines and inhibitors of metalloproteases inhibit tumor progression (14), bone resorption (15), and angiogenesis (16) and may have antiinflammatory properties (17).

In RA, these matrix metalloproteases have been identified in the synovial tissue, synovial fluids, and the proliferative pannus (18). These metalloproteases are also known to be up-regulated in OA-affected joints (19, 20). Interestingly, Yu *et al.* (21) have also shown that prophylactic administration of doxycycline markedly reduced the severity of OA in dog models. In humans, minocycline (a semisynthetic tetracycline) has recently been demonstrated to be superior to placebo in the treatment of RA (22).

Since NO is a putative mediator of inflammation that exerts catabolic effects on cartilage, including the activation of metalloproteases (10), we evaluated the action of tetracycline compounds on the spontaneous release of NO from OA-

Abbreviations: NOS, nitric oxide synthase; iNOS, inducible NOS; L-NMMA, L-N-monomethyl arginine; OA, osteoarthritis; RA, rheumatoid arthritis; IL-1 $\beta$ , interleukin 1 $\beta$ ; LPS, lipopolysaccharide; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; COX-2, cyclooxygenase 2; MMP, matrix metalloprotease; TGF- $\beta$ , transforming growth factor  $\beta$ ; RT-PCR, reverse transcription-coupled PCR; IFN- $\gamma$ , interferon  $\gamma$ .

†To whom reprint requests should be addressed at: Department of Rheumatology, Room 1600, Hospital for Joint Diseases, 301 East 17th Street, New York, NY 10003.

¶Cao, M., Westerhausen-Larson, A., N. Georgescu, H. I., Rizzo, C. F., Stefano (1996) 42nd Annual Meeting of the Or

Dr. Reddy's Laboratories, Ltd., et al.  
v.  
Galderma Laboratories, Inc.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in

## A novel mechanism of action of tetracyclines: Effects on nitric oxide synthases

ASHOK R. AMIN<sup>†‡§¶</sup>, MUKUNDAN G. ATTUR<sup>\*</sup>, GEETA D. THAKKER<sup>\*</sup>, PRAKASH D. PATEL<sup>\*</sup>, PRANAV R. VYAS<sup>\*</sup>, RAJESH N. PATEL<sup>\*</sup>, INDRAVADAN R. PATEL<sup>\*</sup>, AND STEVEN B. ABRAMSON<sup>††</sup>

<sup>\*</sup>Department of Rheumatology, Hospital for Joint Diseases, New York, NY 10003; and Departments of <sup>†</sup>Medicine and <sup>‡</sup>Pathology, and <sup>§</sup>Kaplan Cancer Research Center, New York University Medical Center, New York, NY 10016

Communicated by H. Sherwood Lawrence, New York University Medical Center, New York, NY, September 23, 1996 (received for review April 10, 1996)

**ABSTRACT** Tetracyclines have recently been shown to have “chondroprotective” effects in inflammatory arthritides in animal models. Since nitric oxide (NO) is spontaneously released from human cartilage affected by osteoarthritis (OA) or rheumatoid arthritis in quantities sufficient to cause cartilage damage, we evaluated the effect of tetracyclines on the expression and function of human OA-affected nitric oxide synthase (OA-NOS) and rodent inducible NOS (iNOS). Among the tetracycline group of compounds, doxycycline > minocycline blocked and reversed both spontaneous and interleukin 1 $\beta$ -induced OA-NOS activity in *ex vivo* conditions. Similarly, minocycline  $\geq$  doxycycline inhibited both lipopolysaccharide- and interferon- $\gamma$ -stimulated iNOS in RAW 264.7 cells *in vitro*, as assessed by nitrite accumulation. Although both these enzyme isoforms could be inhibited by doxycycline and minocycline, their susceptibility to each of these drugs was distinct. Unlike acetylating agents or competitive inhibitors of L-arginine that directly inhibit the specific activity of NOS, doxycycline or minocycline has no significant effect on the specific activity of iNOS in cell-free extracts. The mechanism of action of these drugs on murine iNOS expression was found to be, at least in part, at the level of RNA expression and translation of the enzyme, which would account for the decreased iNOS protein and activity of the enzyme. Tetracyclines had no significant effect on the levels of mRNA for  $\beta$ -actin and glyceraldehyde-3-phosphate dehydrogenase nor on levels of protein of  $\beta$ -actin and cyclooxygenase 2 expression. These studies indicate that a novel mechanism of action of tetracyclines is to inhibit the expression of NOS. Since the overproduction of NO has been implicated in the pathogenesis of arthritis, as well as other inflammatory diseases, these observations suggest that tetracyclines should be evaluated as potential therapeutic modulators of NO for various pathological conditions.

Nitric oxide (NO), a multifunctional mediator produced by and acting on various cells, participates in inflammatory and autoimmune-mediated tissue destruction. NO is produced by a family of ubiquitous enzymes, nitric oxide synthases (NOSs). The overexpression of NOS in a variety of inflammatory tissues has led many to conclude that the modulation of NO synthesis and action could represent a new approach to the treatment of inflammatory and autoimmune conditions (1, 2). Where examined, NO formation is found to be increased in autoimmune diseases [rheumatoid arthritis (RA), systemic lupus erythematosus, ulcerative colitis, and Crohn disease], and several classic inflammatory symptoms (erythema and vascular leakiness) are reversed by NOS inhibitors (2–4). The most compelling evidence for NO as a mediator of tissue injury

has been in arthritis, based on studies carried out in animal models (5, 6), human osteoarthritis (OA) (7), and RA (8).

We have recently observed that human OA-affected cartilage can spontaneously release NO under *ex vivo* conditions in quantities sufficient to cause cartilage damage (7). The human OA-affected NOS (OA-NOS) is overexpressed in OA-affected cartilage and not detectable in normal cartilage. The inducible OA-NOS has properties similar to neuronal NOS (based on its molecular weight and antibody cross-reactivity among anti-NOS antibodies) and the 133-kDa inducible NOS [iNOS; sensitive to NF- $\kappa$ B and cycloheximide, up-regulated by interleukin (IL) 1 $\beta$ , tumor necrosis factor  $\alpha$ , and lipopolysaccharide (LPS)]. NO is known to potentiate matrix degradation, which includes inhibition of proteoglycan and collagen type II synthesis (9, 10) and up-regulation of metalloproteinase activity (10).

Doxycycline and minocycline are members of the tetracycline family of broad-spectrum antibiotics. During recent years, it has been established that tetracyclines, which are rapidly absorbed and have a prolonged half-life, exert biological effects independent of their antimicrobial activity (11–13). Such effects include inhibition of matrix metalloproteinases (MMPs) [including collagenase (MMP-1), gelatinase (MMP-2), and stromelysin (MMP-3) activity] and prevention of pathogenic tissue destruction (11). Furthermore, recent studies have also suggested that tetracyclines and inhibitors of metalloproteinases inhibit tumor progression (14), bone resorption (15), and angiogenesis (16) and may have antiinflammatory properties (17).

In RA, these matrix metalloproteinases have been identified in the synovial tissue, synovial fluids, and the proliferative pannus (18). These metalloproteinases are also known to be up-regulated in OA-affected joints (19, 20). Interestingly, Yu *et al.* (21) have also shown that prophylactic administration of doxycycline markedly reduced the severity of OA in dog models. In humans, minocycline (a semisynthetic tetracycline) has recently been demonstrated to be superior to placebo in the treatment of RA (22).

Since NO is a putative mediator of inflammation that exerts catabolic effects on cartilage, including the activation of metalloproteinases (10), we evaluated the action of tetracycline compounds on the spontaneous release of NO from OA-

Abbreviations: NOS, nitric oxide synthase; iNOS, inducible NOS; L-NMMA, L-N-monomethyl arginine; OA, osteoarthritis; RA, rheumatoid arthritis; IL-1 $\beta$ , interleukin 1 $\beta$ ; LPS, lipopolysaccharide; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; COX-2, cyclooxygenase 2; MMP, matrix metalloproteinase; TGF- $\beta$ , transforming growth factor  $\beta$ ; RT-PCR, reverse transcription-coupled PCR; IFN- $\gamma$ , interferon  $\gamma$ .

<sup>¶</sup>To whom reprint requests should be addressed at: Department of Rheumatology, Room 1600, Hospital for Joint Diseases, 301 East 17th Street, New York, NY 10003.

<sup>||</sup>Cao, M., Westerhausen-Larson, A., Niyibizi, C., Kavalkovich, K., Georgescu, H. I., Rizzo, C. F., Stefanovic-Racic, M., & Evans, C. H. (1996) 42nd Annual Meeting of the Orthopaedic Research Society.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked “advertisement” in accordance with 17 U.S.C. 1085.

affected human cartilage in *ex vivo* conditions (7) and on iNOS in LPS-stimulated murine macrophages. Both these enzyme isoforms show distinct susceptibility to pharmacological intervention by hydrocortisone and transforming growth factor  $\beta$  (TGF- $\beta$ ) *in vitro* (7). In the present study we report that (i) doxycycline and minocycline inhibit the activity of murine macrophage iNOS (minocycline  $\geq$  doxycycline) and human inducible OA-NOS (doxycycline  $>$  minocycline), (ii) doxycycline and minocycline inhibit iNOS expression at the level of iNOS mRNA and protein expression, thereby down-regulating its specific activity, and (iii) unlike acetylating agents or competitive inhibitors of L-arginine, doxycycline and minocycline do not directly inhibit the catalytic activity of iNOS *in vitro* in the L-arginine  $\rightarrow$  L-citrulline conversion assay.

## MATERIALS AND METHODS

**Cell Lines and Reagents.** Murine macrophage cells (RAW 264.7) were obtained from the American Type Culture Collection. Anti-murine iNOS and anti-cyclooxygenase (COX-2) antibodies were obtained from Transduction Laboratories (Lexington, KY). OA-affected cartilage was obtained from OA patients who underwent knee replacement surgery and were free of steroidal/nonsteroidal antiinflammatory drugs for at least 2 weeks before surgery. Doxycycline, minocycline, hydrocortisone, and LPS were obtained from Sigma, and murine interferon (IFN)- $\gamma$  and human IL-1 $\beta$  were from Promega.

**Assay of OA-NOS in Organ Cultures.** This assay was basically carried out as described (7). Briefly, OA-affected cartilage was cut into 3-mm discs; four to six discs (100–200 mg) were placed in organ culture in 2 ml of medium (F-12 with 0.1% BSA) for 24–72 h in the incubator. The medium was analyzed for nitrite accumulation by modified Griess reaction (23).

**Western Blot Analysis.** Equal amounts of protein (25–50  $\mu$ g) estimated by BCA reagent (Pierce) were loaded onto SDS/PAGE gels and examined by Western blot analysis with a specific anti-iNOS or anti-cyclooxygenase 2 (COX-2) murine mAb as specified by Transduction Laboratories. Membranes with bound antibodies (e.g., iNOS) were stripped by submersion in stripping buffer (100 mM 2-mercaptoethanol/2% SDS/62.5 mM Tris-HCl, pH 6.7) and incubating at 50°C for 30 min with occasional agitation. Membranes were then washed twice for 10 min at room temperature using large volumes of wash buffer. This filter could then be reprobed with an anti-actin antibody provided by James L. Lessard (Children's Hospital Medical Center, Cincinnati). Blots were developed using the ECL Western blot system (Amersham). Quantitation of the bands was performed using a densitometer from Molecular Dynamics.

**Northern Blot Analysis.** Total RNA was isolated using TRI reagent (MRC, Cincinnati). Northern blot analysis was carried out as described (24, 25). Briefly, 30  $\mu$ g of RNA was subjected to electrophoresis and transferred via capillary action onto a nylon membrane (Zeta-Probe, Bio-Rad). The membrane was hybridized with [<sup>32</sup>P]dCTP-labeled iNOS cDNA (4-kb *Sma*I fragment), a gift from James Cunningham (Harvard Medical School), and the blot was exposed to Kodak x-ray film for 24–48 h with intensifying screens at –70°C. The  $\beta$ -actin and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) probes were purchased from CLONTECH and probed as described above. Quantitation was performed using a PhosphorImager or Personal Densitometer (SI) (Molecular Dynamics).

**Reverse Transcription-Coupled PCR (RT-PCR) Analysis.** The presence of iNOS and  $\beta$ -actin mRNA in cells was analyzed by RT of total RNA followed by PCR amplification of the cDNA as described (26, 27). The sense and antisense oligonucleotides for iNOS were, respectively, 5'-ACG-GAGAAGCTTAGATCTGGAGCAGAAGTG-3' (nt 142–171) and 5'-CTGCAGGTTGGACCACTGGATCCTGC-

63) and 5'-CGTCTCCGGAGTCCATCACA-3' (nt 534–552), respectively. The predicted PCR product of the iNOS cDNA was 654 bp; that of the  $\beta$ -actin cDNA was 508 bp. The cDNA was prepared from equal amounts (1  $\mu$ g) of total RNA using SuperScript RNase H reverse transcriptase (GIBCO/BRL). An equal amount of the cDNA was used to amplify the mRNA by PCR. PCR amplification was performed in 50  $\mu$ l of a solution containing 1.5 mM MgCl<sub>2</sub>, 500 ng of iNOS primer, 100 ng of  $\beta$ -actin primer, all four dNTPs (each at 0.2 mM), 2.5 units of *Taq* DNA polymerase (GIBCO/BRL). The cycle conditions for amplification of cDNA were 1 min at 94°C, 1–2 min at 55°C, and 3 min at 72°C for 30 cycles for both iNOS and  $\beta$ -actin.

**Assays for iNOS in Cell-Free Extracts.** Specific activity of iNOS was determined in cell-free extracts by monitoring the conversion of L-[<sup>3</sup>H]arginine to L-[<sup>3</sup>H]citrulline as described (25, 28). RAW 264.7 cells were induced with LPS (100 ng/ml) in the presence and absence of tetracyclines or hydrocortisone for 14–20 h. The cell-free extracts were prepared in Tris buffer (10 mM, pH 7.4) containing chymostatin at 10  $\mu$ g/ml, antipain at 10  $\mu$ g/ml, leupeptin at 10  $\mu$ g/ml, pepstatin at 10  $\mu$ g/ml, 1 mM DTT, and 1 mM phenylmethylsulfonyl fluoride (25). The protein was measured by BCA assay reagent using BSA as standard (29). The reaction mixture for iNOS assay was as described (25, 28). After 20 min, the assays were terminated by heating the reaction mixture at 90°C for 5 min; 10  $\mu$ l ( $\approx$ 100,000 cpm) of the supernatant was spotted on activated Avicel TLC plates (Analtech Associates). The TLC plates were developed in a solvent system consisting of ethanol/water/ammonia, 80:16:4 (vol/vol). Quantitation of the spot for L-[<sup>3</sup>H]citrulline was performed by a Bioscan System 200 imaging scanner.

## RESULTS AND DISCUSSION

**Effect of Doxycycline and Minocycline on Inducible OA-NOS Activity in Human OA-Affected Cartilage.** Studies in our laboratory have shown (7) that OA-affected cartilage spontaneously releases NO in *ex vivo* conditions sufficient to cause cartilage damage. NO has dual effects on matrix metabolism: not only does it potentiate activity of matrix-degrading metalloproteases (10) but also it inhibits synthesis of matrix components such as proteoglycans and collagen (9, 11). Therefore, we first examined whether doxycycline or minocycline could block human OA-NOS activity in *ex vivo* conditions. Generally accepted pharmacologically relevant concentrations were selected for this study based on previous reports (19, 30–32). OA-affected cartilage was obtained from patients with advanced OA undergoing knee replacement surgery. OA-affected cartilage slices (Fig. 1A) were incubated in 0.1% BSA/endotoxin-free medium with doxycycline or minocycline at 5–80  $\mu$ g/ml for 24, 48, and 72 h in *ex vivo* conditions. Activity of NOS was monitored at different time intervals by estimating nitrite, the stable end-product, as described (7). The amount of NO spontaneously released (as measured by nitrite accumulation) at 0, 24, 48, and 72 h was 0,  $4.8 \pm 0.38$ ,  $16.4 \pm 0.7$ , and  $17.8 \pm 0.9$   $\mu$ M, respectively. The results showed that doxycycline and minocycline significantly inhibited NO production in OA-affected cartilage in a dose-dependent manner (Fig. 1). These data also indicate that doxycycline was more potent in its ability to inhibit OA-NOS activity. For example, at 72 h, the IC<sub>50</sub> for doxycycline was 32  $\mu$ g/ml compared with 54  $\mu$ g/ml for minocycline. These experiments further indicate that doxycycline and minocycline not only blocked the ongoing production of NO by OA-NOS *ex vivo* but also caused a decline in nitrite accumulation in cartilage organ culture for at least 72 h under conditions in which nitrite continues to accumulate in control cultures (Fig. 1). In a separate experiment, OA-affected cartilage was also exposed to IL-1 $\beta$  at 5 ng/ml, which augmented the release of nitrite from  $3.6 \pm 1.7$  to  $16.7 \pm 2.5$   $\mu$ M ( $P < 0.0001$ ) at 72 h. Addition of doxycycline at 20 and 40




FIG. 1. (A) Effect of doxycycline and minocycline on OA-NOS expression in human OA-affected cartilage. OA-affected knee articular cartilage from one OA-affected patient was placed in organ culture in 2 ml of medium in the presence or absence (control) of doxycycline and minocycline at 5–80  $\mu\text{g/ml}$ . The spontaneous release of nitrite was monitored at different time intervals. Data are expressed as micromolar nitrite released (mean  $\pm$  SD,  $n = 3-4$ ). Zero time indicates spontaneous release of NO at 24, 48, and 72 h, which was  $4.8 \pm 0.38$ ,  $16.4 \pm 0.7$ , and  $17.8 \pm 0.9$ , respectively. Statistics were derived using unpaired Student's  $t$  test. Data represent one of three identical experiments with samples from different patients. (B) Effect of doxycycline and minocycline on nitrite accumulation in RAW 264.7 cells stimulated with LPS. Murine macrophage cells (RAW 264.7) were incubated with doxycycline or minocycline (5–80  $\mu\text{g/ml}$ ) for 1–2 h followed by addition of LPS at 100 ng/ml. After 14–20 h of incubation, the medium was used to estimate nitrite accumulation by the modified Griess reaction (23). Data are expressed as micromolar of nitrite accumulated at a given time interval ( $n = 3$ ). Statistics were derived using unpaired Student's  $t$  test. Data represent one of three similar experiments. Graphs were plotted using the DELTA graph curve-fitting program: polynomial of degree 3.

( $P < 0.0006$ ), respectively, when compared with the IL-1 $\beta$ -stimulated cartilage slices at the same time interval. Similarly, addition of minocycline (at 20 and 40  $\mu\text{g/ml}$ ) showed  $16.6 \pm 3.2$  ( $P < 0.47$ ) and  $10.0 \pm 2.0$  ( $P < 0.0024$ )  $\mu\text{M}$  of nitrite accumulation when compared with the IL-1 $\beta$ -stimulated cartilage slices. These experiments also demonstrated that the  $\text{IC}_{50}$  levels for tetracyclines to inhibit NOS activity augmented by IL-1 $\beta$  were similar to the spontaneous release of NO. The concentrations of doxycycline that inhibited NO production in our studies are comparable to those required for the inhibition of matrix metalloproteinases (19, 30–33). In those studies, doxycycline at 20–50  $\mu\text{g/ml}$  inhibited the activity of proteolytic enzymes such as collagenase and gelatinase, blocked proteoglycan degradation, reduced the cell death associated with proteoglycan loss, and augmented cartilage growth (31).

**Effects of Doxycycline and Minocycline on iNOS in Murine Macrophages.** Our recent studies have indicated that human inducible OA-NOS is distinct from murine and human iNOS, based upon its size, immunoreactivity, and susceptibility to TGF- $\beta$  and hydrocortisone (7). Therefore, we also evaluated the effect of tetracyclines on the expression of iNOS in murine macrophages. RAW 264.7 cells were activated with LPS (100 ng/ml) to induce iNOS (34) with and without doxycycline and minocycline at 5–80  $\mu\text{g/ml}$ . Fig. 1B shows a concentration-dependent inhibition of nitrite accumulation in cells stimulated with LPS in the presence of doxycycline at 5–80  $\mu\text{g/ml}$

incubation in these cells. In the same set of experiments, minocycline was also administered at concentrations ranging from 5 to 80  $\mu\text{g/ml}$ . The  $\text{IC}_{50}$  for minocycline was 17  $\mu\text{g/ml}$  at 14 h and 12  $\mu\text{g/ml}$  at 20 h of incubation in RAW 264.7 cells stimulated with LPS. Although a marginal difference in the potency of doxycycline and minocycline was seen at 20 h of incubation (based on  $\text{IC}_{50}$ ), significantly higher concentrations of doxycycline, as compared with minocycline, were generally required to inhibit iNOS by >50% at both time intervals. We also examined the effect of minocycline or doxycycline at 20  $\mu\text{g/ml}$  on murine iNOS expression when stimulated with IFN- $\gamma$  at 100 units/ml for 16 h (which induced  $16.4 \pm 0.5$   $\mu\text{M}$  nitrite). Addition of minocycline and doxycycline significantly decreased IFN- $\gamma$ -induced nitrite production to  $9.7 \pm 0.3$   $\mu\text{M}$  ( $P < 0.0001$ ) and  $13.1 \pm 1.1$   $\mu\text{M}$  ( $P < 0.005$ ), respectively.

These studies indicate that both doxycycline and minocycline inhibit NO production in murine macrophages stimulated with either LPS or IFN- $\gamma$ . Furthermore, these experiments together with our prior observations (7) show that the iNOS and inducible OA-NOS have distinct susceptibility to doxycycline, minocycline, TGF- $\beta$ , and hydrocortisone. This is not surprising since it has been reported that two different forms of collagenase, MMP-8 ( $\text{IC}_{50}$ , 7–15  $\mu\text{g/ml}$ ) and MMP-1 ( $\text{IC}_{50}$ , 140  $\mu\text{g/ml}$ ), in two different cell types (i.e., neutrophils and fibroblasts), show distinct susceptibility to inhibition by tetracyclines (35). Furthermore, it should be noted that the same enzyme expressed in two closely related cell lines can have differential susceptibility to tetracyclines. For example, two osteoblastic cell lines, UMR 106–01 ( $\text{IC}_{50}$ , >200  $\mu\text{g/ml}$ ) and ROS 17/2.8 ( $\text{IC}_{50}$ , 20–30  $\mu\text{g/ml}$ ), showed differential susceptibility to doxycycline when evaluated for gelatinase activity (36). Finally, another factor that may contribute to the differential  $\text{IC}_{50}$  values of tetracyclines on NOS activity in cartilage slices compared with macrophage cells is the ability of each drug to penetrate the cartilage matrix and act on chondrocytes (37).

In view of the recent observation that *in vivo* administration of minocycline in rats augments the percentage of splenocytes exhibiting a rise in intracellular  $\text{Ca}^{2+}$  under *ex vivo* conditions upon stimulation with concanavalin A, depending on the concentration of  $\text{Ca}^{2+}$  in the medium, we examined the effect of extracellular  $\text{Ca}^{2+}$  on the influence of iNOS expression upon stimulation with LPS (38). RAW 264.7 cells were supplemented (–1 h) with an additional 0, 0.34, and 0.68 mM  $\text{Ca}^{2+}$  above that present in the medium (1.3 mM) and stimulated with LPS. The nitrite levels at 24 h were  $24.5 \pm 4.9$ ,  $26.6 \pm 1.8$ , and  $27.3 \pm 0.3$   $\mu\text{M}$  in medium supplemented with 0, 0.34 mM, and 0.68 mM  $\text{Ca}^{2+}$ , respectively. Addition of LPS and minocycline at 40  $\mu\text{g/ml}$  at the time of the addition of  $\text{Ca}^{2+}$  (–1 h) reduced the nitrite levels to  $8.8 \pm 1.04$  ( $P < 0.0029$ ),  $12.7 \pm 0.7$  ( $P < 0.007$ ), and  $12.1 \pm 0.9$  ( $P < 0.006$ )  $\mu\text{M}$ , for the 0, 0.34 mM, and 0.68 mM  $\text{Ca}^{2+}$ -supplemented cultures, respectively. Thus, there was no significant difference between the  $\text{Ca}^{2+}$ -supplemented and nonsupplemented controls, indicating that supplemental  $\text{Ca}^{2+}$  does not have a significant influence on the tetracycline-dependent NOS inhibition in the murine macrophages tested *in vitro*. It should be noted that tetracyclines also inhibit IL-1 $\beta$ - and IFN- $\gamma$ -induced NOS expression; unlike LPS, these cytokines do not flux extracellular  $\text{Ca}^{2+}$  into the cells upon activation (39, 40). These experiments indicate that tetracyclines do not exert their effects on NOS via the chelation of extracellular  $\text{Ca}^{2+}$ . However, it should be noted that intracellular  $\text{Ca}^{2+}$  is critical for the enzyme activity of NOS, which usually appears 3–4 h after LPS stimulation (3).

Based on the above studies, we sought to evaluate the mechanism of action of tetracyclines on NOS expression in the murine macrophage model where the biochemistry, enzymology, and molecular biology of iNOS are well-characterized (3, 34, 41).

sion of OA-NOS directly from the OA-affected cartilage without disturbing the architecture of the cartilage, which plays a significant role in chondrocyte function.

Using LPS-induced RAW 264.7 cells, we examined the following hypotheses. Tetracyclines may (i) decrease only the catalytic activity of iNOS without influencing the expression of iNOS protein; (ii) decrease both the catalytic activity of iNOS and the expression of iNOS protein, which in turn cumulatively leads to decrease in the accumulation of nitrite in the medium; or (iii) decrease the expression of iNOS protein and, therefore, the production of nitrite.

**Effect of Doxycycline and Minocycline on the Enzyme Activity of iNOS in Whole Cells.** RAW 264.7 cells were exposed to doxycycline or minocycline in the presence of LPS for 16–18 h; cell-free extracts prepared at the end of each time period were evaluated for iNOS enzyme activity using the L-arginine → L-citrulline conversion assay in total cell extracts. As shown in Fig. 2, preexposure of cells to either doxycycline or minocycline inhibits the conversion of arginine to citrulline in cell lysates in a dose-dependent manner when compared with the control LPS-stimulated activity. Doxycycline at 20, 40, and 80 μg/ml significantly reduced citrulline accumulation by 57%, 72%, and 85%, respectively; similar inhibition was also observed for minocycline (45%, 69%, and 69%, respectively). As expected, pretreatment of cells with 10 μM hydrocortisone and 75 μM L-N-monomethyl arginine (L-NMMA) blocked iNOS activity by 60% and 64%, respectively.

**Direct Effect of Doxycycline and Minocycline on iNOS Enzyme Activity in Cell-Free Extracts.** Recent studies have indicated that tetracyclines inhibit collagenase activity via direct effects on the enzyme (30, 43), which could be partially reversed by addition of Ca<sup>2+</sup> to the reaction mixture (44). Another mechanism proposed for this phenomenon is that procollagenase is reduced to inactive fragments upon activation in the presence of doxycycline (43). We have recently shown that acetylating agents, such as aspirin and N-acetylimidazole (25), as well as competitive inhibitors of L-arginine (2–4), inhibit iNOS catalytic activity *in vitro*. To evaluate the direct effect of doxycycline and minocycline on iNOS enzyme activity, we induced RAW 264.7 cells with LPS for 16 h in the absence of these agents and prepared cell-free

extracts as a source of iNOS enzyme in L-arginine → L-citrulline conversion assay. Separate aliquots of equal amounts of enzyme (in total cell extracts) were preincubated for 15 min with doxycycline at 20–80 μg/ml, minocycline at 20–80 μg/ml, 1 mM N-acetylimidazole, and 200 μM L-NMMA, respectively, before the enzyme reaction was initiated by adding the cofactors. The experiment showed that, unlike N-acetylimidazole or L-NMMA, doxycycline/minocycline had no significant effect directly on the units of enzyme activity or specific activity of iNOS in cell-free extracts (Fig. 3). These experiments indicate that the action of these drugs on iNOS seems to be distinct from that reported for metalloproteases such as procollagenases (30, 43, 45). Minocycline and doxycycline could not block an ongoing L-arginine → L-citrulline reaction catalyzed by iNOS in cell-free extracts. Furthermore, these experiments also indicate that, unlike the inhibition of collagenase activity, which can be reversed by addition of 10–50 μM Ca<sup>2+</sup> in the enzyme assay (44), the tetracyclines exert no significant effect on the iNOS catalytic activity in cell-free extracts, and therefore, action via their Ca<sup>2+</sup>-chelating properties can be excluded.

**Effect of Doxycycline and Minocycline on iNOS Protein Expression in Murine Macrophages.** Based on the above data, which indicated that tetracyclines did not directly affect the specific activity of NOS, we assumed that the decrease in specific activity in the cells preincubated with tetracyclines might be due to a decrease specifically in the iNOS protein in the total cell extracts. Therefore, we examined iNOS protein expression. RAW 264.7 cells stimulated with LPS in the presence and absence of various concentrations of doxycycline and minocycline were incubated for 16 h; cell-free extracts were prepared. The extracts were examined for 133-kDa iNOS by Western blot analysis, using specific antibodies. Fig. 4 shows a dose-dependent inhibition of iNOS protein expression (IC<sub>50</sub>, <20 μg/ml) in the presence of both doxycycline and minocycline. There was no significant total effect on the levels of β-actin synthesis (data not shown) and COX-2 expression (that is also induced with iNOS by LPS) in the same samples treated with doxycycline or minocycline at 20–40 μg/ml, thus indicating that the effects of doxycycline and minocycline on iNOS are not nonspecific. It should be noted that the IC<sub>50</sub> values for the inhibition of nitrite accumulation and iNOS protein expression for each tetracycline derivative were similar, consis-




FIG. 2. Effect of doxycycline and minocycline on iNOS enzyme activity in RAW 264.7 cells stimulated with LPS. Murine macrophage cells (RAW 264.7) were incubated with doxycycline or minocycline (20–80 μg/ml), hydrocortisone (10 μM), or L-NMMA (75 μM) for 1 h, followed by addition of LPS at 100 ng/ml for 16–18 h. The cell-free extracts were prepared and the L-arginine → L-citrulline conversion assay was carried out in cell-free extracts. The data have been represented as specific activity of the enzyme, which was defined as nmol of citrulline per min per mg of protein. Percent inhibition




FIG. 3. Effect of doxycycline and minocycline on the specific activity of iNOS *in vitro*. RAW 264.7 cells were stimulated with LPS for 16 h and cell-free extracts were prepared as a source of iNOS. Various modulators including methanol (MeOH), the carrier for N-acetylimidazole (NAI), were added 15 min prior to the addition of the cofactors to initiate the iNOS reaction. The specific activity was calculated and percent inhibition compared with the original LPS

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.