



US007211267B2

(12) **United States Patent**
Ashley

(10) **Patent No.:** **US 7,211,267 B2**
(45) **Date of Patent:** **May 1, 2007**

(54) **METHODS OF TREATING ACNE**

(75) Inventor: **Robert A. Ashley**, Newtown, PA (US)

(73) Assignee: **CollaGenex Pharmaceuticals, Inc.**,
Newtown, PA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/117,709**

(22) Filed: **Apr. 5, 2002**

(65) **Prior Publication Data**

US 2003/0130240 A1 Jul. 10, 2003

Related U.S. Application Data

(60) Provisional application No. 60/325,489, filed on Sep. 26, 2001, provisional application No. 60/281,916, filed on Apr. 5, 2001.

(51) **Int. Cl.**

A61K 9/20 (2006.01)
A61K 9/48 (2006.01)
A61K 9/68 (2006.01)
A61K 37/18 (2006.01)

(52) **U.S. Cl.** **424/401; 424/440; 424/451; 424/464; 514/152**

(58) **Field of Classification Search** **424/401, 424/464, 465, 468, 451, 457; 514/152**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,704,383 A * 11/1987 McNamara et al. 514/152

5,122,519 A * 6/1992 Ritter 514/152
5,157,046 A * 10/1992 Van Wauwe et al. 514/397
5,413,777 A * 5/1995 Sheth et al. 424/490
5,532,227 A * 7/1996 Golub et al. 514/152
5,674,539 A 10/1997 Tomas et al.
5,827,840 A * 10/1998 Ramamurthy et al. 514/152
5,908,838 A * 6/1999 Gans 514/152
5,998,390 A * 12/1999 Ramamurthy et al. 514/94
6,664,287 B2 * 12/2003 Avery et al. 514/436
6,673,843 B2 * 1/2004 Arbiser 514/679
7,014,858 B2 * 3/2006 Ashley 424/401
2003/0082120 A1 * 5/2003 Milstein 424/59
2003/0139380 A1 * 7/2003 Robert 514/152

FOREIGN PATENT DOCUMENTS

EP 0 410 099 A1 1/1991
JP 02006437 A * 1/1990
WO WO 83/00628 3/1983
WO WO 99/58131 11/1999

OTHER PUBLICATIONS

Wong et al. "Oral ibuprofen and tetracycline for the treatment of acne vulgaris" (Journal of the American Academy of Dermatology), pp. 1076-1081; 1984.*

(Continued)

Primary Examiner—S. Tran


(74) Attorney, Agent, or Firm—Hoffmann & Baron, LLP

(57) **ABSTRACT**

A method of treating acne in a human in need thereof comprising administering systemically to said human a tetracycline compound in an amount that is effective to treat acne but has substantially no antibiotic activity, without administering a bisphosphonate compound.

31 Claims, 1 Drawing Sheet

PHOTOOTOXICITY INDEX




Dr. Reddy's Laboratories, Ltd., et al.
v.
Galderma Laboratories, Inc.
IPR2015-_____

OTHER PUBLICATIONS

- Akamatsu et al. "Effect of Doxycycline on the Generation of Reactive Oxygen Species" (Stockh) 1992; 72:178-179.*
- Komman et al. "The Effect of Long-Term Low-Dose Tetracycline Therapy on the Subgingival Microflora in Refractory Adult Periodontitis" (J. Periodontol.) 53(10) 604-610; 1982.*
- Stedman's Medical Dictionary 27th Edition.*
- Webster et al., "Inhibition of lipase production in *Propionibacterium acnes* by sub-minimal-inhibitory concentrations of tetracycline and erythromycin", *British Journal of Dermatology* (1981), pp. 453-457.*
- Plewig et al., "Acne: Morphogenesis and Treatment", Springer-Verlag, pp. 297-301 (1975).*
- Webster et al., "Suppression of Polymorphonuclear Leukocyte Chemotactic Factor Production in *Propionibacterium acnes* by Subminimal Inhibitory Concentrations of Tetracycline, Ampicillin, Minocycline, and Erythromycin", *Antimicrobial Agents and Chemotherapy* 21(5):770-772. 1982.*
- Kenneth S. Komman and Edward H. Karl, "The Effect of Long-Term Low-Dose Tetracycline Therapy on the Subgingival Microflora in Refractory Adult Periodontitis," *J. Periodontol.* 53(10) 604-610 (Oct. 1982).
- Akamatsu, et al. "Effect of Keigai-Rengyo-To, a Japanese Kampo Medicine, on Neutrophil Functions: a Possible Mechanism of Action of Keigai-Rengyo-To in Acne," *The Journal of International Medical Research*, 25: 255-265 (1997).
- Baer, et al. "High-Dose Tetracycline Therapy in Severe Acne," *Arch Dermatol*, 112:479-481 (Apr. 1976).
- Cheryl Guttman, "Emerging resistance changes face of antibiotic therapy for acne," *Dermatology Times*, Jan. 2001, p. 22.
- Hirohiko Akamatsu, Maki Asada, Jinro Komura, Yasuo Asada, and Yukie Niwa, "Effect of Doxycycline on the Generation of Reactive Oxygen Species: A Possible Mechanism of Action of Acne Therapy with Doxycycline," *Acta Derm Venereol (Stockh)* 72:178-178 (1992).
- Bodokh, Y. Jacomet, J. Ph. Lacour and J.P. Ortonne, "Minocycline Induces an Increase in the Number of Excreting Pilosebaceous Follicles in Acne Vulgaris," *Acta Derm Venereol (Stockh)*, 77:225-259 (1997).
- W. J. Cunliffe, M.D., F.R.C.P., "Evolution of a Strategy for the Treatment of Acne," *J Am Acad Dermatol*, 16:591-9 (1987).
- E. Anne Eady, Eileen Ingham, Christina E. Walters, Jonathan H. Cove, and William J. Cunliffe, "Modulation of Comedonal Levels of Interleukin-1 in Acne Patients Treated with Tetracyclines," *J. Invest Dermatol*, 101:86-91 (1993).
- Boni E. Elewski, M.D., Beth A.J. Lamb, W. Mitchell Sams, Jr., M.D., and Ray Gammon, M.D., "In Vivo Suppression of Neutrophil Chemotaxis by Systemically and Topically Administered Tetracycline," *J Am Acad Dermatol*, 8:807-812 (1983).
- Nancy B. Esterly, M.D., Nancy L. Furey, M.D., and Lillian E. Flanagan, B.S., "The Effect of Antimicrobial Agents on Leukocyte Chemotaxis," *The Journal of Investigative Dermatology*, 70(1):51-55 (1978).
- Sainte-Marie, I. Tenaud, O. Jumbou and B. Dréno, "Minocycline Modulation of Alpha-MSH Production by Keratinocytes In vitro," *Acta Derm Venereol* 79:265-267 (1999).
- Hoshiki Miyachi, M.D., Akira Yoshioka, M.D., Sadao Imamura, M.D., and Yukie Niwa, M.D., "Effect of Antibiotics on the Generation of Reactive Oxygen Species," *J Invest Dermatol*, 86(4):449-453 (1986).
- Gerd Plewig, M.D., and Erwin Schöpf, M.D., "Anti-Inflammatory Effects of Antimicrobial Agents: An In Vivo Study," *The Journal of Investigative Dermatology*, 65:532-536 (1975).
- M. Toyoda and M. Morohashi, "An Overview of Topical Antibiotics for Acne Treatment," *Dermatology*, 196:130-134 (1998).
- Sheila E. Unkles, and Curtis G. Gemell, "Effect of Clindamycin, Erythromycin, Lincomycin, and Tetracycline on Growth and Extracellular Lipase Production by *Propionibacteria* In Vitro," *Antimicrobial Agents and Chemotherapy*, 21:39-43 (1982).
- G.F. Webster, K.J. McGinley, and J.J. Leyden, "Inhibition of Lipase Production in *Propionibacterium acnes* by Sub-Minimal-Inhibitory Concentrations of Tetracycline and Erythromycin," *British Journal of Dermatology*, 104:453-457 (1981).
- Guy F. Webster, M.D., Ph.D., Susan M. Toso, M.S., and Lutz Hegemann, M.D., Ph.D., "Inhibition of a Model of In Vitro Granuloma Formation by Tetracyclines and Ciprofloxacin," *Arch Dermatol*, 130:748-752 (1994).
- Reynold C. Wong, M.D., Sewon Kang, M.P.H., Jan L. Heezen, L.P.N., John J. Voorhees, M.D., and Charles N. Ellis, M.D., "Oral Ibuprofen and Tetracycline for the Treatment of Acne Vulgaris," *J Am Acad Dermatol*, 11:1076-1081 (1984).
- Skidmore et al., "Effects of Subantimicrobial-Dose Doxycycline in the Treatment of Moderate Acne," *Archives of Dermatology* 139:459-464 (Apr. 2003), XP009047590.
- Bikowski, J.B., "Treatment of rosacea with doxycycline monohydrate," *Curtis*. 2000 Aug., 66(2):149-152.
- Jimenez-Acosta, "Response to tetracycline of telangiectasias in male hemophilic with human immunodeficiency virus infection," *J. Am. Acad. Dermatol.* Aug. 1988, 19(2 Pt. 1):369-379.
- Torresani, C., "Clarithromycin versus doxycycline in the treatment of rosacea," *Int. J. Clin. Dermatol.* Dec. 1997, 36(12):942-946.
- McClellan, K.J., "Topical Metronidazole. A review of its use in rosacea," *Am. J. Clin. Dermatol.* May-Jun. 2000, 1(3):191-199.
- Quarterman, M.J., "Ocular Rosacea. Signs, symptoms and tear studies before and after treatment with doxycycline," *Arch. Dermatol.* Jan. 1997, 133(1):49-54.
- Akamatsu, et al. "Effects of subminimal inhibitory concentrations of minocycline on neutrophil chemotactic factor production in comedonal bacteria, neutrophil phagocytosis and oxygen metabolism," *Arch Dermatol Res* 283: 524-528 (1991).
- Bikowski, et al. "Treatment of rosacea with doxycycline monohydrate" *Cutis*, 66: 149-152 (Aug 2000).
- Golub, et al. "Tetracyclines inhibit connective tissue breakdown: New therapeutic implications for an old family of drugs" *Critical Reviews in Oral Biology and Medicine*, 2(2): 297-322 (1991).
- Illig "Positive side effects of antibiotic and antimicrobial substances in therapy" *Infection* 7 (Suppl.6): S 584-588 (1979) (English translation. Original document in German.)
- Knight, et al. "A follow-up of tetracycline-treated rosacea" *British Journal of Dermatology* 93: 577-580 (1975).
- Marks, et al. "Comparative effectiveness of tetracycline and ampicillin in rosacea" *The Lancet*, 1049-1052 (Nov. 13, 1971).
- Millar, et al. "A general practice study investigating the effect of minocycline (Minocine) 50 mg bd for 12 weeks in the treatment of acne vulgaris" *The British Journal of Clinical Practice* 41(8):882-886 (Aug. 1987).
- Plewig, et al. *Acne: Morphogenesis and Treatment*, Springer-Verlag 297-301 (1975).
- Webster, et al. "Suppression of Polymorphonuclear Leukocyte Chemotactic Factor Production in *Propionibacterium acnes* by Subminimal Inhibitory Concentrations of Tetracycline, Ampicillin, Minocycline, and Erythromycin" *Antimicrobial Agents and Chemotherapy* 21(5):770-772 (1982).

* cited by examiner

FIG. 1 PHOTOTOXICITY INDEX



1

METHODS OF TREATING ACNE

CROSS-REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application No. 60/281,916, filed Apr. 5, 2001, and U.S. Provisional Application No. 60/325,489, filed Sep. 26, 2001, both of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

Acne is a common disease characterized by various types of lesions. The areas affected typically are areas of the skin where sebaceous glands are largest, most numerous, and most active. The lesions associated with acne are usually categorized as either non-inflammatory or inflammatory.


Non-inflammatory lesions include comedones. Comedones appear in two forms, open and closed. Comedones are thought to arise from abnormal follicular differentiation. Instead of undergoing shedding and discharge through the follicular orifice, abnormal desquamated cells (keratinocytes) become unusually cohesive, forming a micro-comedo or a microscopic hyperkeratotic plug in the follicular canal. The progressive accumulation of these microcomedones lead to visible comedones.

In its mildest form, acne is a more or less superficial disorder characterized by slight, spotty skin irritations. In such cases, ordinary skin hygiene is typically a satisfactory treatment. In the more inflammatory types of acne, however, pustules; infected cysts; and in extreme cases, canalizing, inflamed and infected sacs appear. Without effective treatment, these lesions may become extensive and leave permanent, disfiguring scars.

Microorganisms, especially *Propionibacterium acnes*, are strongly implicated in the pathogenesis of acne. The microorganisms are thought to release microbial mediators of inflammation into the dermis or trigger the release of cytokines from ductal keratinocytes.

Accordingly, the efficacy of antibiotics in treating acne is thought to be due, in significant part, to the direct inhibitory effect of the antibiotics on the growth and metabolism of these microorganisms. Systemically-administered tetracycline antibiotics, especially minocycline hydrochloride, are particularly effective in treating acne.


The tetracyclines are a class of compounds of which tetracycline is the parent compound. Tetracycline has the following general structure:



Structure A

2

The numbering system of the multiple ring nucleus is as follows:



Structure B

Tetracycline, as well as the 5-hydroxy (oxytetracycline, e.g. Terramycin) and 7-chloro (chlorotetracycline, e.g. Aureomycin) derivatives, exist in nature, and are all well known antibiotics. Semisynthetic derivatives such as 7-dimethylaminotetracycline (minocycline) and 6 α -deoxy-5-hydroxytetracycline (doxycycline) are also known tetracycline antibiotics. Natural tetracyclines may be modified without losing their antibiotic properties, although certain elements of the structure must be retained to do so.

In addition to the direct antibiotic activity of tetracyclines, further activities of antibiotic tetracyclines have been investigated for possible therapeutic effects on acne.

For example, a study by Elewski et al., *J. Amer. Acad. Dermatol.*, 8:807-812 (1983) suggests that acne therapy, consisting of orally-administered tetracycline at a total daily dose of 1000 mg, may have therapeutic anti-inflammatory effects in addition to antibiotic effects. In particular, it was found that the anti-inflammatory effect of tetracycline was, at least in part, due to inhibition of neutrophil chemotaxis induced by bacterial chemotactic factors.

A more recent study, performed by Eady et al., *J. Invest. Dermatol.*, 101:86-91 (1993), evaluated the effects of oral minocycline or tetracycline therapy on the cytokine and microflora content of open comedones in acne patients. The total daily dose of minocycline administered was 100 mg. The total daily dose of tetracycline administered was 1000 mg.

Eady et al. found that the therapies upregulated the production of bioactive IL-1 α -like material and immunochemical IL-1 β . IL-1 is considered to be a pro-inflammatory cytokine.

Accordingly to Eady et al., no overall decrease in the numbers of propionibacteria/mg of comedonal material was found. It is important to note, however, that the numbers of propionibacteria/mg of comedonal material are not expected to decrease in response to antibiotic therapy. Since the bacteria within comedones are encapsulated by the follicle, they are not susceptible to antibiotic treatment.

Another possible activity of tetracyclines in acne therapy was investigated by Bodokh, I., et al., *Acta. Derm. Venerol.*, 77:255-259 (1997). Their study was designed to evaluate the action of minocycline on sebaceous excretion in acne patients. A 100 mg daily dose of minocycline was administered. A subclinical increase in seborrhoea was reported. The authors propose that minocycline induces an increase in seborrhoea via a reduction in ductal obstruction. The mechanism by which the ductal obstruction is reduced is proposed to be a reduction in ductal irritation. The authors suggest that the reduction of ductal irritation is due to minocycline's direct effect on *P. acnes*, or minocycline's effect on the lipase produced by *P. acnes*.

Bodokh et al. also found that during treatment no correlation exists between seborrhoea intensity and clinical severity of acne. The authors state that the lack of correlation

authors consider the antibiotic activity of minocycline to be therapeutically significant with respect to acne.

Similarly, in a recent clinical study it was reported that tetracycline in sub-antibiotic doses had no clinical effect on acne. (Cunliffe et al., *J. Am. Acad. Dermatol.*, 16:591-9 (1987).) In particular, a 100 mg total daily dose of minocycline and a 1.0 g total daily dose of tetracycline were found to be necessary to successfully treat acne.

The antibiotic effects of antibiotics are generally directly proportional to the dose administered of the antibiotics. Accordingly, in moderate to severe (i.e. inflammatory) forms of acne, oral antibiotics are typically administered at high doses. For example, in conventional acne therapy, tetracycline is administered at an initial dose of 500 to 2,000 mg/day, followed by a maintenance dose of 250-500 mg/day.

Clearly, the state-of-the-art teaching is that the clinical efficacy of systemically-administered tetracyclines in the treatment of acne is due, at least in significant part, to the antibiotic effects of the tetracyclines. In addition to their antibiotic effects, it has been proposed that tetracyclines reduce the number of inflammatory lesions (papules, pustules and nodules) by a variety of non-antibiotic mechanisms. Such mechanisms include interfering with the chemotaxis of polymorphonuclear leukocytes (PMN) into the inflammatory lesion, inhibition of PMN derived collagenase, and by scavenging reactive oxidative species produced by resident inflammatory cells.

There is no disclosure in the prior art of using either a sub-antibiotic dose of an antibiotic tetracycline compound, or of using a non-antibiotic tetracycline compound for the treatment of acne.

The use of tetracycline antibiotics, however, can lead to undesirable side effects. For example, the long term administration of antibiotic tetracyclines can reduce or eliminate healthy microbial flora, such as intestinal flora, and can lead to the production of antibiotic resistant organisms or the overgrowth of yeast and fungi.

Accordingly, there is a need for an effective treatment of acne which causes fewer undesirable side effects produced by the systemically-administered antibiotics used in conventional acne therapy.

SUMMARY OF INVENTION

The present invention provides a method of treating acne in a human in need thereof. The method comprises administering systemically to the human a tetracycline compound in an amount that is effective to treat acne but has substantially no antibiotic activity (i.e. substantially no antimicrobial activity), without administering a bisphosphonate compound.

Additionally, the present invention provides methods for reducing the number of comedones, inhibiting oxidation of melanin, and/or inhibiting lipid-associated abnormal follicular differentiation in a human in need thereof. These methods comprise administering systemically to the human a tetracycline compound in an amount that is effective for its purpose, e.g., to reduce the number of comedones, to inhibit oxidation of melanin, and/or to inhibit lipid-associated abnormal follicular differentiation, but has substantially no antibiotic activity.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the phototoxicity factor (PIF) for some

COL	R7	R8	R9
308	hydrogen	hydrogen	amino
311	hydrogen	hydrogen	palmitamide
306	hydrogen	hydrogen	dimethylamino

For structures L, M, N or O the compounds indicated are as follows:

COL	R7	R8	R9
801	hydrogen	hydrogen	acetamido
802	hydrogen	hydrogen	dimethylaminoacetamido
804	hydrogen	hydrogen	nitro
805	hydrogen	hydrogen	amino

For structure P, R8 is hydrogen and R9 is nitro.

DETAILED DESCRIPTION

The present invention provides methods of treating acne. As used herein, the term "acne" is a disorder of the skin characterized by papules, pustules, cysts, nodules, comedones, and other blemishes or skin lesions. These blemishes and lesions are often accompanied by inflammation of the skin glands and pilosebaceous follicles, as well as, microbial, especially bacterial, infection.

For the purposes of this specification, acne includes all known types of acne. Some types of acne include, for example, acne vulgaris, cystic acne, acne atrophica, bromide acne, chlorine acne, acne conglobata, acne cosmetica, acne detergicans, epidemic acne, acne estivalis, acne fulminans, halogen acne, acne indurata, iodide acne, acne keloid, acne mechanica, acne papulosa, pomade acne, premenstrual acne, acne pustulosa, acne scorbutica, acne scrofulosorum, acne urticata, acne varioliformis, acne venenata, propionic acne, acne excoriee, gram negative acne, steroid acne, nodulocystic acne and acne rosacea. Acne rosacea is characterized by inflammatory lesions (erythema) and permanent dilation of blood vessels (telangiectasia).

The present invention is particularly effective in treating comedones, e.g., reducing the number of comedones. Both open and closed comedones can be treated in accordance with the methods of this invention.

The present invention can also be used to treat certain other types of acneiform dermal disorders, e.g. perioral dermatitis, seborrheic dermatitis in the presence of acne, gram negative folliculitis, sebaceous gland dysfunction, hidradenitis suppurativa, pseudo-folliculitis barbae, or folliculitis.

The method comprises the administration of a tetracycline compound to a human in an amount which is effective for its purpose e.g., the treatment of acne, including reducing the number of comedones, but which has substantially no antibiotic activity.

The tetracycline compound can be an antibiotic or non-antibiotic compound. The tetracycline compound has the general tetracycline structure indicated above, or a derivative thereof.

Some examples of antibiotic (i.e. antimicrobial) tetracy-

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.