

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SUBSYS safely and effectively. See full prescribing information for SUBSYS.

SUBSYS™ (fentanyl sublingual spray), CII  
Initial U.S. Approval: 1968

### WARNING: RISK OF RESPIRATORY DEPRESSION, MEDICATION ERRORS, ABUSE POTENTIAL

See full prescribing information for complete boxed warning.

- Due to the risk of fatal respiratory depression, SUBSYS is contraindicated in opioid non-tolerant patients (1) and in management of acute or postoperative pain, including headache/migraines. (4)
- Keep out of reach of children. (5.3)
- Use with CYP3A4 inhibitors may cause fatal respiratory depression. (7)
- When prescribing, do not convert patients on a mcg per mcg basis from any other oral transmucosal fentanyl product to SUBSYS. ( 5.1)
- When dispensing, do not substitute with any other fentanyl products. (5.1)
- Contains fentanyl, a Schedule II controlled substance with abuse liability similar to other opioid analgesics. (9.1)
- SUBSYS is available only through a restricted program called the TIRF REMS Access program. Outpatients, healthcare professionals who prescribe to outpatients, pharmacies, and distributors are required to enroll in the program. (5.10)

### INDICATIONS AND USAGE

SUBSYS is an opioid agonist indicated for the management of breakthrough pain in cancer patients 18 years of age and older who are already receiving and who are tolerant to opioid therapy for their underlying persistent cancer pain. Patients must remain on around-the-clock opioids when taking SUBSYS. (1)

Limitations of Use:

SUBSYS may be dispensed only to patients enrolled in the TIRF REMS ACCESS program.

### DOSAGE AND ADMINISTRATION

- Patients must require and use around-the-clock opioids when taking SUBSYS. (1)
- Initial dose of SUBSYS: 100 mcg.
- Individually titrate to a tolerable dose that provides adequate analgesia using a single SUBSYS dose per breakthrough cancer pain episode. (2)
- No more than two doses can be taken per breakthrough pain episode. (2.2)
- Wait at least 4 hours before treating another episode of breakthrough pain with SUBSYS. (2.3)
- Limit consumption to four or fewer doses per day once successful dose is found. (2.3)

### DOSAGE FORMS AND STRENGTHS

- Sublingual spray in 100 mcg, 200 mcg, 400 mcg, 600 mcg, 800 mcg dosage strengths. (3)

### CONTRAINDICATIONS

- Opioid non-tolerant patients. (4)
- Management of acute or postoperative pain including headache/migraine and dental pain (4)
- Intolerance or hypersensitivity to fentanyl, SUBSYS, or its components. (4)

### WARNINGS AND PRECAUTIONS

- Clinically significant respiratory and CNS depression can occur. Monitor patients accordingly. (5.1)
- Full and consumed SUBSYS units contain medicine that can be fatal to a child. Ensure proper storage and disposal. (5.3, 16.2)
- Use with other CNS depressants and moderate or strong CYP450 3A4 inhibitors may increase depressant effects including respiratory depression, hypotension, and profound sedation. Consider dosage adjustments if warranted. (5.4)
- Titrate SUBSYS cautiously in patients with chronic obstructive pulmonary disease or preexisting medical conditions predisposing them to respiratory depression and in patients susceptible to intracranial effects of CO<sub>2</sub> retention. (5.6, 5.7)

### ADVERSE REACTIONS

Most common adverse reactions during treatment (frequency ≥5%): vomiting, nausea, constipation, dyspnea, and somnolence. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Insys Therapeutics, Inc., at 1-855-978-2797 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS

- Boxed Warning and Warnings and Precautions (5.4, 7)

### USE IN SPECIFIC POPULATIONS

- Safety and effectiveness in pediatric patients below 18 years of age have not been established. (8.4)
- Administer SUBSYS with caution to patients with liver or kidney dysfunction. (8.6)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 1/2012

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**WARNING: IMPORTANCE OF PROPER PATIENT SELECTION, DOSING, AND POTENTIAL FOR ABUSE**

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Initial Dose
- 2.2 Dose Titration
- 2.3 Maintenance Dosing
- 2.4 Administration of SUBSYS
- 2.5 Discontinuation of SUBSYS

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Important Information Regarding Prescribing and Dispensing
- 5.2 Respiratory Depression
- 5.3 Patient/Caregiver Instructions
- 5.4 Additive CNS Depressant Effects
- 5.5 Effects on Ability to Drive and Use Machines
- 5.6 Chronic Pulmonary Disease
- 5.7 Head Injuries and Increased Intracranial Pressure
- 5.8 Cardiac Disease
- 5.9 MAO Inhibitors
- 5.10 Transmucosal Immediate Release Fentanyl (TIRF) Risk Evaluation and Mitigation Strategy (REMS) ACCESS Program

**6 ADVERSE REACTIONS**

- 6.1 Clinical Studies Experience

**7 DRUG INTERACTIONS**

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Patients with Renal or Hepatic Impairment
- 8.7 Gender

**9 DRUG ABUSE AND DEPENDENCE**

- 9.1 Controlled Substance
- 9.2 Abuse and Addiction
- 9.3 Dependence

**10 OVERDOSAGE**

- 10.1 Clinical Presentation
- 10.2 Immediate Management
- 10.3 Treatment of Overdosage (Accidental Ingestion) in the Opioid NON-Tolerant Person
- 10.4 Treatment of Overdose in Opioid-Tolerant Patients
- 10.5 General Considerations for Overdose

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

**16 HOW SUPPLIED/STORAGE AND HANDLING**

- 16.1 Storage and Handling
- 16.2 Disposal of SUBSYS
- 16.3 How Supplied

**17 PATIENT COUNSELING INFORMATION**

- 17.1 Patient/Caregiver Instructions
- 17.2 SUBSYS Child Safety Kit
- 17.3 Disposal of Used SUBSYS Unit Dose Systems
- 17.4 Disposal of Unopened SUBSYS Unit Dose Systems When No Longer Needed

**\*Sections or subsections omitted from the full prescribing information are not listed.**

FULL PRESCRIBING INFORMATION

**WARNING: RISK OF RESPIRATORY DEPRESSION,  
MEDICATION ERRORS, ABUSE POTENTIAL**

**Respiratory Depression**

Fatal respiratory depression has occurred in patients treated with transmucosal immediate-release fentanyl products such as SUBSYS, including following use in opioid non-tolerant patients and improper dosing. The substitution of SUBSYS for any other fentanyl product may result in fatal overdose.

Due to the risk of respiratory depression, SUBSYS is contraindicated in the management of acute or postoperative pain including headache/migraine and in opioid non-tolerant patients.

Death has been reported in children who have accidentally ingested transmucosal immediate-release fentanyl products. SUBSYS must be kept out of reach of children. [see Patient Counseling Information (17.3) and How Supplied/Storage and Handling (16.1)]

The concomitant use of SUBSYS with CYP3A4 inhibitors may result in an increase in fentanyl plasma concentrations, and may cause potentially fatal respiratory depression [see Drug Interactions (7)].

**Medication Errors**

Substantial differences exist in the pharmacokinetic profile of SUBSYS compared to other fentanyl products that result in clinically important differences in the extent of absorption of fentanyl that could result in fatal overdose.

- When prescribing, do not convert patients on a mcg per mcg basis from any other fentanyl products to SUBSYS. [see Dosage and Administration (2.1), Warnings and Precautions (5.2), and Clinical Pharmacology (12.3)]
- When dispensing, do not substitute a SUBSYS prescription for other fentanyl products.

**Abuse Potential**

SUBSYS contains fentanyl, an opioid agonist and a Schedule II controlled substance, with an abuse liability similar to other opioid analgesics. SUBSYS can be abused in a manner similar to other opioid agonists, legal or illicit. This should be considered when prescribing or dispensing SUBSYS in situations where the physician or pharmacist is concerned about an increased risk of misuse, abuse or diversion.

Because of the risk for misuse, abuse, addiction, and overdose, SUBSYS is available only through a restricted program required by the Food and Drug Administration, called a Risk Evaluation and Mitigation Strategy (REMS). Under the Transmucosal Immediate-Release Fentanyl (TIRF) REMS Access program, outpatients, healthcare professionals who prescribe to outpatients, pharmacies, and distributors must enroll in the program. [See Warnings and Precautions (5.11)] Further information is available at [www.TIRFREMSaccess.com](http://www.TIRFREMSaccess.com) or by calling 1-866-822-1483.

## 1 INDICATIONS AND USAGE

SUBSYS is indicated for the management of breakthrough pain in adult cancer patients who are already receiving and who are tolerant to around-the-clock opioid therapy for their underlying persistent cancer pain. Patients considered opioid tolerant are those who are taking around-the-clock medicine consisting of at least 60 mg of oral morphine daily, at least 25 mcg of transdermal fentanyl/hour, at least 30 mg of oral oxycodone daily, at least 8 mg of oral hydromorphone daily or an equianalgesic dose of another opioid daily for a week or longer. Patients must remain on around-the-clock opioids when taking SUBSYS.

This product **must not** be used in opioid non-tolerant patients because life-threatening respiratory depression and death could occur at any dose in patients not on a chronic regimen of opioids. For this reason, SUBSYS is contraindicated in the management of acute or postoperative pain.

SUBSYS is intended to be used only in the care of cancer patients and only by oncologists and pain specialists who are knowledgeable of and skilled in the use of Schedule II opioids to treat cancer pain.

### Limitations of Use:

As part of the Transmucosal Immediate-Release Fentanyl (TIRF) REMS ACCESS Program, SUBSYS may be dispensed only to outpatients enrolled in the program. [see *Warnings and Precautions* (5.10)]. For inpatient administration (e.g. hospitals, hospices, and long-term care facilities that prescribe for inpatient use) of SUBSYS, patient enrollment is not required.

## 2 DOSAGE AND ADMINISTRATION

Healthcare professionals who prescribe SUBSYS on an outpatient basis must enroll in the TIRF REMS ACCESS program and comply with the requirements of the REMS to ensure safe use of SUBSYS. [see *Warnings and Precautions* (5.10)]

As with all opioids, the safety of patients using such products is dependent on health care professionals prescribing them in strict conformity with their approved labeling with respect to patient selection, dosing, and proper conditions for use.

### 2.1 Initial Dose

Individually titrate SUBSYS to a dose that provides adequate analgesia and minimizes side effects. The initial dose of SUBSYS to treat episodes of breakthrough cancer pain is **always** 100 mcg. . **When prescribing, do not switch patients on a mcg per mcg basis from any other oral transmucosal fentanyl product to SUBSYS** as SUBSYS is not equivalent on a mcg per mcg basis with any other fentanyl product [see *Warnings and Precautions* (5.2) and *Clinical Pharmacology* (12.3)].

Prescribe an initial titration supply of 100 mcg SUBSYS units, which limits the number of units in the home during titration.

Avoid prescribing a higher dose until patients have used up all units to prevent confusion and possible overdose.

### 2.2 Dose Titration

- a. From the 100 mcg initial dose, closely follow patients and change the dosage level until the patient reaches a dose that provides adequate analgesia using a single SUBSYS dose per breakthrough cancer pain episode with tolerable side effects. Patients should record their use of SUBSYS over several episodes of breakthrough cancer pain and review their experience with their physicians to determine if a dosage adjustment is warranted.
- b. For each breakthrough pain episode treated, if pain is not relieved after 30 minutes, patients may take **ONLY ONE** additional dose of the same strength for that episode. Thus patients should take a maximum of two doses of SUBSYS for any breakthrough pain episode.
- c. Patients **MUST** wait **at least 4 hours** before treating another episode of breakthrough pain with SUBSYS.
- d. If there is a need to titrate to a 200 mcg dose, prescribe 200 mcg SUBSYS units.
- e. Subsequent titration steps are 400 mcg, 600 mcg, 800 mcg, 1200 mcg and 1600 mcg. See Table 1.
- f. To reduce the risk of overdose during titration, patients should have only one strength of SUBSYS available at any time.

**Table 1. Titration Steps**

<b>SUBSYS DOSE</b>	<b>Using</b>
100 mcg	1 × 100 mcg unit
200 mcg	1 × 200 mcg unit
400 mcg	1 × 400 mcg unit
600 mcg	1 × 600 mcg unit
800 mcg	1 × 800 mcg unit
1200 mcg	2 × 600 mcg unit
1600 mcg	2 × 800 mcg unit

**SUBSYS Titration Process**

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.