

Biological Psychiatry

A Journal of Psychiatric Research

Joseph Wortis, Editorial Office, Biological Psychiatry is a representation that neither the abstract or are currently under consideration or otherwise, related to the submitted words or less—no abstract necessary) words must be strictly limited.

illustrations and tables), typed double-spaced, and type corresponding author's academic or professional affiliations, etc. If the article title exceeds 45 characters, reference lists, tables, and figure legends must be mentioned with arabic numerals and no longer than 150 words.

originals or glossy copies of originals. On back of cover of "top."

them in the text by name and year in

1968): Isozyme variability in the esterases of *D. aldrichi*

vicus.

Psychophila, Vol. 6. New York:

microextraction and
Scott D (ed), *Methods in Cell*

accepted and returned within 48 hours of receipt of the original form that accompanies proofs.

Do not be asked to transfer copyright to the widest possible dissemination



HEALTH SCIENCES LIBRARY
University of Wisconsin

AUG 03 1989

1305 Linden Dr.
Madison, Wis. 53706

A PUBLICATION OF THE SOCIETY OF BIOLOGICAL PSYCHIATRY

VOLUME 26, NUMBER 4
Wockhardt 1032

AUGUST 1989
BIPCBF 26(4)1989
ISSN 0006-3223

HEALTH S
Univer

AU

EDITORIAL

13
Madi

GHB—New Hope for

Narcolepsy is a relatively rare Americans. Studies have suggested that it is characterized by sleepiness and a series of auxiliary hypnagogic hallucinations. High-dose treatment of cataplexy and other symptoms, when used in treating both types of symptoms, may be contraindicating factors (such as managing this life-long disorder).

In this issue of *Biological Psychiatry*, the authors report that gammahydroxybutyrate (GHB) treatment of narcolepsy with cataplexy significantly reduced the frequency of hypnagogic hallucinations and daytime dizziness. In their study of 20 patients, the frequency of hypnagogic hallucinations decreased by over 50% and the frequency of daytime dizziness decreased by over 50%.

Although gammahydroxybutyrate treatment of narcolepsy with cataplexy is a new development, the current study is the first demonstration of the effectiveness of GHB in narcolepsy.

Gammahydroxybutyrate is found in mammalian tissue. It is found in decreasing levels noted after treatment. High-affinity brain receptor binding, synthesis, release, and reuptake of gammahydroxybutyrate in brain tissue may be involved in hypnagogic hallucinations, daytime dizziness, and administration. Its short half-life and non-stimulant effects suggest that GHB has non-stimulant effects. It is usually administered 3-4 hr apart. The compound is often associated with a reduction in slow-wave or delta sleep. Scrima et al.'s study showed effects on delta sleep may be due to its effects on delta sleep. In our own experience with GHB, early reports suggested that it may be effective in narcolepsy. The data from the

Biological Psychiatry is cited in *Beck Medical Information*, *Biological Abstracts*, *Chemical Abstracts*, *Current Contents*, *Excerpta Medica*, *Index Medicus*, *Mental Health Abstracts*, *Psychological Abstracts*, *Referativnyi Zhurnal*, *Science Citation Index*, and *Selected List of Tables of Contents of Psychiatric Periodicals*.

Biological Psychiatry is published in two volumes mailed monthly and biweekly in January, February, March, and April by Elsevier Science Publishing Co., Inc., 655 Avenue of the Americas, New York, NY 10010. Institutional subscriptions: \$448.00; individual subscriptions: \$218.00 and \$135.00 for renewals; special in-training rate to residents, fellows, interns, and students: \$64.00; society subscriptions: \$25.00. Subscribers outside of the United States add \$56.00 for surface postage and handling. Subscriptions outside of the United States are sent by air.

Claims for missing issues can be honored only up to three months from date of issue. Duplicate copies will not be sent to replace ones undelivered through failure to notify Elsevier of change of address.

Single copy and back volume information available upon request. Please direct orders for the journal, changes of address, and claims for missing issues to: Journals Fulfillment Department, Elsevier Science Publishing Co., Inc., 655 Avenue of the Americas, New York, NY 10010.

Please direct inquiries regarding the placement of advertising in this journal to: Synergistic Media Sales, 50 East 42nd Street, Suite 504, New York, NY 10017. (212) 682-2200.

No responsibility is assumed by the Publisher or by the Society of Biological Psychiatry for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein. No suggested test or procedure should be carried out unless, in the reader's judgment, its risk is justified. Because of rapid advances in the medical sciences, we recommend that the independent verification of diagnoses and drug dosages should be made. Discussions, views and recommendations as to medical procedures, choice of drugs and drug dosages are the responsibility of the authors.

© 1989 Society of Biological Psychiatry. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal or internal use of specific clients. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotional purposes, or for creating new collective works. For such purposes, consent will be given on the condition that the copier pay through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by the U.S. Copyright Law. If no code appears on an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author.

Second-class postage paid at New York, NY and additional mailing offices. Postmaster: send address changes to *Biological Psychiatry*, Elsevier Science Publishing Co., Inc., 655 Avenue of the Americas, New York, NY 10010.

Manufactured in the U.S.A.

HB seem to require lower levels of GHB. Conjecture is that any decrease in nighttime sleep. As patients with sleep, they may experience more GHB.

Standard pharmacotherapies, the 13 narcoleptic patients to date has shown that, laboratory determination of blood constituents obtained in 3-years of patient experience with treated abnormalities.

The pharmacotherapy of narcolepsy, alcohol and codeine in the management of antidepressants, which recently in symptoms of the narcoleptic type is usually a predictable and progressive cataplexy manifested by a marked attack. Although cataplexy is ordinarily cataplexy can result in cataplexies attacks.

Essential agent to provide relief of associated with REM rebound, and in any of our patients who have drug, which is thus far available. One of a series of studies necessary to spread availability as a possible cure of narcolepsy. Its easier access to REM-related mechanisms and its clinical utility that extends far

Martin B. Scharf
Kathleen A. Fletcher

gamma-hydroxybutyrate on sleep. *Biol Psychiatry* 1985;22:100-105.

(1985): The effects and effectiveness of gamma-hydroxybutyrate on sleep. *Am J Psychiatry* 142:100-105, 1985.

Efficacy of gamma-hydroxybutyrate versus placebo in treating narcolepsy-cataplexy: Double-blind subjective measures. *Biol Psychiatry* 1989;26:331-343.

Clin Neurophysiol 7:75-89.

Efficacy of Gamma-Hydroxybutyrate versus Placebo in Treating Narcolepsy-Cataplexy: Double-Blind Subjective Measures

Lawrence Scrima, Paul G. Hartman, Frank H. Johnson, Jr., and F. Charles Hiller

The efficacy of gamma-hydroxybutyrate (GHB) versus placebo for treating narcolepsy was evaluated in 20 patients with narcolepsy, 10 men and 10 women, using a double-blind counterbalanced crossover design. Each patient completed a daily sleep-wake log and questionnaire during a 14-day baseline, a 29-day placebo period, a 29-day GHB period (50 mg GHB/kg/night given 25 mg/kg h.s. and 25 mg/kg 3 hr later), and a 6-day washout period after each treatment. Cataplexy frequency was significantly lower during GHB treatment than during placebo treatment ($p = 0.022$). Compared to baseline values, the number of cataplexy attacks per day declined by 52% and 69% during GHB treatment weeks 1 and 4, respectively. The number of subjective arousals from sleep was less with GHB than with placebo ($p = 0.035$), and the number of sleep attacks was not significantly different during GHB versus placebo treatment. GHB did not have a significant effect on subjective estimates of sleep onset latency, total sleep time, Stanford Sleepiness Scale ratings at morning wake-up, methylphenidate usage, or the number of naps per day. The results indicate that GHB is efficacious for reducing the frequency of cataplexy attacks and subjective nocturnal arousals in patients with narcolepsy within the first 4 weeks of treatment.

Introduction

Narcolepsy is a chronic, incurable disorder characterized by intermittent excessive daytime sleepiness and abnormal rapid eye movement (REM) sleep manifestations, such as sleep-onset REM periods, cataplexy, sleep paralysis, and/or hypnagogic hallucinations (Association of Sleep Disorders Centers 1979). Cataplexy is a sudden loss of muscle tone that occurs primarily during emotional arousal; sleep paralysis is an inability to move upon first-lying down or upon waking; and hypnagogic hallucinations are dream-like hallucinations that occur at sleep onset. Most patients with narcolepsy also have disrupted nocturnal sleep (Montplaisir 1976). Narcolepsy is generally treated with a central nervous system stimulant

From the Sleep Disorders Center, Pulmonary Division, Department of Medicine, University of Arkansas for Medical Sciences, Little Rock, AR.

Supported in part by Orphan Products Grant FD-R-000115 from the Food and Drug Administration.

Address reprint requests to Dr. L. Scrima, Sleep Disorders Center, Slot 594, University of Arkansas for Medical Sciences, 4301 West Markham Street, Little Rock, AR 72205.

Received July 29, 1988; revised December 14, 1988.

(e.g., amphetamine, methylphenidate, or pemoline) to reduce excessive daytime sleepiness (EDS) (Parkes 1976) and an antidepressant (e.g., imipramine or protriptyline) to control cataplexy and other REM sleep-related symptoms (Takahashi 1976). Stimulants, however, do not fully control EDS in most narcolepsy patients (Parkes 1976) and can have undesirable side effects on the cardiovascular, gastrointestinal, and central nervous systems (Gilman et al. 1985). Treatment of cataplexy with antidepressants has been reported to be successful in most cases (Takahashi 1976; Billiard et al. 1983), but these drugs have adverse effects, including (1) prolonged cardiac conduction times that may promote dangerous ventricular arrhythmias; (2) postural hypotension; (3) anticholinergic effects, such as blurred vision, dry mouth, and impotence (Gilman et al. 1985); (4) suppression of REM sleep (Zung 1969; Cadilhac 1976); and (5) increased nocturnal myoclonus (Guilleminault et al. 1976). The anticholinergic side effects often result in patient self-withdrawal from antidepressants, which is usually followed by an increase in the frequency and severity of cataplexy events (Scrima 1981; Scharf and Fletcher 1988). Both stimulants and anticataplexy drugs may become less effective as tolerance increases (Parkes 1976; Broughton and Mamelak 1979).

Gamma-hydroxybutyrate (GHB) is a four-carbon fatty acid that occurs naturally in the mammalian central nervous system (Muyard and Laborit 1977) and has been termed a "putative neurotransmitter" (Mandel et al. 1987). GHB was reported to induce anesthesia at 60-70 mg/kg (Vickers 1969), but the report does not make it clear whether the doses were given orally or intravenously. Lower oral doses of GHB were reported to induce sleep in psychiatric patients (Mamelak et al. 1977), but the minimum GHB dose that will induce sleep has not been systematically determined. Unlike other hypnotics, GHB given orally induces and maintains sleep without suppressing REM or delta stages of sleep (Mamelak et al. 1977). It was first reported in 1976 (Broughton and Mamelak) that GHB, given orally h.s. and two to three additional times during the sleep period, improved nighttime sleep and reduced cataplexy and sleep attacks in patients with narcolepsy. Subsequent studies confirmed that most narcolepsy patients had moderate to large reductions in cataplexy frequency and daytime sleepiness, as well as reduced sleep disruption, hypnagogic hallucinations, and sleep paralysis after taking GHB in divided dose, i.e., a dose h.s. and one to two additional times during the night (Broughton and Mamelak 1979, 1980; Scharf et al. 1985). Polysomnographic recordings indicated that narcolepsy patients taking a divided dose of GHB had increased sleep continuity, decreased REM fragmentation, and increased amounts of delta sleep (Broughton and Mamelak 1980; Scharf et al. 1985). However, 1 month of oral administration of a single h.s. dose of GHB improved daytime sleepiness in only 39% of patients with narcolepsy, though cataplexy frequency was reduced in 83% of the patients (Montplaisir and Godbout 1986). Tolerance to GHB has not been found to develop, even after daily use by patients with narcolepsy for as long as 9 years (Mamelak et al. 1986). Adverse side effects have been infrequent, mild, and have occurred mainly during the first few days of treatment (Broughton and Mamelak 1979; Scharf et al. 1985; Mamelak et al. 1986).

This report describes the results of the first double-blind study of the effects of GHB on subjective symptoms of narcolepsy as compared to those of a placebo.

Methods

Subjects

Ten women and 10 men with narcolepsy, diagnosed at the accredited Sleep Disorders Center (SDC) of the University of Arkansas for Medical Sciences (UAMS), participated

in the study. All patients were given a physical examination (Guilleminault 1982) (PSG) and an evaluation of sleepiness (MSLT). The criteria for inclusion were (1) sleepiness, ≥ 75 on the MSLT, (2) at least 2 weeks of baseline sleep without cataplexy, (3) excluded if they had other sleep disorders, with the exception of sleep paralysis, mild to moderate nocturnal myoclonus.

The age mean \pm standard deviation was 49.1 ± 4.0 (21-64) years for men and 45.1 ± 2.5 (17.6-45.4) for women. All patients were on stimulant medication (11 were on a combination of amphetamine and protriptyline), and 2 patients were on methylphenidate (≤ 30 mg/day). Patients who were taking medication for cataplexy were excluded. Patients remaining 4 patients on a combination of amphetamine and protriptyline, 6, 5, and 3 days prior to baseline at the first 2 days of the baseline period, 5.7 days prior to baseline at the first 2 days of the baseline period: 5.7 days prior to baseline at the first 2 days of the baseline period. One patient continued taking amphetamine and protriptyline a dose (40 mg/day) that was reduced to reduce narcolepsy symptoms (Scharf et al. 1985). This patient was included in the study. This patient excluded yield

Procedures

A double-blind, crossover study was utilized. The study was conducted at the UAMS pharmacy, so that the first treatment period and placebo first and GHB second treatment period. Each treatment period contained 25 mg of GHB. During the placebo treatment period, subjects received an equivalent amount of fluid during the treatment period, subjects

duce excessive daytime sleepiness (amphetamine or protriptyline) to control (Kashi 1976). Stimulants, however, (Kashi 1976) and can have undesirable effects on central nervous systems (Gilman et al. 1976). It has been reported to be successful in the treatment of these drugs have adverse effects, they may promote dangerous ventricular arrhythmias, such as blurred vision, depression of REM sleep (Zung 1969; Guillemineault et al. 1976). The withdrawal from antidepressants, which the severity of cataplexy events (Scrima et al. 1979). Anticatataplexy drugs may become less effective (Mamelak 1979).

Gamma-hydroxybutyrate (GHB) is a naturally occurring amino acid that occurs naturally in the brain (Broughton 1977) and has been termed a "natural hypnotic" (Broughton 1977). It was reported to induce anesthesia in patients with narcolepsy. It is not clear whether the doses of GHB were reported to induce anesthesia without the minimum GHB dose that was reported. Unlike other hypnotics, GHB does not suppress REM or delta stages of sleep (Broughton and Mamelak 1976). During the sleep period, improved sleep continuity in patients with narcolepsy. Patients had moderate to large reductions in sleep latency, as well as reduced sleep disturbance after taking GHB in divided doses during the night (Broughton and Mamelak 1976). Recordings indicated that narcolepsy symptoms improved, decreased REM sleep (Broughton and Mamelak 1980). Administration of a single h.s. dose of GHB to patients with narcolepsy, though (Montplaisir and Godbout 1986). Improvement in sleep after daily use by patients with narcolepsy. Adverse side-effects have been reported during the first few days of treatment (Broughton et al. 1986).

A double-blind study of the effects of GHB compared those of a placebo.

Patients were recruited from the accredited Sleep Disorders and Narcolepsy Clinic at the University of Sciences (UAMS), participated

in the study. All patients were interviewed by an accredited clinical polysomnographer, were given a physical examination by a physician, and had sleep disorders diagnostic tests (Guilleminault 1982). The diagnostic tests included an overnight polysomnogram (PSG) and an evaluation of their daytime sleepiness with the multiple sleep latency test (MSLT). The criteria for inclusion in this study were: (1) a history of excessive daytime sleepiness and cataplexy, (2) ≥ 2 REM onsets on the MSLT, (3) a sleepiness index of ≥ 75 on the MSLT, (4) at least 10 cataplexy attacks subjectively reported on a daily log during a 2-week baseline period, and (5) age between 16 and 65 years. Patients were excluded if they had other major health problems; were fertile women who were not practicing birth control; were nursing mothers; or had previously taken GHB or had other sleep disorders, with the exception of those commonly associated with narcolepsy, such as sleep paralysis, mild to moderate sleep apnea (arterial oxygen saturation $\geq 80\%$), and nocturnal myoclonus.

The age mean \pm standard error was 45.9 ± 4.6 years (range 16-64) for the women and 49.1 ± 4.0 (21-64) for the men. Weight (kg) mean \pm SE was 85.1 ± 5.2 (range 57-113) for women and 80.4 ± 3.6 (54-90) for men. Body mass index was 31.8 ± 2.5 (17.6-45.4) for women and 26.2 ± 0.9 (20.3-29.1) for men. Prior to the study, 7 patients were on stimulants alone (methylphenidate, pemoline, or dextroamphetamine), 11 were on a combination of stimulants and anticatataplexy medications (imipramine or protriptyline), and 2 patients were not taking stimulants or anticatataplexy medications. Patients who were taking stimulants other than methylphenidate switched to methylphenidate (≤ 30 mg/day) for the duration of the study. Seven patients were withdrawn from anticatataplexy medications at least 2 weeks before the baseline period, and the remaining 4 patients on anticatataplexy medications were withdrawn from imipramine 6, 6, 5, and 3 days prior to baseline, respectively. Only the patient who stopped imipramine 5 days prior to baseline appeared to have elevated amounts of cataplexy events during the first 2 days of the baseline (18 and 12 events, respectively; mean and SD for the rest of the baseline period: 5.7 ± 3.5 events); these 2 days were excluded from the analysis. One patient continued taking propranolol to control hypertension throughout the study at a dose (40 mg/day) that was half the lowest dose of propranolol (80-480 mg/day) reported to reduce narcolepsy symptoms for some patients (Kales et al. 1979; Meier-Ewert et al. 1985). This patient was included in the analysis of results, but analysis of the data with this patient excluded yielded the same pattern of statistically significant results.

Procedures

A double-blind, crossover design, with order of treatment counterbalanced and randomly assigned, was utilized. Thus, each subject provided data for all phases of the study: baseline (14 days), first treatment (29 days), first washout (6 days), second treatment (29 days), and second washout (6 days). Order of treatment was randomly assigned by the UAMS pharmacy, so that half of the men and half of the women received GHB in the first treatment period and placebo in the second, whereas the remaining subjects received placebo first and GHB second. All SDC staff were blind to the order of treatment for subjects. During GHB treatment, subjects received 58 bottles prepared by the pharmacy, each containing 25 mg GHB/kg body weight, mixed with distilled water and syrup of orange. During the placebo treatment, subjects received 58 identical bottles with an equivalent amount of fluid, consisting of syrup of orange in distilled water. During each treatment period, subjects were instructed orally and in writing to (1) refrigerate, but not

Wockhardt 1032

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.