

346



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A01N 43/40, A61K 31/445</b>	<b>A1</b>	(11) International Publication Number: <b>WO 98/31225</b> (43) International Publication Date: 23 July 1998 (23.07.98)
<p>(21) International Application Number: PCT/US98/00618</p> <p>(22) International Filing Date: 13 January 1998 (13.01.98)</p> <p>(30) Priority Data: 60/035,591 17 January 1997 (17.01.97) US</p> <p>(71) Applicant: BRISTOL-MYERS SQUIBB COMPANY [US/US]; P.O. Box 4000, Princeton, NJ 98543-4000 (US).</p> <p>(72) Inventor: GREGG, Richard, E.; 7 Linden Lane, Pennington, NJ 08534 (US).</p> <p>(74) Agents: RODNEY, Burton et al.; Bristol-Myers Squibb Company, P.O. Box 4000, Princeton, NJ 08543-4000 (US).</p>	<p>(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report.</i></p>	
(54) Title: A METHOD OF INHIBITING OR TREATING PHYTOSTEROLEMIA WITH AN MTP INHIBITOR		
(57) Abstract		
<p>A method is provided for inhibiting onset of or treating phytosterolemia by administering to a patient an MTP inhibitor, alone or optionally, in combination with another cholesterol lowering drug, such as pravastatin.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
DJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## A METHOD OF INHIBITING OR TREATING PHYTOSTEROLEMIA WITH AN MTP INHIBITOR

5

Field of the Invention

The present invention related to a method for inhibiting onset of or treating phytosterolemia, by administering an MTP inhibitor alone or in combination with another cholesterol lowering drug, such as pravastatin.

10

Background of the Invention

As indicated in Scriver, C.R. et al "Metabolic Basis of Inherited Diseases" Vol. II (1995), Chap. 65, Inborn Errors in Bile Acid Biosynthesis and Storage of Sterols Other than Cholesterol, Bjorkhem, I. and Boberg, K.M., pp. 2073-2099, phytosterolemia (also referred to as sitosterolemia) is a rare inherited sterol storage disease involving increased intestinal absorption of phytosterol or shellfish sterols and decreased fecal secretion. It is characterized by "tendon and tuberous xanthomas and by a strong predisposition to premature coronary atherosclerosis.... Increased amounts of phytosterols (plant sterols), such as sitosterol and campesterol and their 5 $\alpha$ -stanols, are found in blood, plasma, erythrocytes, and different tissues, especially in the xanthomas and arteries of affected subjects. Increased serum cholesterol and cholesterol have also been found in many patients." (p. 2073)

Patients afflicted with phytosterolemia have been found to have an increased incidence of coronary heart disease at an early age most likely due to early development of atherosclerosis at an early age. Bjorkhem et al, supra, indicate at page 2090 that "the mechanism behind the atherosclerosis is unexplained, but a high content of plant sterols in the circulating lipoproteins might promote their deposition in the arterial wall."

The microsomal triglyceride transfer protein (MTP) catalyzes the transport of triglyceride (TG), cholesteryl ester (CE), and phosphatidylcholine (PC) between small unilamellar vesicles (SUV). Wetterau & Zilversmit, Chem. Phys. Lipids 38, 205-22 (1985). When transfer rates are expressed as the percent of the donor lipid transferred per time, MTP expresses a distinct preference for neutral lipid transport (TG and CE), relative to phospholipid transport. The microsomal triglyceride transfer protein from bovine liver has been isolated and extensively characterized (1). This has led to the cloning of cDNA expressing the protein from several species, including humans (2). MTP is composed of two subunits. The small subunit is the previously characterized multifunctional protein, protein disulfide isomerase. This is supported by biochemical analysis of the protein (3) as well as co-expression studies performed in insect Sf9 cells using the baculovirus expression system. Expression of soluble active MTP requires the co-expression of PDI and the unique large subunit of MTP (4).

1: Wetterau, J.R. and Zilversmit, D.B. (1985) Chem. Phys. Lipids 38, 205-222.

Wetterau, J.R., et al, (1990) J. Biol. Chem. 265, 9800-9807.

Wetterau, J.R., et al, (1991) Biochemistry 30, 4406-4412.

Atzel, A., and Wetterau, J.R. (1993) Biochemistry 32, 10444-10450.

Atzel, A., and Wetterau, J.R. (1994) Biochemistry 33, 15382-15388.

Jamil, H., et al, (1995) J. Biol. Chem. 270, 6549-6554.

2. Sharp, D. et al, (1993) Nature 365, 65-69.

Lin, M.C.M., et al, J. Biol. Chem. 269, 29138-29145.

Nakamuta, M., et al, (1996) Genomics 33, 313-316.

3. Wetterau, J.R., et al, (1990) J. Biol. Chem. 265, 9800-9807.

5 Wetterau, J.R., et al, (1991) Biochemistry 30, 9728-9735.

4. Ricci, B., et al, (1995) J. Biol. Chem. 270, 14281-14285.

10 In vitro, MTP catalyzes the transport of lipid molecules between phospholipid membranes. Presumably, it plays a similar role in vivo, and thus plays some role in lipid metabolism. The subcellular (lumen of the microsomal fraction) and tissue distribution (liver and intestine) of  
15 MTP have led to speculation that it plays a role in the assembly of plasma lipoproteins, as these are the sites of plasma lipoprotein assembly. Wetterau & Zilversmit, Biochem. Biophys. Acta 875, 610-7 (1986). The ability of  
20 MTP to catalyze the transport of TG between membranes is consistent with this hypothesis, and suggests that MTP may catalyze the transport of TG from its site of synthesis in the endoplasmic reticulum (ER) membrane to nascent lipoprotein particles within the lumen of the ER.

Abetalipoproteinemia is an autosomal recessive  
25 disease characterized by a virtual absence of plasma lipoproteins which contain apolipoprotein B (apoB). Kane & Havel in The Metabolic Basis of Inherited Disease, Sixth edition, 1139-64 (1989). Plasma TG levels may be as low as a few mg/dL, and they fail to rise after fat ingestion.  
30 Plasma cholesterol levels are often only 20-45 mg/dL. These abnormalities are the result of a genetic defect in the assembly and/or secretion of very low density lipoproteins (VLDL) in the liver and chylomicrons in the intestine. The molecular basis for this defect had not  
35 been previously determined. In subjects examined, triglyceride, phospholipid, and cholesterol synthesis appear normal. At autopsy, subjects are free of

# Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

## Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

## Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

## Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

## API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

## LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

## FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

## E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.