

4311



Consommation et
Affaires commerciales Canada

Consumer and
Corporate Affairs Canada

Bureau des brevets

Patent Office

Ottawa, Canada
K1A 0C9

(21)	(A1)	2,091,102
(22)		1993/03/05
(43)		1993/09/07

(51) INTL.CL. ⁵ C12N-015/12; C12Q-001/68; C07K-013/00; C07K-015/12;
C07H-021/04; G01N-001/68

(19) (CA) APPLICATION FOR CANADIAN PATENT (12)

(54) Microsomal Triglyceride Transfer Protein

(72) Wetterau, John R., II - ;
Sharp, Daru Y. - ;
Gregg, Richard E. - ;

(73) Same as inventor

(30) (US) 847,503 1992/03/06

(57) 32 Claims

Notice: This application is as filed and may therefore contain an
incomplete specification.

Canada

CCA 3764 (10-07) 01 P.30 P.1-308-3764

2091102

DC21a

Abstract

MICROSOMAL TRIGLYCERIDE TRANSFER PROTEIN

Nucleic acid sequences, particularly DNA sequences, coding for all or part of the high molecular weight subunit of microsomal triglyceride transfer protein, expression vectors containing the DNA sequences, host cells containing the expression vectors, and methods utilizing these materials. The invention also concerns polypeptide molecules comprising all or part of the high molecular weight subunit of microsomal triglyceride transfer protein, and methods for producing these polypeptide molecules. The invention additionally concerns novel methods for preventing, stabilizing or causing regression of atherosclerosis and therapeutic agents having such activity. The invention additionally concerns novel methods for lowering serum lipid levels and therapeutic agents having such activity.

MICROSOMAL TRIGLYCERIDE TRANSFER PROTEIN

5 **Cross-Reference to Related Application**

This is a continuation-in-part of U. S. patent application Ser. No. 847, 503, filed March 6, 1992.

10 **Field of the Invention**

This invention relates to microsomal triglyceride transfer protein, genes for the protein, expression vectors comprising the genes, host cells comprising the vectors, methods for producing the protein, methods for detecting inhibitors of the protein, and methods of using the protein and/or its inhibitors.

15 **Background of the Invention**

The microsomal triglyceride transfer protein (MTP) catalyzes the transport of triglyceride (TG), cholesteryl ester (CE), and phosphatidylcholine (PC) between small unilamellar vesicles (SUV). Wetterau & Zilversmit, Chem. Phys. Lipids **38**, 205-22 (1985). When transfer rates are expressed as the percent of the donor lipid transferred per time, MTP expresses a distinct preference for neutral lipid transport (TG and CE), relative to phospholipid transport. The protein from bovine liver has been isolated and characterized. Wetterau & Zilversmit, Chem. Phys. Lipids **38**, 205-22 (1985). Polyacrylamide gel electrophoresis (PAGE) analysis of the purified protein suggests that the transfer protein is a complex of two subunits of apparent molecular weights 58,000 and 88,000, since a single band was present when
25 purified MTP was electrophoresed under nondenaturing condition,
30

while two bands of apparent molecular weights 58,000 and 88,000 were identified when electrophoresis was performed in the presence of sodium dodecyl sulfate (SDS). These two polypeptides are hereinafter referred to as 58 kDa and 88 kDa, respectively, or the 58 kDa and the 88 kDa component of MTP, respectively, or the low molecular weight subunit and the high molecular weight subunit of MTP, respectively.

Characterization of the 58,000 molecular weight component of bovine MTP indicates that it is the previously characterized multifunctional protein, protein disulfide isomerase (PDI). Wetterau *et al.*, *J. Biol. Chem.* 265, 9800-7 (1990). The presence of PDI in the transfer protein is supported by evidence showing that (1) the amino terminal 25 amino acids of the bovine 58,000 kDa component of MTP is identical to that of bovine PDI, and (2) disulfide isomerase activity was expressed by bovine MTP following the dissociation of the 58 kDa - 88 kDa protein complex. In addition, antibodies raised against bovine PDI, a protein which by itself has no TG transfer activity, were able to immunoprecipitate bovine TG transfer activity from a solution containing purified bovine MTP.

PDI normally plays a role in the folding and assembly of newly synthesized disulfide bonded proteins within the lumen of the endoplasmic reticulum. Bulleid & Freedman, *Nature* 335, 649-51 (1988). It catalyzes the proper pairing of cysteine residues into disulfide bonds, thus catalyzing the proper folding of disulfide bonded proteins. In addition, PDI has been reported to be identical to the beta subunit of human prolyl 4-hydroxylase. Koivu *et al.*, *J. Biol. Chem.* 262, 6447-9 (1987). The role of PDI in the bovine transfer protein is not clear. It does appear to be an essential component of the transfer protein as dissociation of PDI from the 88 kDa component of bovine MTP by either low concentrations of a denaturant (guanidine HCl), a chaotropic agent (sodium perchlorate), or a non-denaturing detergent (octyl glucoside) results in a loss of transfer activity. Wetterau *et al.*,

Biochemistry 30, 9728-35 (1991). Isolated bovine PDI has no apparent lipid transfer activity, suggesting that either the 88 kDa polypeptide is the transfer protein or that it confers transfer activity to the protein complex.

5 The tissue and subcellular distribution of MTP activity in rats has been investigated. Wetterau & Zilversmit, Biochem. Biophys. Acta 875, 610-7 (1986). Lipid transfer activity was found in liver and intestine. Little or no transfer activity was found in plasma, brain, heart, or kidney. Within the liver, MTP was a soluble protein
10 located within the lumen of the microsomal fraction. Approximately equal concentrations were found in the smooth and rough microsomes.

 Abetalipoproteinemia is an autosomal recessive disease characterized by a virtual absence of plasma lipoproteins which
15 contain apolipoprotein B (apoB). Kane & Havel in The Metabolic Basis of Inherited Disease, Sixth edition, 1139-64 (1989). Plasma TG levels may be as low as a few mg/dL, and they fail to rise after fat ingestion. Plasma cholesterol levels are often only 20-45 mg/dL. These abnormalities are the result of a genetic defect in
20 the assembly and/or secretion of very low density lipoproteins (VLDL) in the liver and chylomicrons in the intestine. The molecular basis for this defect has not been previously determined. In subjects examined, triglyceride, phospholipid, and cholesterol synthesis appear normal. At autopsy, subjects are free
25 of atherosclerosis. Schaefer et al., Clin. Chem. 34, 89-12 (1988). A link between the apoB gene and abetalipoproteinemia has been excluded in several families. Talmud et al., J. Clin. Invest. 82, 1803-6 (1988) and Huang et al., Am. J. Hum. Genet. 46, 1141-8 (1990).

30 Subjects with abetalipoproteinemia are afflicted with numerous maladies. Kane & Havel, supra. Subjects have fat malabsorption and TG accumulation in their enterocytes and hepatocytes. Due to the absence of TG-rich plasma lipoproteins, there is a defect in the transport of fat-soluble vitamins such as

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.