



US005952498A

United States Patent [19]

[11] **Patent Number:** 5,952,498

Lenfers et al.

[45] **Date of Patent:** *Sep. 14, 1999

[54] **PROCESS FOR THE PREPARATION OF ENANTIOMERICALLY PURE CYCLOALKANO-INDOL -AND AZAINDOL -AND PYRIMIDO [1,2A] INDOLCARBOXYCLIC ACIDS AND THEIR ACTIVATED DERIVATIVES**

[75] Inventors: **Jan-Bernd Lenfers; Peter Fey; Paul Naab**, all of Wuppertal; **Kai Van Laak**, Köln, all of Germany

[73] Assignee: **Bayer Aktiengesellschaft**, Leverkusen, Germany

[*] Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

[21] Appl. No.: **08/829,566**

[22] Filed: **Mar. 31, 1997**

[30] **Foreign Application Priority Data**

Apr. 4, 1996 [DE] Germany 196 13 549

[51] **Int. Cl.⁶** **C07D 471/04**; C07D 487/04; C07C 69/616; C07C 69/612

[52] **U.S. Cl.** **544/252**; 544/249; 546/84; 546/85; 546/86; 546/87; 548/400; 548/416; 548/427; 514/267; 514/292

[58] **Field of Search** 544/249, 252; 514/267, 292, 312; 546/84-87; 548/427

[56] **References Cited**

U.S. PATENT DOCUMENTS

4,970,215 11/1990 Mohrs et al. 514/311
5,527,809 6/1996 Müller-Gliemann et al. 514/303
5,684,014 11/1997 Müller et al. 514/292

FOREIGN PATENT DOCUMENTS

344519 11/1990 European Pat. Off. .
513533 11/1992 European Pat. Off. .
560163 9/1993 European Pat. Off. .
610698 8/1994 European Pat. Off. .
705831 4/1996 European Pat. Off. .

Primary Examiner—Jose' G. Dees

Assistant Examiner—Sabiha N. Qazi

Attorney, Agent, or Firm—Sprung Kramer Schacfer & Briscoe

[57] **ABSTRACT**

The invention relates to a process and intermediates for the preparation of enantiomerically pure cycloalkanoindolecarboxylic acids and azaindolecarboxylic acids and pyrimido [1,2a]indolecarboxylic acids and their activated derivatives, characterized in that the tolyl acetic acid is first esterified with a chiral alcohol, then diastereoselective substitution at α -carbon atoms is carried out and this product is halogenated in the tolyl group and then reacted with appropriate cycloalkanoindoles, cycloalkanoazaindoles or pyrimido [1, 2a]indoles. It is possible by this method to prepare specifically, in high purity, the enantiomerically pure carboxylic acids which are intermediates for valuable medications.

10 Claims, No Drawings

1

**PROCESS FOR THE PREPARATION OF
ENANTIOMERICALLY PURE
CYCLOALKANO-INDOL -AND AZAINDOL
-AND PYRIMIDO [1,2A]
INDOLCARBOXYCYCLIC ACIDS AND THEIR
ACTIVATED DERIVATIVES**

The invention relates to a process for the preparation of enantiomerically pure cycloalkano-indolecarboxylic acids and azaindolecarboxylic acids and pyrimido[1,2a]


indolecarboxylic acids and their activated derivatives, which represent important intermediates for the synthesis of anti-atherosclerotically active cycloalkanoindole derivatives and azaindole derivatives and pyrimido[1,2a]indole derivatives.

It is known that enantiomerically pure cycloalkano-indolecarboxylic acids and azaindole-carboxylic acids and their activated derivatives can be separated into the corresponding enantiomers by diastereomeric separation by conventional methods, for example by chromatography or fractional crystallization.

This process has a number of disadvantages: both the chromatographic diastereomeric separation and the fractional crystallization of the diastereomers are associated with high equipment requirements. In addition, in this case, generally 50% of the "wrong" diastereomer arises, which can no longer be recycled to the original preparation process.


This 50% loss of yield considerably impairs the economic efficiency of a (large) industrial-scale process, quite apart from the fact that 50% of "by-product" must be disposed of. Furthermore, the customary chiral auxiliary reagents are generally very expensive even in small amounts and can then usually only be prepared via a complex synthetic pathway.

It has now been found that enantiomerically pure cycloalkano-indolecarboxylic acids and azaindolecarboxylic acids and pyrimido[1,2a]indole-carboxylic acids and their activated derivatives of the general formula (I)




in which

A represents a radical of the formula



or




J, D, E, G, L and M are identical or different and denote hydrogen, halogen, trifluoromethyl, carboxyl,

2

hydroxyl, linear or branched alkoxy or alkoxycarbonyl each having up to 6 carbon atoms, or linear or branched alkyl having up to 6 carbon atoms, which itself can be substituted by hydroxyl or by linear or branched alkoxy having up to 4 carbon atoms,

in which

R¹ and R², including the double bond linking them, together form a phenyl ring or pyridyl ring or a ring of the formula



where

R⁵ denotes hydrogen or linear or branched alkyl having up to 4 carbon atoms,


R³ and R⁴, including the double bond linking them, together form a phenyl ring or a 4- to 8-membered cycloalkene or oxocycloalkene radical, where all the ring systems listed under R¹/R² and R³/R⁴ are optionally up to trisubstituted identically or differently by halogen, trifluoromethyl, carboxyl, hydroxyl, by linear or branched alkoxy or alkoxycarbonyl each having up to 6 carbon atoms, or by linear or branched alkyl having up to 6 carbon atoms, which itself can be substituted by hydroxyl or by linear or branched alkoxy having up to 4 carbon atoms,

T represents cycloalkyl having 4 to 12 carbon atoms, or represents linear or branched alkyl having up to 12 carbon atoms,

Q represents hydroxyl or an activating radical, and their salts are obtained

by firstly converting compounds of the general formula (II),

(II)



in which

R⁶ together with the oxygen atom represents a chiral alcohol radical, by means of compounds of the general formula (III)

T-Z

(III)


in which

T has the meaning specified and

Z represents a typical leaving group, such as bromine, chlorine, iodine, mesyl, tosyl, or trifluoromethylsulphonyl, preferably iodine or bromine,


in inert solvents in the presence of a base by diastereoselective alkylation into the enantiomerically pure compounds of the general formula (IV)

3



in which

T and R⁶ have the meaning specified, then converting these, by halogenation, into the enantiomerically pure compounds of the general formula (V)



in which


T and R⁶ have the meaning specified and R⁷ represents halogen, such as chlorine, bromine, iodine, preferably bromine, reacting these in a further step with compounds of the general formula (VI)

A-H

(VI)


in which

R¹, R², R³ and R⁴ have the meaning specified, to give the enantiomerically pure compounds of the general formula (VII)



4

(IV)



5

in which

10

A, T and R⁶ have the meaning specified, and, in the case of compounds of the general formula (I) where Q=OH, carrying out a hydrolysis, and in the case where Q=activating radical, starting from the enantiomerically pure acids reacting with activating reagents.


15

These can be reacted in a further step with D- or L-phenylglycinol to give compounds of the general formula (VIII)

20

(VIII)

25



30


where these are in this case active compounds for medicaments.

The process according to the invention can be described by way of example by the following formula diagram:

5


6

-continued



7

8



Surprisingly, the process according to the invention gives the wanted enantiomerically pure cycloalkano-indolecarboxylic acids and azaindole-carboxylic acids and pyrimido-indolecarboxylic acids and their activated derivatives with-
out great equipment requirements in very good yields and high purity.

Depending on the configuration of the radical R^6 and steric effects of the alkyl halide (II) used, the alkylation of the compound (II) proceeds in high yields and in a simple manner diastereoselectively for the first time. The compounds (IV) arise with high diastereomeric excess and crystallize out of the reaction mixture directly, as a result of which even the simple crystallization of crude products gives the compounds of the formula (IV) in diastereomerically pure form.

A further advantage of the process according to the invention is that, by suitable choice of the solvent and a base, the unwanted diastereomer can be epimerized to the desired diastereomer, which in turn crystallizes out directly. Thus, further (wanted) diastereomerically pure product can be produced from the mother liquors by repeated epimerization and crystallization. Direct addition of the mother liquors to the alkylation step can optimize the entire process in the form of a cyclic process.

A further advantage of the process according to the invention is that the halogenated compounds of the general formula (V) surprisingly react with the compounds of the general formula (VI) without racemization at the carbon atom in the 2 position to the carboxylic acid function, to give the compounds of the general formula (VII).

A further advantage of the process according to the invention is the racemization-free reaction at the carbon atom at the 2 position to the carboxylic acid function of the compounds of the general formula (I) where Q=activated radical, preferably chlorine, to give the compounds of the general formula (VIII).

Furthermore, it is a great advantage of the process according to the invention that the starting compounds are very readily accessible. They may be prepared in good yields from relatively simple building blocks with low equipment requirements. Furthermore, the process according to the invention enables amounts of known racemates of the compounds of the general formula (I) present to be converted into the corresponding enantiomers. The process according to the invention enables the preparation of the compounds according to the invention of the general formula (I) using few synthetic stages and in a considerably higher overall yield than by processes known from the prior art.


R^6 , in the context of the above specified definition, represents a chiral alcohol radical, such as (+)- or (-)-

menthyl, (+)- or (-)-bornyl, (+)- or (-)-isobornyl or (-)-8-phenylmenthyl. Preferably, R^6 represents (+)- or (-)-menthyl.

Activating radicals (Q), in the context of the invention, generally represent chloride, bromide, mesylate, tosylate or triflate. Preference is given to chloride.

Preferably, by the process according to the invention, compounds of the general formula (I) are prepared, in which


A represents a radical of the formula



in which

J, D, E, G, L and M are identical or different and denote hydrogen, fluorine, chlorine, bromine trifluoromethyl, carboxyl, hydroxyl, linear or branched alkoxy or alkoxy-carbonyl each having up to 4 carbon atoms, or linear or branched alkyl having up to 4 carbon atoms which itself can be substituted by hydroxyl or by linear or branched alkoxy having up to 3 carbon atoms,

R^1 and R^2 , including the double bond linking them, together form a phenyl ring or pyridyl ring or a ring of the formula



in which

R^5 denotes hydrogen or linear or branched alkyl having up to 3 carbon atoms,

R^3 and R^4 , including the double bond linking them, together form a phenyl ring or a cyclopentene,

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.