



US007932268B2

(12) **United States Patent Rader**

(10) **Patent No.:** US 7,932,268 B2
(45) **Date of Patent:** Apr. 26, 2011

(54) **METHODS FOR TREATING DISORDERS OR DISEASES ASSOCIATED WITH HYPERLIPIDEMIA AND HYPERCHOLESTEROLEMIA WHILE MINIMIZING SIDE EFFECTS**

(75) Inventor: **Daniel J. Rader**, Philadelphia, PA (US)
(73) Assignee: **The Trustees of the University of Pennsylvania**, Philadelphia, PA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 895 days.

(21) Appl. No.: **10/591,923**

(22) PCT Filed: **Mar. 7, 2005**

(86) PCT No.: **PCT/US2005/007435**

§ 371 (c)(1),
(2), (4) Date: **Jun. 21, 2007**

(87) PCT Pub. No.: **WO2005/087234**

PCT Pub. Date: **Sep. 22, 2005**

(65) **Prior Publication Data**

US 2009/0042941 A1 Feb. 12, 2009

Related U.S. Application Data

(60) Provisional application No. 60/550,915, filed on Mar. 5, 2004.

(51) **Int. Cl.**
A61K 31/445 (2006.01)

(52) **U.S. Cl.** **514/321; 514/325**

(58) **Field of Classification Search** 514/252.03, 514/255.03, 263.22, 321, 325, 824, 210.02
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,983,140 A	9/1976	Endo et al.
4,231,938 A	11/1980	Monaghan et al.
4,346,227 A	8/1982	Terahara et al.
4,448,784 A	5/1984	Glamkowski et al.
4,450,171 A	5/1984	Hoffman et al.
4,499,289 A	2/1985	Baran et al.
4,613,610 A	9/1986	Wareing
4,647,576 A	3/1987	Hoefle et al.
4,686,237 A	8/1987	Anderson
4,716,175 A	12/1987	Hoefle et al.
4,871,721 A	10/1989	Biller
4,924,024 A	5/1990	Biller
5,015,644 A	5/1991	Roth et al.
5,026,554 A	6/1991	Bartizal et al.
5,117,080 A	5/1992	Lee
5,510,379 A	4/1996	Lee et al.
5,595,872 A	1/1997	Wetterau, II et al.
5,684,014 A	11/1997	Muller et al.
5,712,279 A	1/1998	Biller et al.
5,712,396 A	1/1998	Magnin et al.
5,739,135 A *	4/1998	Biller et al. 514/252.03

5,767,115 A *	6/1998	Rosenblum et al. 514/210.02
5,786,361 A	7/1998	Muller et al.
5,789,197 A	8/1998	Wetterau, II et al.
5,811,429 A	9/1998	Connell et al.
5,827,875 A	10/1998	Dickson, Jr. et al.
5,883,109 A *	3/1999	Gregg et al. 514/321
5,885,983 A	3/1999	Biller et al.
5,952,498 A	9/1999	Lenfers et al.
5,990,110 A	11/1999	Firestone
6,034,115 A	3/2000	Connell et al.
6,057,339 A	5/2000	Gregg
6,066,650 A	5/2000	Biller et al.
6,066,653 A	5/2000	Gregg et al.
6,114,341 A	9/2000	Muller et al.
6,121,283 A	9/2000	Chang et al.
6,140,343 A	10/2000	DeNinno et al.
6,194,454 B1 *	2/2001	Dow 514/522
6,245,775 B1	6/2001	Muller et al.
6,265,431 B1	7/2001	Muller et al.
6,297,233 B1	10/2001	Stein et al.
6,344,450 B1	2/2002	Bisacchi et al.
6,479,503 B2	11/2002	Muller et al.
6,492,365 B1	12/2002	Wetterau, II et al.
6,498,156 B2	12/2002	Glombik et al.
6,620,821 B2	9/2003	Robl
6,627,636 B2	9/2003	Robl
6,720,351 B2	4/2004	Bertinato et al.
6,774,236 B1	8/2004	Lenfers et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU 727895 7/1998

(Continued)

OTHER PUBLICATIONS

Catapano, Ezetimibe: a selective inhibitor of cholesterol absorption, *European Heart Journal Supplements* 2001 (3, Supplemental E): E6-E10).

Knopp RH, Drug treatment of lipid disorders. *New England J. Med.* 1999; 341(7): 498-511; electronic pp. 1-25.

Aggarwal, et al; *BMC Cardiovasc. Disord.* 27;5(1):30 (2005).

Chandler, et al., *J. Lipid. Res.* 44(10):1887-901 (2003).

Cuchel et al., "Inhibition of Microsomal Triglyceride Transfer Protein in Familial Hypercholesterolemia," *N Engl J Med.*, (2007); 356:148-156.

Funatsu et al. "Atorvastatin Increases Hepatic Fatty Acid Beta-Oxidation in Sucrose-Fed Rats: Comparison with an MTP Inhibitor." *Eur. J. Pharm.* 2002 455:161-167.

(Continued)

Primary Examiner — Kevin Weddington
(74) *Attorney, Agent, or Firm* — Goodwin Procter LLP

(57) **ABSTRACT**

The present invention provides methods and compositions for treating hyperlipidemia and/or hypercholesterolemia comprising administering to the subject an effective amount of an MTP inhibitor to inhibit hyperlipidemia and/or hypercholesterolemia in said subject, wherein said administration comprises an escalating series of doses of the MTP inhibitor. In some embodiments the method comprises administering at least three step-wise, increasing dosages of the MTP inhibitor to the subject. In some embodiments, the method further comprises the administration of one or more other lipid modifying compounds.

U.S. PATENT DOCUMENTS

6,812,345 B2 11/2004 Robl et al.
 6,846,836 B2 1/2005 Hamann et al.
 6,858,622 B2 2/2005 Muller et al.
 6,875,782 B2 4/2005 Cheng et al.
 6,884,812 B2 4/2005 Glombick et al.
 6,916,809 B2 7/2005 Chen et al.
 6,916,813 B2 7/2005 Atwal et al.
 6,949,572 B2 9/2005 Bertinato et al.
 6,979,692 B2 12/2005 Bertinato et al.
 7,053,080 B2 5/2006 Davis et al.
 7,056,906 B2 6/2006 Strony
 7,358,254 B2 4/2008 Robl et al.
 7,394,501 B2 7/2008 Iwata et al.
 7,645,732 B2 1/2010 Ye et al.
 2002/0035064 A1 3/2002 Robl et al.
 2002/0045271 A1 4/2002 Hussain et al.
 2003/0069221 A1 4/2003 Kosoglou et al.
 2003/0109543 A1 6/2003 Ogletree
 2003/0153541 A1 8/2003 Dudley et al.
 2003/0162788 A1 8/2003 Thomas et al.
 2003/0187053 A1 10/2003 Bertinato et al.
 2004/0014748 A1 1/2004 Grutzmann et al.
 2004/0058908 A1 3/2004 Keller et al.
 2005/0075367 A1 4/2005 Hagiwara et al.
 2005/0090426 A1 4/2005 Blumberg
 2005/0101561 A1 5/2005 Tunac
 2006/0069161 A1 3/2006 Lee et al.
 2006/0135460 A1 6/2006 Widder et al.
 2006/0153913 A1 7/2006 Yamane et al.
 2006/0160834 A1 7/2006 Fong et al.
 2006/0166999 A1 7/2006 Grutzmann et al.
 2006/0205726 A1 9/2006 Hagiwara et al.
 2006/0211020 A1 9/2006 Farrer et al.
 2006/0211762 A1 9/2006 Rongen et al.
 2006/0252733 A1 11/2006 Jansen
 2006/0270655 A1 11/2006 Swick et al.
 2007/0027093 A1 2/2007 Ogawa et al.
 2007/0032404 A1 2/2007 Sweet et al.
 2007/0088089 A1 4/2007 Wisler
 2007/0093468 A1 4/2007 Wisler
 2007/0093527 A1 4/2007 Wisler
 2007/0098778 A1 5/2007 Borsadia
 2007/0099884 A1 5/2007 Erondu et al.
 2008/0016127 A1 1/2008 Field
 2008/0033019 A1 2/2008 Stamler
 2008/0051427 A1 2/2008 Schuckler
 2008/0103122 A1 5/2008 Veltri
 2008/0161279 A1 7/2008 Wisler
 2008/0175864 A1 7/2008 Ye et al.
 2008/0241869 A1 10/2008 Davis
 2008/0248070 A1 10/2008 Tunac
 2008/0253985 A1 10/2008 Wisler
 2008/0255084 A1 10/2008 Webb
 2008/0280992 A1 11/2008 Kunz et al.
 2009/0042835 A1 2/2009 Davis
 2009/0042941 A1 2/2009 Rader
 2009/0054393 A1 2/2009 Wisler
 2009/0093527 A1 4/2009 Li et al.

FOREIGN PATENT DOCUMENTS

CA 2091102 9/1993
 CA 2291471 6/2000
 CA 2325201 5/2001
 DE 19951022 4/2001
 EP 0142146 5/1985
 EP 0221025 5/1987
 EP 0325130 7/1989
 EP 0705831 4/1996
 EP 0779276 6/1997
 EP 0779279 6/1997
 EP 0799828 10/1997
 EP 0802198 10/1997
 EP 1099442 5/2001
 EP 1 181 954 A 2/2002
 FR 2596393 10/1987

WO WO-86/03488 6/1986
 WO WO-86/07054 12/1986
 WO WO-96/26205 8/1996
 WO WO-96/26948 A1 9/1996
 WO WO-96/40640 12/1996
 WO WO-97/41111 11/1997
 WO WO-98/03069 1/1998
 WO WO-98/03174 1/1998
 WO WO-98/23593 6/1998
 WO WO-98/27979 7/1998
 WO WO-98/31225 7/1998
 WO WO-98/31366 7/1998
 WO WO-98/31367 7/1998
 WO WO-98/50028 11/1998
 WO WO-00/38725 7/2000
 WO WO-01/08679 2/2001
 WO WO-2004/028544 4/2004
 WO WO-2004/110368 12/2004
 WO WO-2004/110375 12/2004
 WO WO-2005/000217 1/2005
 WO WO-2005/033100 A1 4/2005
 WO WO-2005/051382 6/2005
 WO WO-2005/072740 8/2005
 WO WO-2005/084666 9/2005
 WO WO-2005/087234 9/2005
 WO WO-2005/094864 10/2005
 WO WO-2005/097131 10/2005
 WO WO-2006/046623 5/2006
 WO WO-2006/062748 6/2006
 WO WO-2006/063128 6/2006
 WO WO-2006/108666 10/2006
 WO WO-2006/111238 10/2006
 WO WO-2007/047724 4/2007
 WO WO-2007/047880 4/2007
 WO WO-2008/012056 1/2008
 WO WO-2008/021353 2/2008
 WO WO-2008/030382 3/2008
 WO WO-2008/072061 6/2008
 WO WO-2008/075949 6/2008
 WO WO-2008/079398 A1 7/2008
 WO WO-2008/090198 7/2008
 WO WO-2008/115574 9/2008

OTHER PUBLICATIONS

Li, et al., "Discovery of Potent and Orally Active MTP Inhibitors as Potential Anti-Obesity Agents," *Bioorganic & Medicinal Chemistry Letters*, Oxford, GB, vol. 16, No. 11, Jun. 1, 2006, pp. 3039-3042.
 Looije, Norbert A., et al., "Disodium Ascorbyl Phytostanyl Phosphates (FM-VP4) Reduces Plasma Cholesterol Concentration, Body Weight and Abdominal Fat Gain Within a Dietary-Induced Obese Mouse Model," *Journal of Pharmacy & Pharmaceutical Sciences: A Publication of the Canadian Society for Pharmaceutical Sciences*, vol. 8, No. 3, 2005, pp. 400-408.
 Samaha, et al., "Inhibition Of Microsomal Triglyceride Transfer Protein Alone Or With Ezetimibe In Patients With Moderate Hypercholesterolemia," *Nature Clinical Practice*, (2008), pp. 1-9, Nature Publishing Group.
 Atzel, A., et al., "Mechanism of Microsomal Triglyceride Transfer Protein Catalyzed Lipid Transport", *Biochemistry* (1993), 32, 10444-10450.
 Bakillah A. et al., "Decreased Secretion of ApoB follows inhibition of ApoB-MTP Binding by a Novel Antagonist", *Biochemistry (Mosc)*, (2000) 39(16):4892-4899.
 Barclay, "Hyperlipidemia", *NMT Briefs*, 2003.
 Bayes, M., et al., "Gateways to Clinical Trials", *Methods and Findings in Experimental and Clinical Pharmacology*, 24(1):2002, 37-55.
 Bays et al., "Pharmacotherapy for Dyslipidaemia—Current Therapies and Future Agents", *Expert Opin Pharmacother*, Nov. 2003;4(11):1901-38.
 Biller et al., "Isoprenoid (Phosphinylmethyl) Phosphonates as Inhibitors of Squalene synthetase", *J. Med. Chem.*, 31 (10):1988, 1869-71.
 Bruckert, "New Lipid-Modifying Therapies", *Expert Opin. Investig. Drugs*, (2003)12(3):325-35.
 G. ...

- Chang et al., "Microsomal Triglyceride Transfer Protein (MTP) Inhibitors: Discovery of Clinically Active Inhibitors Using High-Throughput Screening and Parallel Synthesis Paradigms", *Current Opinion in Drug Discovery & Development*, (2002) 5(4):562-70.
- Corey and Volante, *J. Am. Chem. Soc.*, (1976) 98:1291-93.
- de Montellano et al., "Inhibition of Squalene Synthetase by Farnesyl Pyrophosphate Analogues" *J. Med. Chem.*, (1977) 20:243-49.
- Earl et al., "Ezetimibe", *Nature Reviews*, 2003, 2:97-98.
- Evans M. et al. "Medical Lipid-Regulating Therapy: Current Evidence, Ongoing Trials and Future Developments", *Drugs*: 2004; 64 (11): pp. 1181-1196.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. National Cholesterol Education Program; Adult Treatment Panel III Report. Publication No. 01-3095, I-I-IX-11. 2001. Bethesda, MD, National Heart, Lung, and Blood Institute.
- Farrell., "Drugs and Steatohepatitis", *Semin Liver Dis*, (2002) 22(2):185-194.
- Gagne, et al. "Efficacy and Safety of Ezetimibe Coadministered with Atorvastatin or Simvastatin in Patients With Homozygous Familial Hypercholesterolemia", *Circulation*, (2002) 105 (21):2469-75.
- International Search Report for Application No. PCT/US05/07435 dated Jul. 14, 2005 (7 pages).
- International Search Report for Application No. PCT/US06/040637 dated Jun. 12, 2007 (8 pages).
- International Search Report for Application No. PCT/US06/040639 dated Jun. 12, 2007 (9 pages).
- International Search Report for Application No. PCT/US06/040640 dated May 23, 2007 (9 pages).
- International Search Report for Application No. PCT/US06/040953 dated Mar. 3, 2007 (8 pages).
- Jamil et al., "An inhibitor of the Microsomal Triglyceride Transfer Protein Inhibits apoB Secretion From HepG2 Cells", *Proc Natl Acad Sci U S A*, (1996) 93(21):11991-11995.
- Kastelein J., "What future for Combination Therapies", *Int J. Clin Pract. Suppl. Mar.* 2003; (134): pp. 45-50.
- Kirkpatrick et al., "Market indicators", *Nature*, 2003, 2:98.
- Liao et al., "Blocking Microsomal Triglyceride Transfer Protein Interferes with apoB Secretion Without Causing Retention or Stress in the ER", *Journal of Lipid Research*, (2003) 44(5):978.
- McClard et al., *J.A.C.S.*, (1987) 109:5544.
- Ritter et al., "Heterocyclic Ring Scaffolds as Small-Molecule Cholesterol Absorption Inhibitors", *Org. Biomol. Chem.*, 2005, 3:3514-3523.
- Robl et al., "A Novel Series of Highly Potent Benzimidazole-Based Microsomal Triglyceride Transfer Protein Inhibitors", *Journal of Medicinal Chemistry*, 2001, 44(6):851-856.
- Shiomi et al., "MTP Inhibitor Decreases Plasma Cholesterol Levels in LDL Receptor-Deficient WHHL Rabbits by Lowering the VLDL Secretion", *Euro. Journal of Pharma.* 2001, 431:127-131.
- Sorbera, L.A., et al., Implitapide "Hypolipidemic Treatment of Atherosclerosis MTP Inhibitor ApoB Secretion Inhibitor", *Drugs of the Future*, (2000) 25(11):1138-1144.
- Subhop et al., "Cholesterol Absorption Inhibitors For the Treatment of Hypercholesterolaemia", *Drugs*, 2002, 62(16):2333-2347.
- Thomas et al., "Alleviation of MTP Inhibitor-Induced Hepatic Steatosis in Hyperlipidemic *fa/fa* Rats by Fenofibrate", Dept. of Metabolic Diseases and Dept. of Chemical Research, Boehringer Ingelheim Pharma GmbH & Co. KG., (2005).
- Wetterau et al., "An MTP Inhibitor that Normalizes Atherogenic Lipoprotein Levels in WHHL Rabbits", *Science*, (1998) 282(5389):751-754.
- Wetterau et al., "Microsomal Triglyceride Transfer Protein", *Biochimica Biophysica Acta*, (1997) 1345(2):136-150.
- Wierzbicki A.S., "New Lipid-Lowering Agents", *Expert Opinion on Emerging Drugs*, Ashley Publications, GB, 8 (2):2003, 365-376.

* cited by examiner

1

**METHODS FOR TREATING DISORDERS OR
DISEASES ASSOCIATED WITH
HYPERLIPIDEMIA AND
HYPERCHOLESTEROLEMIA WHILE
MINIMIZING SIDE EFFECTS**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

This application is a national phase application under 35 U.S.C. §371 of PCT/US05/007435 filed Mar. 7, 2005 which in turn claims priority benefit of U.S. Ser. No. 60/550,915, filed Mar. 5, 2004, all of which are hereby incorporated by reference in their entireties.

FIELD OF THE INVENTION

The present invention generally relates to therapy for hypercholesterolemia and hyperlipidemia.

BACKGROUND OF THE INVENTION

Hypercholesterolemia is a well-known risk factor for ASCVD, the major cause of mortality in the Western world. Numerous epidemiological studies have clearly demonstrated that pharmacological lowering of total cholesterol (TC) and Low-density Lipoprotein (LDL) Cholesterol (LDL-C) is associated with a significant reduction in clinical cardiovascular events. Hypercholesterolemia is often caused by a polygenic disorder in the majority of cases and modifications in lifestyle and conventional drug treatment are usually successful in reducing cholesterol levels. However, in few cases, as in familial hypercholesterolemia (FH), the cause is a monogenic defect and the available treatment in homozygous patients can be much more challenging and far from optimal because LDL-C levels remain extremely elevated despite aggressive use of combination therapy. Therefore, for this group of high-risk patients, effective medical therapy is urgently needed.

Triglycerides are common types of fats (lipids) that are essential for good health when present in normal amounts. They account for about 95 percent of the body's fatty tissue. Abnormally high triglyceride levels may be an indication of such conditions as cirrhosis of the liver, underactive thyroid (hypothyroidism), poorly controlled diabetes, or pancreatitis (inflammation of the pancreas). Researchers have identified triglycerides as an independent risk factor for heart disease.

Higher-than-normal triglyceride levels are often associated with known risk factors for heart disease, such as low levels of HDL ("good") cholesterol, high levels of LDL ("bad") cholesterol and obesity. Triglycerides may also contribute to thickening of artery walls—a physical change believed to be a predictor of atherosclerosis.

Therefore, high triglyceride levels are at least a warning sign that a patient's heart health may be at risk. In response, physicians may be more likely to stress the importance of losing weight, getting enough exercise, quitting smoking, controlling diabetes and other strategies that patients can use to protect their own cardiovascular health.

A large number of genetic and acquired diseases can result in hyperlipidemia. They can be classified into primary and secondary hyperlipidemic states. The most common causes of the secondary hyperlipidemias are diabetes mellitus, alcohol abuse, drugs, hypothyroidism, chronic renal failure, nephrotic syndrome, cholestasis and bulimia. Primary hyperlipidemias have also been classified into common hypercho-

2

hypercholesterolemia, remnant hyperlipidemia, chylomicronemia syndrome and familial hypertriglyceridemia.

A number of treatments are currently available for lowering serum cholesterol and triglycerides. However, each has its own drawbacks and limitations in terms of efficacy, side-effects and qualifying patient population.

Bile-acid-binding resins are a class of drugs that interrupt the recycling of bile acids from the intestine to the liver; e.g., cholestyramine (Questran Light®, Bristol-Myers Squibb), and colestipol hydrochloride (Colestid®, The Upjohn Company). When taken orally, these positively-charged resins bind to the negatively charged bile acids in the intestine. Because the resins cannot be absorbed from the intestine, they are excreted carrying the bile acids with them. The use of such resins, however, at best only lowers serum cholesterol levels by about 20%, and is associated with gastrointestinal side-effects, including constipation and certain vitamin deficiencies. Moreover, since the resins bind other drugs, other oral medications must be taken at least one hour before or four to six hours subsequent to ingestion of the resin; thus, complicating heart patient's drug regimens.

The statins are cholesterol-lowering agents that block cholesterol synthesis by inhibiting HMGCoA reductase—the key enzyme involved in the cholesterol biosynthetic pathway. The statins, e.g., lovastatin (Mevacor®, Merck & Co., Inc.), simvastatin (Zocor®, Merck & Co., Inc.), atorvastatin (Lipitor®, Pfizer), rosuva (Crestor®, Astra Zeneca) and pravastatin (Pravachol®, Bristol-Myers Squibb Co.), and combinations thereof, are sometimes used in combination with bile-acid-binding resins. Statins significantly reduce serum cholesterol and LDL-serum levels, and slow progression of coronary atherosclerosis. However, serum HDL cholesterol levels are only moderately increased. The mechanism of the LDL lowering effect may involve both reduction of VLDL concentration and induction of cellular expression of LDL-receptor, leading to reduced production and/or increased catabolism of LDLs. Side effects, including liver and kidney dysfunction are associated with the use of these drugs (Physicians Desk Reference, Medical Economics Co., Inc., Montvale, N.J., 2004; hereinafter "PDR"). The FDA has approved atorvastatin to treat rare but urgent cases of familial hypercholesterolemia.

Ezetimibe is a cholesterol absorption inhibitor which reduces the amount of cholesterol absorbed by the body. Ezetimibe is used to reduce the amount of total cholesterol, LDL cholesterol (by about 18%), and apolipoprotein B. Ezetimibe is often used with a low cholesterol diet and, in some cases, other cholesterol lowering medications.

Niacin, or nicotinic acid, is a water soluble vitamin B-complex used as a dietary supplement and antihyperlipidemic agent. Niacin diminishes production of VLDL and is effective at lowering LDL. In some cases, it is used in combination with bile-acid binding resins. NIAPAN® has been approved to prevent recurrent heart attacks in patients with high cholesterol. Niacin can increase HDL when used at adequate doses, however, its usefulness is limited by serious side effects when used at such high doses.

Fibric acid derivatives ("fibrates") are a class of lipid-lowering drugs used to treat various forms of hyperlipidemia (i.e., elevated serum triglycerides) which may also be associated with hypercholesterolemia. Fibrates appear to reduce the VLDL fraction and modestly increase HDL. However, the effects of these drugs on serum cholesterol is variable. Fibrates are mainly used to lower high triglyceride levels. Although fibrates typically do not appear as effective as statins in lowering total cholesterol and LDL cholesterol lev-

3

other medications to lower very high cholesterol levels. For example, fibrates are also sometimes added to statins to raise HDL cholesterol levels. In the United States, fibrates have been approved for use as antilipidemic drugs, but have not received approval as hypercholesterolemia agents. For example, clofibrate (Atromid-S®, Wyeth-Ayerst Laboratories) is an antilipidemic agent which acts to lower serum triglycerides by reducing the VLDL fraction. Although serum cholesterol may be reduced in certain patient subpopulations, the biochemical response to the drug is variable, and is not always possible to predict which patients will obtain favorable results. Atromid-S® has not been shown to be effective for prevention of coronary heart disease. The chemically and pharmacologically related drug, gemfibrozil (Lopid®, Parke-Davis) is a lipid regulating agent which moderately decreases serum triglycerides and VLDL cholesterol, and moderately increases HDL cholesterol—the HDL₂ and HDL₃ subfractions as well as both ApoA-I and A-II (i.e., the AI/AII-HDL fraction). However, the lipid response is heterogeneous, especially among different patient populations. Moreover, while prevention of coronary heart disease was observed in male patients between 40-55 without history or symptoms of existing coronary heart disease, it is not clear to what extent these findings can be extrapolated to other patient populations (e.g., women, older and younger males). Indeed, no efficacy was observed in patients with established coronary heart disease. Fenofibrate (Tricor, Secalip) is also used to reduce levels of cholesterol and triglycerides. Serious side-effects have been associated with the use of several fibrates including toxicity such as malignancy, (especially gastrointestinal cancer), gallbladder disease and an increased incidence in non-coronary mortality. Fibrates are often not indicated for the treatment of patients with high LDL or low HDL as their only lipid abnormality (Physician's Desk Reference, 2004, Medical Economics Co., Inc. Montvale, N.J.).

Oral estrogen replacement therapy may be considered for moderate hypercholesterolemia in post-menopausal women. However, increases in HDL may be accompanied with an increase in triglycerides. Estrogen treatment is, of course, limited to a specific patient population (postmenopausal women) and is associated with serious side effects including induction of malignant neoplasms, gall bladder disease, thromboembolic disease, hepatic adenoma, elevated blood pressure, glucose intolerance, and hypercalcemia.

Homozygous familial hypercholesterolemia (hoFH) is a serious life-threatening genetic disease caused by homozygosity or compound heterozygosity for mutations in the low density lipoprotein (LDL) receptor. Total plasma cholesterol levels are generally over 500 mg/dl and markedly premature atherosclerotic vascular disease is the major consequence. Untreated, most patients develop atherosclerosis before age 20 and generally do not survive past age 30. The primary goal of therapy consists of controlling the hypercholesterolemia to delay the development of atherosclerotic cardiovascular disease (ASCVD). However, patients diagnosed with hoFH are largely unresponsive to conventional drug therapy and have limited treatment options. A mean LDL-C reduction of only about 5.5% has been recently reported in patients with genotype-confirmed hoFH treated with the maximal dose of statins (atorvastatin or simvastatin 80 mg/day). The addition of ezetimibe 10 mg/day to this regimen resulted in a total reduction of LDL-C levels of 27%, which is still far from optimal. Several non-pharmacological options have also been tested. Surgical interventions, such as portacaval shunt and ileal bypass have resulted only in partial and transient LDL-C lowering. Orthotopic liver transplantation has been demon-

4

patients, but obvious disadvantages and risks are associated with this approach. Although hoFH could be an excellent model for gene therapy, this modality of treatment is not foreseeable in the near future due to the limitations on the availability of safe vectors that provide long-term expression of LDL receptor gene. Thus, the current standard of care in hoFH is LDL apheresis, a physical method of filtering the plasma of LDL-C which as monotherapy can transiently reduce LDL-C by about 50%. Apheresis uses affinity columns to selectively remove apoB-containing lipoproteins. However, because of rapid re-accumulation of LDL-C in plasma, apheresis has to be repeated frequently (every 1-2 weeks) and requires 2 separate sites for IV access. Although anecdotally this procedure may delay the onset of atherosclerosis, it is laborious, expensive, and not readily available. Furthermore, although it is a procedure that is generally well tolerated, the fact that it needs frequent repetition and IV access can be challenging for many of these young patients. Therefore, there is a tremendous unmet medical need for new medical therapies for hoFH.

Patients with heterozygous FH can usually be successfully treated with combination drug therapy to lower the LDL-C to acceptable levels. In contrast, hoFH is unresponsive to conventional drug therapy and thus there are limited treatment options. Specifically, treatment with statins, which reduce LDL-C by inhibiting cholesterol synthesis and upregulating the hepatic LDL receptor, have negligible effect in patients whose LDL receptors are non-existent or defective.

In July 2004, the NCEP published a paper entitled "Implications of Recent Clinical Trials for the National Cholesterol Education Program Adult Treatment Panel III Guidelines", updating certain elements of the "Adult Treatment Panel III (ATP III)" cholesterol guidelines released in 2001. For high-risk patients, individuals who have coronary heart disease (CHD) or disease of the blood vessels to the brain or extremities, or diabetes, or multiple (2 or more) risk factors that give them a greater than 20 percent chance of having a heart attack within 10 years, the ATP III update recommends that the overall goal for high-risk patients is still an LDL less than 100 mg/dL with a therapeutic option to set the goal at an LDL less than 70 mg/dL for very high-risk patients, those who have had a recent heart attack, or those who have cardiovascular disease combined with either diabetes, or severe or poorly controlled risk factors (such as continued smoking), or metabolic syndrome (a cluster of risk factors associated with obesity that includes high triglycerides and low HDL cholesterol). The ATP III update also recommends consideration of drug treatment in addition to lifestyle therapy for LDL levels 100 mg/dL or higher in high-risk patients, and characterizes drug treatment as optional for LDL less than 100 mg/dL. For moderately high-risk patients, individuals who have multiple (2 or more) CHD risk factors together with a 10-20 percent risk for a heart attack within 10 years, the ATP III update recommends the overall goal for moderately high-risk patients to be an LDL less than 130 mg/dL. There is a therapeutic option to set the treatment goal at an LDL less than 100 mg/dL, and to use drug treatment if LDL is 100-129 mg/dL. For high-risk and moderately high-risk patients, the ATP III update advises that the intensity of LDL-lowering drug treatment in high-risk and moderately high-risk patients be sufficient to achieve at least a 30 percent reduction in LDL levels.

Patients suffering from severe hypercholesterolemia may also be unable to reach the new goals for LDL and HDL described above. For example, a large number of patients may be unable to attain LDL levels less than 70 using maximally

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.