

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use XYREM safely and effectively. See full prescribing information for XYREM.

XYREM® (sodium oxybate) oral solution, CIII
Initial U.S. Approval: 2002

WARNING: CENTRAL NERVOUS SYSTEM (CNS) DEPRESSION and MISUSE AND ABUSE.

See full prescribing information for complete boxed warning.

- Respiratory depression can occur with Xyrem use (5.4)
- Xyrem is a Schedule III controlled substance and is the sodium salt of gamma hydroxybutyrate (GHB), a Schedule I controlled substance. Abuse or misuse of illicit GHB is associated with CNS adverse reactions, including seizure, respiratory depression, decreased consciousness, coma and death (5.2, 9.2)
- Because of the risks of CNS depression, abuse, and misuse, Xyrem is available only through a restricted distribution program called the Xyrem Success Program® using a centralized pharmacy. Prescribers and patients must enroll in the program. (5.3)

RECENT MAJOR CHANGES

Dosage and Administration, Dose Adjustment with Co-administration of Divalproex Sodium (2.4)
04/2014

INDICATIONS AND USAGE

Xyrem is a central nervous system depressant indicated for the treatment of:

- Cataplexy in narcolepsy (1.1)
- Excessive daytime sleepiness (EDS) in narcolepsy (1.2)

Xyrem may only be dispensed to patients enrolled in the Xyrem Success Program (1).

DOSAGE AND ADMINISTRATION

- Initiate dose at 4.5 grams (g) per night administered orally in two equal, divided doses: 2.25 g at bedtime and 2.25 g taken 2.5 to 4 hours later (2.1)
- Titrate to effect in increments of 1.5 g per night at weekly intervals (0.75 g at bedtime and 0.75 g taken 2.5 to 4 hours later) (2.1).
- Recommended dose range: 6 g to 9 g per night orally (2.1).

Total Nightly Dose	Take at Bedtime	Take 2.5 to 4 Hours Later
4.5 g per night	2.25 g	2.25 g
6 g per night	3 g	3 g
7.5 g per night	3.75 g	3.75 g
9 g per night	4.5 g	4.5 g

- Take each dose while in bed and lie down after dosing (2.2).

- Allow 2 hours after eating before dosing (2.2).
- Prepare both doses prior to bedtime; dilute each dose with approximately ¼ cup of water in pharmacy-provided vials (2.2).
- Patients with Hepatic Impairment: starting dose is 2.25 g per night administered orally in two equal, divided doses of approximately 1.13 g at bedtime and approximately 1.13 g taken 2.5 to 4 hours later (2.3).
- Concomitant use with divalproex sodium: an initial reduction in Xyrem dose of at least 20% is recommended (2.4, 7.1).

DOSAGE FORMS AND STRENGTHS

Oral solution, 0.5 g per mL (3)

CONTRAINDICATIONS

- Succinic semialdehyde dehydrogenase deficiency (4)
- In combination with sedative hypnotics or alcohol (4)

WARNINGS AND PRECAUTIONS

- CNS depression: Use caution when considering the concurrent use of Xyrem with other CNS depressants (5.1).
- Caution patients against hazardous activities requiring complete mental alertness or motor coordination within the first 6 hours of dosing or after first initiating treatment until certain that Xyrem does not affect them adversely (5.1).
- Depression and suicidality: Monitor patients for emergent or increased depression and suicidality (5.5).
- Confusion/Anxiety: Monitor for impaired motor/cognitive function (5.6).
- Parasomnias: evaluate episodes of sleepwalking (5.7).
- High sodium content in Xyrem: Monitor patients with heart failure, hypertension, or impaired renal function (5.8).

ADVERSE REACTIONS

Most common adverse reactions (≥ 5% and at least twice the incidence with placebo) were nausea, dizziness, vomiting, somnolence, enuresis, and tremor (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Jazz Pharmaceuticals at 1-800-520-5568, or FDA at 1-800-FDA-1088 or www.fda.gov/Medwatch.

USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm (8.1).
- Geriatric patients: Monitor for impaired motor and/or cognitive function when taking Xyrem (8.5).

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 4/2014

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: CENTRAL NERVOUS SYSTEM (CNS) DEPRESSION and MISUSE AND ABUSE

1 INDICATIONS AND USAGE

- 1.1 Cataplexy in Narcolepsy
- 1.2 Excessive Daytime Sleepiness in Narcolepsy

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing Information
- 2.2 Important Administration Instructions
- 2.3 Dose Modification in Patients with Hepatic Impairment
- 2.4 Dose Adjustment with Co-administration of Divalproex Sodium

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Central Nervous System Depression
- 5.2 Abuse and Misuse
- 5.3 Xyrem Success Program
- 5.4 Respiratory Depression and Sleep-Disordered Breathing
- 5.5 Depression and Suicidality
- 5.6 Other Behavioral or Psychiatric Adverse Reactions
- 5.7 Parasomnias

- 5.8 Use in Patients Sensitive to High Sodium Intake

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Alcohol, Sedative Hypnotics, and CNS Depressants
- 7.2 Divalproex Sodium

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

- 10.1 Human Experience
- 10.2 Signs and Symptoms
- 10.3 Recommended Treatment of Overdose

Reference ID: A4285521

- 10.4 Poison Control Center
- 11 DESCRIPTION**
- 12 CLINICAL PHARMACOLOGY**
 - 12.1 Mechanism of Action
 - 12.3 Pharmacokinetics
- 13 NONCLINICAL TOXICOLOGY**
 - 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 14 CLINICAL STUDIES**
 - 14.1 Cataplexy in Narcolepsy

- 14.2 Excessive Daytime Sleepiness in Narcolepsy
- 16 HOW SUPPLIED/STORAGE AND HANDLING**
 - 16.1 How Supplied
 - 16.2 Storage
 - 16.3 Handling and Disposal
- 17 PATIENT COUNSELING INFORMATION**

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

WARNING: CENTRAL NERVOUS SYSTEM DEPRESSION and MISUSE AND ABUSE.

Xyrem (sodium oxybate) is a CNS depressant. In clinical trials at recommended doses obtundation and clinically significant respiratory depression occurred in Xyrem-treated patients. Almost all of the patients who received Xyrem during clinical trials in narcolepsy were receiving central nervous system stimulants [see *Warnings and Precautions (5.1)*].

Xyrem[®] (sodium oxybate) is the sodium salt of gamma hydroxybutyrate (GHB). Abuse of GHB, either alone or in combination with other CNS depressants, is associated with CNS adverse reactions, including seizure, respiratory depression, decreases in the level of consciousness, coma, and death [see *Warnings and Precautions (5.2)*].

Because of the risks of CNS depression, abuse, and misuse, Xyrem is available only through a restricted distribution program called the Xyrem Success Program[®], using a centralized pharmacy. Prescribers and patients must enroll in the program. For further information go to www.XYREM.com or call 1-866-XYREM88[®] (1-866-997-3688). [see *Warnings and Precautions (5.3)*].

1 INDICATIONS AND USAGE

Limitations of Use

Xyrem may only be dispensed to patients enrolled in the Xyrem Success Program [see *Warnings and Precautions (5.3)*].

1.1 Cataplexy in Narcolepsy

Xyrem (sodium oxybate) oral solution is indicated for the treatment of cataplexy in narcolepsy.

1.2 Excessive Daytime Sleepiness in Narcolepsy

Xyrem (sodium oxybate) oral solution is indicated for the treatment of excessive daytime sleepiness (EDS) in narcolepsy.

2 DOSAGE AND ADMINISTRATION

Healthcare professionals who prescribe Xyrem must enroll in the Xyrem Success Program and must comply with the requirements to ensure safe use of Xyrem [see *Warnings and Precautions (5.3)*].

2.1 Dosing Information

The recommended starting dose is 4.5 grams (g) per night administered orally in two equal, divided doses: 2.25 g at bedtime and 2.25 g taken 2.5 to 4 hours later (see Table 1). Increase the dose by 1.5 g per night at weekly intervals (additional 0.75 g at bedtime and 0.75 g taken 2.5 to 4 hours later) to the effective dose range of 6 g to 9 g per night orally. Doses higher than 9 g per night have not been studied and should not ordinarily be administered.

Table 1: Xyrem Dose Regimen (g = grams)

If A Patient's Total Nightly Dose is:	Take at Bedtime:	Take 2.5 to 4 Hours Later:
4.5 g per night	2.25 g	2.25 g
6 g per night	3 g	3 g
7.5 g per night	3.75 g	3.75 g
9 g per night	4.5 g	4.5 g

2.2 Important Administration Instructions

Take the first dose of Xyrem at least 2 hours after eating because food significantly reduces the bioavailability of sodium oxybate.

Prepare both doses of Xyrem prior to bedtime. Prior to ingestion, each dose of Xyrem should be diluted with approximately $\frac{1}{4}$ cup (approximately 60 mL) of water in the empty pharmacy vials provided. Patients should take both doses of Xyrem while in bed and lie down immediately after dosing as Xyrem may cause them to fall asleep abruptly without first feeling drowsy. Patients will often fall asleep within 5 minutes of taking Xyrem, and will usually fall asleep within 15 minutes, though the time it takes any individual patient to fall asleep may vary from night to night. Patients should remain in bed following ingestion of the first and second doses, and should not take the second dose until 2.5 to 4 hours after the first dose. Patients may need to set an alarm to awaken for the second dose. Rarely, patients may take up to 2 hours to fall asleep.

2.3 Dose Modification in Patients with Hepatic Impairment

The recommended starting dose in patients with hepatic impairment is 2.25 g per night administered orally in two equal, divided doses: approximately 1.13 g at bedtime and approximately 1.13 g taken 2.5 to 4 hours later [see *Use in Specific Populations (8.6)*; *Clinical Pharmacology (12.3)*].

2.4 Dose Adjustment with Co-administration of Divalproex Sodium

Pharmacokinetic and pharmacodynamic interactions have been observed when Xyrem is co-administered with divalproex sodium. For patients already stabilized on Xyrem, it is recommended that addition of divalproex sodium should be accompanied by an initial reduction in the nightly dose of Xyrem by at least 20%. For patients already taking divalproex sodium, it is recommended that prescribers use a lower starting Xyrem dose when introducing Xyrem. Prescribers should monitor patient response and adjust dose accordingly. [see *Drug Interactions (7.1)* and *Clinical Pharmacology (12.3)*].

3 DOSAGE FORMS AND STRENGTHS

Xyrem is a clear to slightly opalescent oral solution, in a concentration of 0.5 g per mL.

4 CONTRAINDICATIONS

Xyrem is contraindicated in patients being treated with sedative hypnotic agents.

Patients should not drink alcohol when using Xyrem.

Xyrem is contraindicated in patients with succinic semialdehyde dehydrogenase deficiency. This is a rare disorder of inborn error of metabolism variably characterized by mental retardation, hypotonia, and ataxia.

5 WARNINGS AND PRECAUTIONS

5.1 Central Nervous System Depression

Xyrem is a central nervous system (CNS) depressant. Alcohol and sedative hypnotics are contraindicated in patients who are using Xyrem. The concurrent use of Xyrem with other CNS depressants, including but not limited to opioid analgesics, benzodiazepines, sedating antidepressants or antipsychotics, sedating anti-epileptic drugs, general anesthetics, muscle relaxants, and/or illicit CNS depressants, may increase the risk of respiratory depression, hypotension, profound sedation, syncope, and death. If use of these CNS depressants in combination with Xyrem is required, dose reduction or discontinuation of one or more CNS depressants (including Xyrem) should be considered. In addition, if short-term use of an opioid (e.g. post- or perioperative) is required, interruption of treatment with Xyrem should be considered.

Healthcare providers should caution patients about operating hazardous machinery, including automobiles or airplanes, until they are reasonably certain that Xyrem does not affect them adversely (e.g., impair judgment, thinking, or motor skills). Patients should not engage in hazardous occupations or activities requiring complete mental alertness or motor coordination, such as operating machinery or a motor vehicle or flying an airplane, for at least 6 hours after taking the second nightly dose of Xyrem. Patients should be queried about CNS depression-related events upon initiation of Xyrem therapy and periodically thereafter [see *Warnings and Precautions (5.3)*].

5.2 Abuse and Misuse

Xyrem is a Schedule III controlled substance. The active ingredient of Xyrem, sodium oxybate or gamma-hydroxybutyrate (GHB), is a Schedule I controlled substance. Abuse of illicit GHB, either alone or in combination with other CNS depressants, is associated with CNS adverse reactions, including seizure, respiratory depression, decreases in the level of consciousness, coma, and death. The rapid onset of sedation, coupled with the amnesic features of Xyrem, particularly when combined with alcohol, has proven to be dangerous for the voluntary and involuntary user (e.g., assault victim). Because illicit use and abuse of GHB have been reported, physicians should carefully evaluate patients for a history of drug abuse and follow such patients closely, observing them for signs of misuse or abuse of GHB (e.g. increase in size or frequency of dosing, drug-seeking behavior, feigned cataplexy) [see *Warnings and Precautions (5.3)* and *Drug Abuse and Dependence (9.2)*].

5.3 Xyrem Success Program

Because of the risks of central nervous system depression and abuse/misuse, Xyrem is available only through a restricted distribution program called the Xyrem Success Program.

Required components of the Xyrem Success Program are:

- Use of a centralized pharmacy
- Healthcare Providers who prescribe Xyrem must complete the enrollment forms and comply with the requirements.
- To receive Xyrem, patients must understand the risks and benefits of Xyrem.

Further information is available at www.XYREM.com or 1-866-XYREM88[®] (1-866-997-3688).

5.4 Respiratory Depression and Sleep-Disordered Breathing

Xyrem may impair respiratory drive, especially in patients with compromised respiratory function. In overdoses, life-threatening respiratory depression has been reported [see *Overdosage (10)*].

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.