

A randomized trial of abiraterone acetate (AA) administered with 1 of 4 glucocorticoid (GC) regimens in metastatic castration-resistant prostate cancer (mCRPC) patients (pts).

Subcategory:

Prostate Cancer - Advanced Disease

Category:

Genitourinary Cancer

Meeting:

2016 Genitourinary Cancers Symposium

Session Type and Session Title:

Poster Session A: Prostate Cancer

Abstract Number:

261

Poster Board Number:

Poster Session A Board #L10

Citation:

J Clin Oncol 34, 2016 (suppl 2S; abstr 261)

Author(s):

Gerhardt Attard, Axel S. Merseburger, Cora N. Sternberg, Linda Cerbone, Federica Recine, Robert J. Jones, Susan Feyerabend, Alfredo Berruti, Steven Joniau, Peter Schatteaman, Lajos Geczi, Peter Tenke, Patrick Werbrout, Florence Lefresne, Florence Nave Shelby, Marjolein Lahaye, Corinna Pick, Bertrand F. Tombal; The Institute of Cancer Research and The Royal Marsden Hospital, London, United Kingdom; Department of Urology, University of Lübeck, Lübeck, Germany; Department of Medical Oncology, San Camillo and Forlanini Hospital, Rome, Italy; University of Glasgow, Beatson West of Scotland Cancer Centre, Glasgow, United Kingdom; Studienpraxis Urologie, Nürtingen, Germany; Medical Oncology, University of Brescia, Spedali Civili Hospital, Brescia, Italy; University Hospitals Leuven, Leuven, Belgium; Onze-Lieve-Vrouweziekenhuis, Aalst, Belgium; Chemotherapy C and Clinical Pharmacology Department, National Institute of Oncology, Budapest, Hungary; Jahn Ferenc Dél-Pesti Kórház, Urology Department, Budapest, Hungary; AZ Groeninge, Kortrijk, Belgium; Janssen-Cilag, Paris, France; Janssen-Cilag, Tilburg, Netherlands; Janssen-Cilag, Neuss, Germany; Institut de Recherche Expérimental et Clinique, Université Catholique de Louvain, Brussels, Belgium

Abstract Disclosures**Abstract:**

Background: AA is approved for mCRPC, coadministered with prednisone (P) (5 mg BID) to prevent adverse events (AEs) associated with mineralocorticoid excess (ME). Lower GC doses had not previously been formally evaluated in combination with AA. **Methods:** This was an open-label, multicenter, phase 2 trial (NCT01867710) of asymptomatic chemotherapy-naïve mCRPC pts randomized 1:1:1:1 to AA (1000 mg QD) plus P 5 mg BID or P 5 mg QD or P 2.5 mg BID or dexamethasone (DEX) 0.5 mg QD. Pts who had previously received GC or ketoconazole were excluded. The primary end point was no ME (ME of pts experiencing neither hypokalemia nor hypertension during the first 24 weeks of treatment). Secondary end points included additional safety, as well as response rate in the first 24 weeks, defined as a decline in prostate-specific antigen (PSA) \geq 50% confirmed after 4 weeks. **Results:** 164 pts were randomized; 133 (81.6%) completed 24 weeks' treatment. Median age: 70 years. Table 1 shows the rates of ME, hypertension, hypokalemia and PSA response. Changes in HbA1c values were minimal and observed in 16 (10.7%) pts. **Conclusions:** These data suggest that P 5 mg BID, which is approved in combination with AA, and DEX 0.5 mg QD, are effective in preventing ME-associated AEs, and that P 2.5 mg BID and P 5 mg QD can be safely used with appropriate monitoring. The suggestion of a higher PSA response rate with DEX 0.5 mg QD arm warrants further validation. Clinical trial information: **NCT01867710**

	P 5 mg BID	P 5 mg QD	P 2.5 mg BID	DEX 0.5 mg QD
	n = 41	n = 41	n = 40	n = 42
No ME^a, n/N (%)	27/34 (79.4)	17/37 (45.9)	22/35 (62.9)	28/37 (75.7)
Hypokalemia^a, n/N (%)				
Overall	3/41 (7.3)	6/41 (14.6)	3/39 (7.7)	4/42 (9.5)
Grade 3	0	2/41 (4.9)	0	0
Hypertension^a, n/N (%)				
Overall	6/41 (14.6)	15/41 (36.6)	12/39 (30.8)	5/42 (11.9)
Grade 3	3/41 (7.3)	6/41 (14.6)	5/39 (12.8)	1/42 (2.4)
\geq 50% PSA response at Week 12, n/N (%)	20/35 (57.1)	24/34 (70.6)	17/37 (45.9)	31/39 (79.5)
No change in HbA1c, n/N (%)^b	28/38 (73.7)	34/36 (94.4)	37/37 (100)	35/39 (89.7)

N = Evaluable pts. ^aAccording to the National Cancer Institute Common Terminology Criteria for Adverse Events V4.0. No grade 4 occurred. ^bChange at end point versus baseline.

Abstracts by Gerhardt Attard:

Amerigen Exhibit 1128

Loco-regional treatment (LRT) for M1 at diagnosis prostate cancer (PCa) patients (pts) and impact on overall survival (OS): A retrospective analysis.

Meeting: 2016 Genitourinary Cancers Symposium | Abstract No: 280 | First Author: Diletta Bianchini

Category: Genitourinary Cancer - Prostate Cancer - Advanced Disease


PSA levels after dexamethasone withdrawal (DW) in castration resistant prostate cancer (CRPC).

Meeting: 2016 Genitourinary Cancers Symposium | Abstract No: 278 | First Author: Zafeiris Zafeiriou

Category: Genitourinary Cancer - Prostate Cancer - Advanced Disease

More

ASSOCIATED POSTER



Meeting: 2016 Genitourinary Cancers Symposium

Presenter: Gerhardt Attard

[View Poster](#)