

- [54] **CLONED HIGH SIGNAL STRENGTH PROMOTERS**
- [75] Inventors: **Hermann G. Bujard**, Heidelberg, Fed. Rep. of Germany; **Annie C. Y. Chang**, Palo Alto; **Stanley N. Cohen**, Portola Valley, both of Calif.
- [73] Assignee: **The Board of Trustees of the Leland Stanford Jr. University**, Stanford, Calif.
- [21] Appl. No.: **265,276**
- [22] Filed: **May 20, 1981**
- [51] Int. Cl.³ **C12Q 1/68; C12Q 1/02; C12P 21/00; C12P 19/34; C12N 15/00**
- [52] U.S. Cl. **435/6; 435/29; 435/68; 435/172.3; 435/91; 536/27; 935/6; 935/36; 935/37; 935/41; 935/76**
- [58] Field of Search **435/29, 68, 6, 172, 435/240, 253, 254, 257, 258, 317, 91; 536/27, 28, 29**

[56] **References Cited**

PUBLICATIONS

Stuber et al., PNAS, 78(1), 167-171, (Jan. 1981).


Casadaban et al., J. Mol. Biol., 138, 179-207, (1980).
 West et al., Gene, 9, 175-193, (1980).
 Gabain et al., PNAS, 76(1), 189-193, (1979).
 Stüber et al., Molec. Gen. Genet., 166, 141-149, (1978).
 Stüber et al., PNAS USA, 78(1), 167-171, (1981).

Primary Examiner—Esther M. Kepplinger
Attorney, Agent, or Firm—Bertram I. Rowland

[57] **ABSTRACT**

Method for preparing high signal strength promoters and terminators and DNA compositions employing such promoters and terminators. T5 phage is cleaved to provide for DNA sequences having intact promoters. These promoters are inserted into vectors separated from a balanced terminator by a gene of interest and the terminator is desirably followed by a marker allowing for selection of transformants. High efficiencies in transcription of DNA can be achieved with the highly active T5 promoters. The promoters and terminators are used in hybrid DNA for efficient expression of structural genes and transcription to provide RNA sequences.

15 Claims, 1 Drawing Figure



CLONED HIGH SIGNAL STRENGTH PROMOTERS

The government has rights in the invention pursuant to Grant Nos. AI 08619 and GM 27241 awarded by the National Institute of Health.

The research was supported in part by a grant from the Deutsche Forschungsgemeinschaft.

BACKGROUND OF THE INVENTION

1. Field of the Invention

Having established the feasibility of producing a wide variety of naturally occurring and synthetic polypeptides by means of hybrid DNA technology, there are continuing and extensive efforts to provide for more efficient and economic methods for producing the polypeptides. In developing a process for the commercial production of polypeptides, many factors will be involved in optimizing the economic and efficient production of the polypeptides. Included among these factors are regulatory signals, which are DNA sequences involved with the regulation of replication, transcription and translation.

One area of interest is at the level of transcription. Transcription involves the enzyme RNA polymerase. The RNA polymerase binds to a site called a promoter. It has been observed that promoters vary in their activity, as evidenced by the number of initiations of RNA per unit time or the strength of binding of the enzyme to the promoter site. The promoter may have one or more sequences that bind, which may or may not be contiguous. The more active promoters are referred to as strong promoters.

It was found that when introducing a strong promoter into a vector and employing the resulting plasmid for transformation, one could not select transformants based on expression of markers which allowed for selection. Therefore, cloning of the strong promoters was not feasible. It is therefore desirable that methods be provided which would allow for the screening of strong promoters and terminators and their subsequent cloning to be used in conjunction with the replication, transcription and translation of the genes for production of DNA, RNA, and polypeptides.

2. Description of the Prior Art

Promoters from bacterial and viral sources have been cloned in *E. coli*, and their signal strength in vitro has been studied using expression from distal promoterless sequences encoding β -galactosidase or other proteins (Casadaban and Cohen (1980) *J. Mol. Biol.* 138, 179-207; West and Rodriguez (1980) *Gene* 9, 175-193). Attempts to clone small DNA fragments carrying the strong promoters of bacteriophage T5 have been unsuccessful (v. Gabain and Bujard (1979) *PNAS USA* 76, 189, 193). Fragments of T5 DNA having both a strong promoter and a strong termination signal have been cloned. (Breunig (1979) Dissertation (Universitat Heidelberg, Heidelberg, Germany)) Analysis has shown that transcriptional regions of several *E. coli* plasmids are organized in units where initiation and termination signals, are balanced. (Stuber and Bujard (1981) *PNAS USA* 78: 167-171) P₂₅ and P₂₆ promoters of the T5 bacteriophage are reported as among the most efficient RNA polymerase binding sequences. (Stuber et al (1978) *Mol. gen. Genet.* 166 141-149; Niemann (1981) *Diplomarbeit* (Universitat Heidelberg, Heidelberg, Germany)).

SUMMARY OF THE INVENTION

Methods for cloning, sequencing and using strong promoters and terminators are provided, as well as compositions resulting from the methods. By cleaving T5 phage and selecting fragments specifically binding to RNA polymerase, fragments containing promoters are isolated. A vector is constructed having a strong promoter, followed by a DNA sequence of interest, optionally followed by one or more translational stop codons in one or more reading frames, followed by a balanced terminator, followed by a marker allowing for selection of transformants. Upon introducing the resulting plasmid into a microorganism host, efficient transcription of the gene is obtained with substantially lesser expression of the marker as compared to the gene. The level of expression of the marker permits selection of transformants having the above described construct. The construct or regulatory portions thereof are used for efficient transcription of RNA or gene expression.

BRIEF DESCRIPTION OF THE DRAWING

FIG. 1 is a flow chart of the preparation of a plasmid for cloning strong promoters and terminators.

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Methods and compositions are provided for preparing and cloning strong promoter and terminator regulatory signals and utilization of the strong regulatory sequences in the transcription and expression of genes of interest.

Particularly, T5 phage promoters are isolated, cloned in conjunction with a strong terminator, and appropriate vectors developed for insertion of DNA sequences of interest, usually structural genes, to provide for high and efficient transcription and/or expression of the sequence.

The compositions of this invention are characterized as having in the downstream direction of transcription the following units: a strong T5 phage promoter; optionally a structural gene which may be a marker; a balancing terminator; and optionally a marker allowing for selection of transformants containing the construct, which marker has a relatively low level of expression in comparison with the amount of RNA polymerase initiation at the promoter. In the absence of a promoter in the construct, the construct can be used for the cloning and characterization of promoters of different strengths.

In referring to strong promoters, it is intended that the binding affinity of RNA polymerase is stronger than the commonly employed promoters such as lac and trp and at least comparable to and normally greater than the combination of lac and trp promoters. For the most part, the strongest promoters among prokaryotes are the T5 phage promoters and these will be employed as exemplary of naturally occurring or synthetic strong promoters. It is to be understood, that other prokaryotic and eukaryotic promoters, either naturally occurring or synthetic, could find application in the subject invention.

The compositions of this invention will include linear segments for insertion of DNA having the strong regulatory signal sequences (i.e., the promoter and terminator) adjacent opposite ends of the linear segment and plasmids formed by introducing a DNA sequence from a source other than the source of the promoter as a bridge between the strong regulatory signal sequences.

The termini may be blunt or staggered ended, having the same or different termini to allow for directed positioning of inserted sequences.

The compositions which are employed as already indicated have a promoter, a balanced terminator, and desirably a marker which are in the direction of transcription when the signal sequences are joined. The terminator is balanced, so that a small but useful amount of expression of the marker can occur. Normally, in most situations, a gene will bridge the promoter and terminator to provide circular DNA. The gene may be a non-structural gene or a structural gene providing RNA e.g. ribosomal or messenger, or providing a poly(amino acid).

Desirably, the gene is followed by one or a plurality of translational stop codons e.g. oop or nonsense codons, or preferably a plurality, usually up to about six, more usually from about two to five, where there is at least one stop codon in each reading frame. The stop codons aid in the efficiency of termination, both at the level of transcription and expression.

Next in the direction of transcription is the terminator sequence. The terminator sequence is balanced with the promoter in the sense that the marker is transcribed not more than about one-fourth of the times that a gene upstream from the terminator is transcribed, where both genes are under the control of the same promoter. Usually, it will be desirable that a sufficient amount of the marker is transcribed to allow for selection of transformants. In view of the fact that the marker is transcribed from a strong promoter, it will usually be sufficient that the marker is transcribed in relation to the transcription of the gene intermediate the promoter and terminator in only about 1 to 20, usually only 5 to 15 number % of the gene. The strength of the promoter is reflected in the level of expression of the marker which is transcribed from the same promoter as the gene whose expression is desired.

In addition to the DNA sequences indicated above, there will normally be other regulatory signals necessary for expression involved with the DNA sequence, such as translational start and stop sites. In addition to the foregoing regulatory signals, other regulatory signals may be included, such as additional promoters, operators, initiators, catabolite activator protein binding sites, etc. Furthermore, the promoter and terminator may be separated by more than one gene, that is, a plurality of genes, including multimers and operons.

The above DNA sequence construct will have a replicating system or be cleaved and be inserted into a vector to provide a plasmid. The vector is distinguished by having one or more DNA sequences which serve to insure stable replication of the plasmid and may also provide opportunities for high copy numbers of the plasmid in the microorganism host. The vectors may be derived from chromosomal or extrachromosomal sources. The sources include plasmids, viruses (phage), chromosomes, or the like. In addition, the vector or the essential portions thereof may be prepared synthetically.

The plasmids may then be used for transformation of an appropriate microorganism host. Methods of introducing DNA into an appropriate host are well known. Illustrative of such methods, but not exhaustive of such methods, are transformation e.g. calcium shock, transfection, and conjugation. Descriptions of these methods may be found in Genetic Engineering, ed. Setlaw and Hollaender, Vol. 1, Plenum Press, New York and Lon-

don, 1979; Molecular Cloning of Recombinant DNA, ed. Scott and Werner, Vol. 13, Academic Press, Inc. New York, 1973, and references cited therein.

In order to allow for flexibility in preparing the construct and self-replicating sequence or plasmid containing the construct restriction sites should be present to allow for unique insertions and isolation of the various elements. The restriction sites may be naturally present, introduced by linkers, result by partial sequential nucleotide removal from a chain using an exonuclease, or the like. Desirably, the restriction sites will provide for different ends to permit only the proper orientation of the inserted fragment.

A wide variety of structural genes are of interest for production of proteins, including but not limited to proteins of physiological interest, proteins as chemicals, and enzymes which may be of direct interest or of interest in transforming another product, which may be proteinaceous or non-proteinaceous. The proteins may be prepared as a single unit or as individual subunits and then joined together in appropriate ways. Furthermore, as appropriate, the protein products may be modified by glycosylation, acylation with aliphatic acids, e.g. lipid acids, phosphorylation, sulfonation or the like. The different classes of proteins which may be prepared include protamines, histones, albumins globulins, scleroproteins, phosphoproteins, mucoproteins, chromoproteins, lipoproteins, nucleoproteins, and the remaining proteins which are unclassified.

The following is a representative list of proteins of interest.

Insulin; growth hormone; interferon e.g. leukocyte, lymphoblastoid, T-immune and fibroblast; β -lipotropin; β -endorphin; dynorphin; histocompatibility proteins; immunoglobulins e.g. IgA, IgD, IgE, IgG and IgM and fragments thereof; hemoglobin, somatomedins; lymphokines; growth factors e.g. epidermal, fibroblast, platelet-derived, multiplication stimulating and nerve; hematopoietic-stimulating factors e.g. erythropoietin, colony-stimulating, erythroid potentiating activity or burst-promoting activity and lymphopoietins; albumin and prealbumin;

Prealbumin
Albumin
 α_1 -Lipoprotein
 α_1 -Acid glycoprotein
 α_1 -Antitrypsin
 α_1 -Glycoprotein
Transcortin
4.6S-Postalbumin
Tryptophan-poor
 α_1 -glycoprotein
 α_1 - χ -Glycoprotein
Thyroxin-binding globulin
Inter- α -trypsin-inhibitor
Gc-globulin:
(Gc 1-1),
(Gc 2-1),
(Gc 2-2),
Haptoglobin:
(Hp 1-1),
(Hp 2-1),
(Hp 2-2),
Ceruloplasmin
Cholinesterase
 α_2 -Lipoprotein(s)
 α_2 -Macroglobulin
 α_2 -HS-Glycoprotein

Zn- α_2 -glycoprotein
 α_2 -Neuramino-glycoprotein
 Erythropoietin
 β -lipoprotein
 Transferrin
 Hemopexin
 Fibrinogen
 Plasminogen
 α_2 -glycoprotein I
 α_2 -glycoprotein II
 Immunoglobulin G
 (IgG) or γ G-globulin
 Mol. formula:
 $\gamma_2\kappa_2$ or $\gamma_2\lambda_2$
 Immunoglobulin A (IgA) or γ A-globulin
 Mol. formula:
 $(\alpha_2\kappa_2)^n$ or $(\alpha_2\lambda_2)^n$
 Immunoglobulin M (IgM) or γ M-globulin
 Mol. formula:
 $(\mu_2\kappa_2)^5$ or $(\mu_2\lambda_2)^5$
 Immunoglobulin D (IgD) or γ D-Globulin (γ D)
 Mol. formula:
 $(\delta_2\kappa_2)$ or $(\delta_2\lambda_2)$
 Immunoglobulin E (IgE) or γ E-Globulin (γ E)
 Mol. formula:
 $(\epsilon_2\kappa_2)$ or $(\epsilon_2\lambda_2)$
 Free light chains
 Complement factors:
 C'1
 C'1q
 C'1r
 C'1s
 C'2
 C'3
 β_1 A
 α_2 D
 C'4
 C'5
 C'6
 C'7
 C'8
 C'9.

Important protein hormones include:

Peptide and Protein Hormones

Parathyroid hormone: (parathormone)
 Thyrocalcitonin
 Insulin
 Glucagon
 Relaxin
 Erythropoietin
 Melanotropin: (Melanocyte-stimulating hormone; inter-
 medin)
 Somatotropin: (growth hormone)
 Corticotropin: (adrenocorticotropic hormone)
 Thyrotropin
 Follicle-stimulating hormone
 Leuteinizing hormone: (interstitial cell-simulating hor-
 mone)
 Luteomammotropin hormone: (Luteotropin, prolactin)
 Gonadotropin: (chorionic gonadotropin).

Tissue Hormones

Secretin
 Gastrin
 Angiotensin I and II
 Bradykinin
 Human placental lactogen

Peptide Hormones from the Neurohypophysis

Oxytocin
 Vasopressin
 5 Releasing factors (RF): CRF, LRF, TRF, Somatotro-
 pin-RF, GRF, FSHRF, PIF, MIF.

In addition to various non-enzymatic proteins of
 physiological interest, enzymes can also be produced as
 10 an end product or for intracellular transformation of a
 substrate present in the host or substrate introduced
 extracellularly, or for enzymatic transformation in vi-
 tro.

In accordance with the I.U.B. classification, the en-
 15 zymes fall into varying categories such as 1. oxidore-
 ductases; 2. transferases; 3. hydrolases; 4. lyases; 5. isom-
 erases; 6. ligases. Enzymes of particular interest will be
 hydrolases and oxidoreductases for use in commercial
 processing, for example, hydrolases for hydrolysing
 20 polysaccharides, lipids and polypeptides; oxidoreduc-
 tases for oxidation of alcohols and aldehydes, epoxida-
 tion, and the like.

The microorganism host may be bacteria, such as
 Escherichia, Bacillus, Aerobacter, Klebsiella, Proteus,
 Pseudomonas, Streptococcus, Staphylococcus, Clo-
 25 stridium, Mycobacterium, Streptomyces and Actino-
 myces; Fungi e.g. Gymnomycota, Dimastigomycota,
 Eumycota, Zygomycetes, Ascomycetes and Basidomy-
 cetes, such as, Candida, Aspergillus, Rhizobus, Micro-
 sporium, and Fonsecaea; Protozoa e.g. Mastigophora,
 30 Sarcodina, Sporozoa and Celiophora, such as, Trypano-
 soma, Codosiga, Protospongra and Entameba, and Alga
 e.g. Dinoflagellates, Euglenoids, and Diatoms.

Higher cells, e.g., mammalian, may also be employed
 35 as hosts, where viral, e.g., bovine papilloma virus or
 other DNA sequence is available which has plasmid-
 like activity.

Depending upon the nature of the host, a wide vari-
 ety of vectors may be employed. The vector will have
 40 an intact replicon and be capable of replication in the
 host. In addition, replicons can be developed which
 may have one or more other regulatory signals. Regula-
 tory signals can include temperature sensitive replicons,
 runaway-replication sequences, temperature sensitive
 45 operators, and the like. Various additional DNA se-
 quences may be present providing for restriction sites,
 markers, termination sequences, or the like. Desirably,
 the vector should be of a substantially different size
 from the construct to allow for excision of the construct
 50 and ease of separation of the construct from the vector
 by molecular weight separation techniques e.g. electro-
 phoresis and density gradient centrifugation.

The subject invention also provides for a technique
 for screening the strength of promoters and terminators,
 55 thus allowing for the determination of the use of a pro-
 moter or terminator in a particular application, where it
 may be desirable to have promoters or terminators or
 combinations thereof of varying strength. In screening
 promoters and/or terminators one employs a promoter
 60 or terminator of known activity. A DNA construct is
 made having the sequence described previously, where
 appropriate restriction sites are provided for introduc-
 ing the various elements. The construct provides in the
 direction of transcription the promoter, a first gene
 65 marker, optionally termination codons, such as non-
 sense codons and oop terminator, the terminator, and a
 second gene marker. This DNA construct is inserted
 into an appropriate vector. Where the promoter and

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.