

Disappearing and Reappearing Polymorphs. The Benzocaine:Picric Acid System

Jan-Olav Henck,^{*,†,||} Joel Bernstein,^{*,‡} Arkady Ellern,[‡] and Roland Boese[§]

Contribution from the Institut für Pharmakognosie der Universität Innsbruck Innrain 52, Josef-Moeller-Haus, 6020 Innsbruck, Austria, Department of Chemistry, Ben-Gurion University of the Negev, Beer Sheva, Israel 84105, and FB8/Anorganische Chemie, Universität-GH Essen, Universitätstrasse 5, 45117 Essen, Germany

Received June 12, 2000

Abstract: The low-melting polymorphic modification of the 1:1 complex of benzocaine (BC) and picric acid (PA) had earlier been reported to be an example of a “disappearing polymorph”. The BC:PA system has been reinvestigated by thermomicroscopy, calorimetry, solid-state NMR, and X-ray crystallography. The phase diagram has been derived, and robust procedures for the crystallization of the two 1:1 complexes, a hydrate of the 1:1 complex, and a 2:1 complex have been devised. The structures of all four phases have been determined and compared using graph set analysis to characterize the hydrogen-bonding patterns. It is shown that the thorough microscopic investigation of the thermal behavior, combined with calorimetric methods, can lead to the development of strategies to crystallize metastable polymorphic forms which may be difficult to obtain once their stable congeners have been obtained.

Introduction

Polymorphism, the ability of a substance to exist in several different crystal forms or modifications, is a frequently observed phenomenon in molecular compounds.¹ If a solid substance includes a solvent during crystallization, this structure is known as a pseudopolymorph, and in case of, e.g., water such crystal forms are generally called hydrates.² The polymorphic modifications of a compound are chemically identical but usually differ in their physical and chemical properties, such as density, vibrational spectra, and diffraction patterns.

Since a particular polymorph may have desirable properties, it may be useful to develop a robust method to obtain that polymorphic modification consistently and reproducibly. There are many documented cases of difficulties in obtaining crystals of a particular known modification.^{3,4} Such difficulties may have serious practical consequences: a material designed to give a particular structure may exhibit an undesired polymorphic one;⁵ a plant manufacturing a particular modification may have to close because it has been “poisoned” by an undesired polymorph;⁶ a commercial drug formulation may no longer be

available on the market due to the appearance of an undesired polymorphic modification.⁷ We recently documented a number of representative cases in which it was difficult to obtain a given polymorphic form even though previously it had often been obtained routinely over long periods of time—so-called “disappearing polymorphs”.³ In the conclusion to that review, we stated our belief “that once a particular polymorph has been obtained, it is always possible to obtain it again; it is only a matter of finding the right experimental conditions”. One of the challenges to those dealing with the preparation of solids is to be able to rationally develop those conditions and to control the polymorphic form obtained—in many instances to recover the disappeared polymorph. In this paper we present a case study of how such a challenge might be met.

One of the examples cited in the above-mentioned review was the case of benzocaine picrate (BC:PA). This binary system has been studied previously at least three separate times.^{8–10} In 1972, Nielsen and Borka⁸ reported the first preparation of the complex (now known as Mod. II) with a melting point of 130–132 °C. At the time, this modification was used as a pharmacopoeial standard for the identification of benzocaine (a topical anesthetic). Subsequently, the same authors prepared a high-melting modification (now known as Mod. I), mp 162–163 °C, whereupon they encountered severe difficulties in preparing and maintaining Mod. II for more than several hours before it transformed into Mod. I.¹¹ The low-melting modification is thermodynamically unstable at 20 °C.⁸

In 1974, Borka and Kuhnert-Brandstätter⁹ reinvestigated the system using thermomicroscopy and found four modifica-

(7) Chemburkar, S R ; Bauer, J ; Deming, K ; Spiwek, H ; Patel, K ; Morris, J ; Henry, R ; Spanton, S ; Dziki, W ; Porter, W ; Quick, J ; Bauer, P ; Donaubaer, J ; Narayanan, B A ; Soldani, M ; Riley, D ; McFarland, K *Org. Process Res. Dev.* **2000**, *4*, 413–417

(8) Nielsen, T K ; Borka, L *Acta Pharm. Suecica* **1972**, *9*, 503–505

(9) Borka, L ; Kuhnert-Brandstätter, M *Arch. Pharmaz.* **1974**, *307*, 377–384

(10) Togashi, A ; Matsunaga, Y *Bull. Chem. Soc. Jpn.* **1987**, *60*, 1171–1173

* To whom correspondence should be addressed

† Universität Innsbruck

‡ Ben-Gurion University of the Negev

§ Universität-GH Essen

|| Current address: Bayer AG, PH-PD PQCD1, Building E39, 51368 Leverkusen, Germany

(1) The April 1, 1998, release of the Cambridge Structural Database contains 181 309 entries, of which 5641 contain the qualifier “form”, 163 the qualifier “polymorph”, 172 the qualifier “modification”, and 146 the qualifier “mod”. Not all of these are entries for the structures of all polymorphs for a particular substance, but these numbers indicate that roughly 3% of the entries in the database are for polymorphic materials


(2) Byrn, S R *Solid State Chemistry of Drugs*, 2nd ed ; SSCI, Inc : West Lafayette, IN, 1999; pp 513–514

(3) Dunitz, J D ; Bernstein, J *Acc. Chem. Res.* **1995**, *28*, 193–200

(4) Webb, J ; Anderson, B *J. Chem. Educ.* **1978**, *55*, 644

(5) Aakeröy, C B ; Nieuwenhuysen, M ; Price, S L *J. Am. Chem. Soc.* **1998**, *120*, 8986–8993

(6) Anonymous *Ind. Eng. Chem.* **1953**, *11a*



tions of **BC:PA**. They further identified two modifications of a $(\text{BC})_2\text{:PA}$ complex as well as a monohydrate. Apparently unaware of both of these earlier works, Togashi and Matsunaga also reported a $(\text{BC})_2\text{:PA}$ complex and an additional charge-transfer complex between the two compounds, observed by using differential scanning calorimetry (DSC).¹⁰ No structural studies on any of the complexes were reported.

The aims of the work we report here were (1) to develop a general strategy and specific methods for controllably and reproducibly obtaining the “disappearing” modification II of **BC:PA**; (2) to develop methods for obtaining single crystals suitable for X-ray structure determination of as many as possible of the **BC:PA** species reported by earlier authors; and (3) to fully characterize the **BC:PA** phases by spectroscopic and structural techniques.

Experimental Strategy

Since the discovery of polymorphism in 1822,¹² one of the principal techniques for investigating the phenomenon has been optical microscopy and, in the latter part of the 20th century, thermomicroscopy.^{13–16} These methods are particularly useful in detecting the existence of polymorphs and polymorphic transitions quickly and easily. In the last few decades, more sophisticated analytical techniques have supplanted the two classic ones,¹⁷ but the net result of years of effort by a number of groups is that many polymorphic materials have been identified and characterized.^{16,18–20} Despite the success of these methods, very few of these findings have been utilized as the starting point for structural studies.²¹ The general problem

(11) The authors reported the disappearing phenomenon and their attempts to overcome it as follows:⁸ “As a matter of curiosity, it ought to be mentioned that once the stable modification was obtained, the metastable modification could no longer be isolated. It was first observed by one of us (TKN). This observation was later confirmed in a separate laboratory (LB). Further, it was found that after discarding all samples, washing the equipment and laboratory benches and waiting for 8–12 days, the low melting modification could be isolated again. This has now been repeated several times in our laboratories. Obviously the seeding effect during the formation of the primary crystals (or during the very procedure of determination of the melting point) is exceptionally strong.”

(12) Mitscherlich, E. *Ann. Chim. Phys.* **1822**, *19*, 350–419.

(13) Chamot, E. M.; Mason, C. W. *Handbook of Chemical Microscopy, Volume I, Principles and Use of Microscopes and Accessories Physical Methods for the Study of Chemical Problems*, 3rd ed.; John Wiley and Sons: New York, 1958.

(14) Kofler, L.; Kofler, A. *Thermo-MikroMethoden zur Kennzeichnung organischer Stoffe und Stoffgemische*; Wagner: Innsbruck, 1954.

(15) McCrone, W. C. *Fusion Methods in Chemical Microscopy*; Interscience: New York, 1956.

(16) (a) Kuhnert-Brandstätter, M. *Thermomicroscopy in the Analysis of Pharmaceuticals*; Pergamon Press: Oxford, 1971 Kuhnert-Brandstätter, M. In *Thermomicroscopy of Organic Compounds*; Svehla, G., Ed.; Comprehensive Analytical Chemistry, Vol. XVI; Elsevier: Amsterdam, 1982; pp 329–513.

(17) Threlfall, T. *Analyst* **1995**, *120*, 2435–2460.

(18) Burger, A.; Ramberger, R. *Mikrochim. Acta* **1979**, *II*, 259–271.

(19) Haleblan, J. K.; Borika, L. *Acta Pharm. Jugosl.* **1990**, *40*, 71–94.

(20) Borika, L. *Pharm. Acta, Helv.* **1991**, *66*, 16–22.

(21) For a more detailed account of the use and decline of the use of the

apparently lies in the difficulty of translating the microscopic observations to the growth of single crystals that can be used for structure determination. In this work, we adopted a strategy for crystal growth experiments aimed at obtaining crystals for X-ray determination of the various phases of **BC:PA** based on the observations made using optical and thermomicroscopy and on the appearance and disappearance of those phases. We believe that this approach can be widely utilized and thus can provide the means for greatly expanding our understanding of the structural basis for polymorphic behavior.

Results

The Phase Diagram. The phase diagram can provide very useful information and guidelines for the preparation of a particular polymorphic modification. Much of the data for the preparation of the phase diagram may be readily obtained from thermomicroscopic studies. In fact, for a two-component system, such as **BC:PA**, Kuhnert-Brandstätter pointed out that “Kofler’s contact method of thermal analysis offers the possibility of investigating organic two component systems qualitatively in the shortest possible time in the simplest possible manner.”^{16,22} More quantitative details may be added from DSC measurements.

Details on the preparation of the **BC:PA** contact sample and analysis are given in the Experimental Procedures section. Figure 1a shows the photomicrograph of the recrystallized contact preparation of **BC** and **PA** at 25 °C. The interference colors are due to the use of crossed polarizers. The pure compounds are at the extremities of the preparation, while in the region where the original compounds have merged, a number of different areas may be observed, due to the formation of different crystalline species combining the two components. Heating this preparation on the hot-stage microscope to a temperature of about 88 °C shows (Figure 1b) the eutectic melt of **BC** and the broad dark yellow crystals of a new compound. Due to the crossed polarizers, the isotropic melt appears black. As can be seen in Figure 1c, at about 120 °C on the left side of the preparation **BC** is melted, and on the right side the eutectic between **PA** and the remaining crystals of the **BC:PA** 1:1 complex melts. Figure 1d shows the situation which is observed at 122 °C. **PA** is almost melted, and in the middle of the preparation a eutectic melt appears. Thus, two chemically different kinds of complexes between **BC** and **PA** have been formed. The one on the right side is the 1:1 complex, while the small strip on the left side (the **BC** side) is a complex with composition $(\text{BC})_2\text{:PA}$.⁹ The former melts at 129 °C, while the latter shows a melting point at 124 °C.

The results obtained by means of thermomicroscopy and DSC experiments on different mixtures of benzocaine and picric acid led to the phase diagram (Figure 2). These investigations indicate the existence of two modifications of the 1:1 complex as well as of one modification of the 2:1 complex. The eutectic points of Mod. II of the 1:1 complex with benzocaine and picric acid are located at mole fractions of 0.07 (88 °C) and 0.85 (109 °C), respectively. The peritectic of Mod. II with the 2:1 complex appears at a mole fraction 0.33 and about 120 °C, and the eutectic between the 2:1 complex and benzocaine is located at 0.3 and at 110 °C.

Preparation of Single Crystals. The thermodynamic phase relationship between the two forms of **BC:PA** may be

(22) For a summary of the contact method, see also: Emons, H.-H.; Keune, H.; Seyfarth, H.-H. In *Chemical Microscopy*; Svehla, G., Ed.




Figure 1. Photographs of the microscope slide Kofler preparations showing the various phases of the **BC:PA** binary system: (a) 25 °C, (b) 87 °C, (c) 120 °C, and (d) 123 °C (see text for explanation and discussion of details of the phenomena observed here).

enantiotropic or monotropic.¹⁶ In the former case, the choice of crystallization conditions for one of the two polymorphs would be more facile, because thermodynamically controlled crystallization conditions could be chosen at temperatures below the thermodynamic transition point. To determine the monotropism/enantiotropism, we crystallized microcrystalline **BC:PA** Mod. II by means of thermomicroscopy and carried out DSC measurements on the resulting solid. The DSC trace shows an exothermic transition of Mod. II into Mod. I. We could then apply the empirical heat-of-transition rule²³ to conclude that these two crystal forms are monotropically related (Table 1). This means that Mod. I is the thermodynamically stable crystal form from absolute zero up to its melting point.

Since the lower melting Mod. II is thermodynamically less stable at all temperatures below the melting point, “kinetic” conditions must be created to attempt to crystallize this form. Therefore, a high-temperature (80 °C) crystallization was attempted, with water as the solvent, yielding single crystals up to 1 mm in maximum dimension. Since the prior work of

(23) According to the “heat-of-transition rule”,¹⁸ if the phase transition is exothermic, then there is no transition point below the experimentally observed transition temperature. This is generally observed if two transitions

Nielsen and Borcka⁸ indicated that seeds of the stable form must be excluded, we obtained the less stable form prior to carrying out crystallization experiments aimed at obtaining the more stable one.

As Mod. I is the thermodynamically preferred form, an “equilibrium” crystallization is preferred over a “kinetic” one. Nielsen and Borcka⁸ obtained Mod. I by “excessive drying of [Mod. II] at 105 °C”, conditions which are not particularly conducive to an “equilibrium” situation, nor to the growth of single crystals. Also, the existence of the hydrate (determined in the thermomicroscopic studies) suggests that the presence of water might be problematic in attempting to obtain nonhydrated phases. Hence, we resorted to a nonaqueous gel-diffusion crystallization,²⁴ using Sephadex as the gel medium, and obtained large single crystals (1–2 mm maximum dimension) after 3 days at 20 °C.

The rather drastic drying procedure described by Nielsen and Borcka⁸ to obtain Mod. I, Borcka and Kuhnert-Brandstätter’s report,⁹ and our own observations on the hot-stage microscope indicated the existence of a hydrate of **BC:PA**. To obtain single crystals of this material, we attempted a crystallization from a




Figure 2. Phase diagram of the BC:PA system.

Table 1. Physicochemical Data of BC:PA Mod. I, BC:PA Mod. II, and (BC)₂:PA

	BC:PA Mod. I	BC:PA Mod. II	(BC) ₂ :PA
mp (°C), DSConset therm microscopy	161	129	119
enthalpy of fusion (kJ mol ⁻¹)	63	56	120
enthalpy of transition (kJ mol ⁻¹)		-9	
		→ Mod. I	
entropy of fusion ^a (J mol ⁻¹ K ⁻¹)	145	118	

^a Calculated by $\Delta S_f = \Delta H_f / \text{mp}^{-1}$.





Figure 3. Line drawing representation of the morphologies of the four different crystalline materials obtained, as labeled.



saturated water solution in a sealed virgin flask (to prevent the unintentional incursion of seeds of any of the other forms) at 20 °C; needle-shaped crystals (up to 4 mm long) appeared after 48 h.

Crystals of the 2:1 complex were obtained by slow evaporation (ca. 4 weeks) of a 1:1 mixture of the components in 2-propanol at 4 °C. Admittedly, this was not an experiment designed to obtain the 2:1 complex, but the ample evidence to indicate its existence increased our interest in examining all the crystals obtained and motivated our attempts to obtain the crystals, using conditions different from those used for the other three species.

The crystals of all four complexes were yellow but were easily distinguished by their morphology (Figure 3) and, very clearly, by their melting points.

Crystal data for the four structures are given in Table 2; details of the structure determination are given in the Experimental Procedures section.

Crystal Structures. (a) General Features. All four structures are ionic picrates, the acidic hydrogen of the hydroxyl group on picric acid having been transferred to an amino group on benzocaine. The atomic numbering, given in a the ORTEP diagram (for (BC)₂:PA), is identical for all four structures

Table 2. Summary of Crystallographic Data for the Four Compounds Studied

	BC:PA Mod. I	BC:PA Mod. II	(BC) ₂ :PA	BC:PA·H ₂ O
formula	C ₁₅ H ₁₄ N ₄ O ₉	C ₁₅ H ₁₄ N ₄ O ₉	C ₂₄ H ₂₅ N ₅ O ₁₁	C ₁₅ H ₁₆ N ₄ O ₁₀
formula weight	394.3	394.3	559.5	412.3
crystal system	monoclinic	triclinic	triclinic	monoclinic
space group	<i>P</i> 2 ₁ / <i>n</i>	<i>P</i> 1	<i>P</i> 1	<i>P</i> 2 ₁
color of crystal	yellow	yellow	yellow	yellow
<i>Z</i>	4	2	2	2
<i>a</i> (Å)	7.149(3)	7.304(3)	8.190(2)	4.1478(8)
<i>b</i> (Å)	12.240(7)	8.231(2)	8.857(3)	12.732(3)
<i>c</i> (Å)	19.594(9)	15.361(5)	19.747(5)	17.518(4)
α (°)	90	99.91(2)	78.59(2)	90
β (°)	96.77(3)	99.53(2)	85.13(2)	93.84(3)
γ (°)	90	106.73(3)	67.93(2)	90
<i>R</i> -factor (%)	5.1	3.3	4.0	3.8
GOF (on <i>F</i> ²)	0.988	0.969	0.908	0.980




Figure 5. Packing diagrams the four crystal structures. For ease of comparison, in all cases the view is chosen on the best plane of the shaded picrate anion as a reference, with the C–O vector pointing down. Some slight rotational adjustments of the views have been made to facilitate better viewing of the packing and the hydrogen bonds. Carbon atoms are not explicitly drawn. Oxygens are represented by empty circles, and nitrogens are represented by filled circles. Hydrogens have been eliminated for clarity except where noted. The unit cell axes are marked as X, Y, Z, while hydrogen bonds are identified by lowercase letters (see text). (a) BC:PA Mod. I, (b) BC:PA Mod. II, (c) BC:PA·H₂O (hydrogens are included on the water molecule and on N1 of BC), and (d) (BC)₂:PA (hydrogens on N1 and N1' are included).

Table 3. Rotation Angles (deg) of Nitro Groups in PA in the Four Crystal Structures

angle about C–N bond	BC:PA Mod. I	BC:PA Mod. II	BC:PA·H ₂ O	(BC) ₂ :PA
C11–N2	–176.0	–144.4	–168.5	–152.8
C13–N3	4.7	11.3	8.2	0
C15–N4	–37.9	147.4	19.2	–49.4

(Figure 4). There are no unusual bond lengths or angles. Only two of the torsion angle values for the ethyl ester chain of the BC moiety suggest some conformational variation, namely that for C5–C6–C7–O1 of BC:PA Mod. I (157.6°) and that for C7–O1–C8–C9 of BC:PA Mod. II (161.0°) compared to the others, which are all in the range 170–180°. These conformational differences were not investigated further.

The rotations of the nitro groups in PA are summarized in Table 3. That para to the phenolate oxygen is essentially coplanar with the benzene ring in all cases. The ortho nitro groups are rotated out of the phenyl plane, those about C11–N2 all in the same sense but to varying degrees, while those about C15–N4 are also rotated to approximately the same degree, but the rotation in the hydrate is opposite in sense from the other three instances.

BC:PA Mod. I in (Figure 5a) may be described as mixed

BC⁺ and PA[–] within a stack make an angle of 2.3° with each other, the distance between centers of rings being 4.04 Å. BC:PA Mod. II (Figure 5b) also crystallizes in mixed π – π stacks of \cdots BC⁺ \cdots PA[–] \cdots , again essentially along [100]. The mode of overlap along the stack is such that the four atom chain of the ester group is nearly parallel to the long (O–C \cdots p–NO₂) axis of the molecule, while in Mod. I the chain of the ester group is nearly perpendicular to that axis. The angle between the phenyl rings (7.6°) is slightly larger, but the distance between centers is smaller (3.70 Å), indicating a greater degree of overlap between neighboring molecules in Mod. II.

In contrast to the two anhydrous complexes, in BC:PA·H₂O (Figure 5c) the cations and anions are arranged in segregated stacks \cdots BC⁺ \cdots BC⁺ \cdots and \cdots PA[–] \cdots PA[–] \cdots . The perpendicular distance between planes in the BC⁺ stack is 3.73 Å, while that in the PA[–] stack is 3.45 Å. Sheets of cations or anions are generated by the screw operation along the *b* axis, and in these sheets cations and anions alternate along the *c* axis. The angle between the phenyl rings of BC⁺ and PA[–] is 35°. There are sheets of cations and sheets of anions, both parallel to the stacking direction, so that the structure is built up of parallel sheets of cations and anions, in a manner similar to zinc blende.

Explore Litigation Insights

Docket Alarm provides insights to develop a more informed litigation strategy and the peace of mind of knowing you're on top of things.

Real-Time Litigation Alerts



Keep your litigation team up-to-date with **real-time alerts** and advanced team management tools built for the enterprise, all while greatly reducing PACER spend.

Our comprehensive service means we can handle Federal, State, and Administrative courts across the country.

Advanced Docket Research



With over 230 million records, Docket Alarm's cloud-native docket research platform finds what other services can't. Coverage includes Federal, State, plus PTAB, TTAB, ITC and NLRB decisions, all in one place.

Identify arguments that have been successful in the past with full text, pinpoint searching. Link to case law cited within any court document via Fastcase.

Analytics At Your Fingertips



Learn what happened the last time a particular judge, opposing counsel or company faced cases similar to yours.

Advanced out-of-the-box PTAB and TTAB analytics are always at your fingertips.

API

Docket Alarm offers a powerful API (application programming interface) to developers that want to integrate case filings into their apps.

LAW FIRMS

Build custom dashboards for your attorneys and clients with live data direct from the court.

Automate many repetitive legal tasks like conflict checks, document management, and marketing.

FINANCIAL INSTITUTIONS

Litigation and bankruptcy checks for companies and debtors.

E-DISCOVERY AND LEGAL VENDORS

Sync your system to PACER to automate legal marketing.